


1 What is a Graph

Graphs present a key concept of discrete mathematics. Informally, a graph consists of

- *vertices*, sometimes called nodes (“dots”),
- and *edges* (“arcs”) between pairs of vertices.



Importantly, graphs are **easy to draw** and understand from a picture, and **easy to process** by computer programs, too. □

Brief outline of this lecture

- What is a graph: definition of a graph and its basic terms, examples and trivial classes of graphs.
- Vertex degrees, degree sequence of a graph (score).
- Subgraphs and isomorphism, recognizing (non-)isomorphic graphs.
- Directed graphs and multigraphs.

1.1 Defining a graph

Definition 1.1. Graph (actually, a *simple undirected* graph) is a pair $G = (V, E)$ where V is the *vertex* set and E is the *edge* set – a subset of pairs of vertices.

Notation: An edge between vertices u and v is denoted by $\{u, v\}$, or shortly uv .

The vertices joined by an edge are *adjacent* or *neighbours*.


The vertex set of a (known) graph G is referred to as $V(G)$, the edge set as $E(G)$. □

Fact: A graph is, in algebra terms, a symmetric irreflexive binary relation. □

Remark: How can we describe a graph?

Either by listing the vertices and edges, or with a nice picture. . .

$$V = \{1, 2, 3, 4\}, \quad E = \left\{ \{1, 2\}, \{1, 3\}, \{1, 4\}, \{3, 4\} \right\}$$




□

Which one do you like more?


Basic graph classes

It is a custom to refer to some basic special graphs by descriptive names such as the following.

The cycle of length n has $n \geq 3$ vertices joined in a cyclic fashion with n edges:




The path of length n has $n + 1$ vertices joined consecutively with n edges:




The complete graph on $n \geq 1$ vertices has n vertices, all pairs forming edges (i.e. $\binom{n}{2}$ edges altogether):

K_n



The complete bipartite graph on $m \geq 1$ plus $n \geq 1$ vertices has $m + n$ vertices in two parts, and the edges join all the $m \cdot n$ pairs coming from different parts:



$K_{m,n}$



1.2 Vertex degrees in a graph

Definition 1.2. The **degree** of a vertex v in a graph G , denoted by $d_G(v)$, equals the number of edges of G incident with v . An edge e is **incident** with a vertex v if v is one of the ends of e . \square

In this example, the degrees are written at the vertices.



Definition: A graph is **d -regular** if all its vertices have the same degree d .

Notation: The **maximum** degree in a gr. G is denoted by $\Delta(G)$ and **minimum** by $\delta(G)$.

Theorem 1.3. *The sum of the degrees of all vertices of a graph is always even, equal to twice the number of edges.* \square

Proof. When summing the degrees, every edge has two ends and hence it is counted twice. \square

The degree sequence

Definition: The *degree sequence* (called also the score) of a graph G is the collection of degrees of the vertices of G , written in a sequence of natural numbers which is (usually) sorted as nondecreasing or nonincreasing.

In abstract graphs, their vertices usually have no names, and so we have to sort their degree sequence somehow. The particular custom is not important. \square

Just to quickly ask, why the sequence 1, 2, 3, 4, 5, 6 cannot be a degree sequence of a graph? \square
(Is the sum even? No...)

And what about the sequence 1, 2, 3, 4, 5, 6, 7? \square

Theorem 1.4. *Let $d_1 \leq d_2 \leq \dots \leq d_n$ be a sequence of natural numbers. There exists a simple graph on n vertices having a degree sequence*

$$d_1, d_2, \dots, d_n$$

if, and only if, there exists a simple graph on $n - 1$ vertices having the degree sequence

$$d_1, d_2, \dots, d_{n-d_n-1}, d_{n-d_n} - 1, \dots, d_{n-2} - 1, d_{n-1} - 1.$$

Example 1.5. *Is there a simple graph with degree sequence*

$(1, 1, 1, 2, 3, 4)$?

Using Theorem 1.4 we modify the sequence to $(1, 0, 0, 1, 2)$, \square

then sort again as $(0, 0, 1, 1, 2)$,

and continue analogously with the next step, arriving at the seq. $(0, 0, 0, 0)$. \square

A graph with the last degree sequence clearly exists, and so by the equivalence claim in Theorem 1.4, our graph exists as well. \square

How can we construct such a graph? See...



$(1, 1, 1, 1, 2, 3, 4, 6, 7)$? \square

The first step analogously translates to $(1, 0, 0, 0, 1, 2, 3, 5)$,

sorted as $(0, 0, 0, 1, 1, 2, 3, 5)$. \square

One more step and we get $(0, 0, -1, 0, 0, 1, 2)$. What does it mean? \square

Since the degrees of a graph cannot be negative, a graph with the last sequence does not exist, and neither does the original one!



\square

1.3 Subgraphs and Isomorphism

Definition: A *subgraph* of a graph G is any graph H on a subset of vertices $V(H) \subseteq V(G)$, such that the edges of H form a subset of the edges of G and have both ends in $V(H)$.

We write $H \subseteq G$ as for the set inclusion. \square

Why, on the left hand side, the red subsets do not form a subgraph?



\square


On the right hand side, we have got a well-formed subgraph in green colour. \square

Definition: An *induced subgraph* is a subgraph $H \subseteq G$ such that all edges of G between pairs of vertices from $V(H)$ are included in H .

Definition 1.6. An **isomorphism** between graphs G and H is a bijective (*one to one*) mapping $f : V(G) \rightarrow V(H)$ such that, for every pair of vertices $u, v \in V(G)$, it holds that $\{u, v\}$ is an edge of G if and only if $\{f(u), f(v)\}$ is an edge of H .

Graphs G and H are **isomorphic** if there exists an isomorphism between them. We write $G \simeq H$. \square



Fact: Isomorphic graphs have the same numbers of vertices and of edges.



Which two of the graphs are isomorphic and why the third one is not? \square



Fact: If a bijection f is an isomorphism, then f must map the vertices of same degrees, i.e. $d_G(v) = d_H(f(v))$. The converse is not sufficient, however!

Example 1.7. Are the following two graphs isomorphic?



We shall first compare the numbers of vertices and of edges. (Agree.) □ Then we compare their degree sequences. (Again, they agree $2, 2, 2, 2, 3, 3$.) □ This means we have found no easy distinction, and the graphs **might** (or may not!) be isomorphic. We have to start looking for all possible bijections between them. □

In this particular case, it helps to notice that the two degree-3 vertices on the left are isomorphic, and so we may choose any one of them to map it to the leftmost vertex of the second graph. □ Numbering the vertices by $1, 2, 3, 4, 5, 6$, we can construct our mapping (already have $1 \rightarrow 1'$) as follows, see on the picture below.



□

Theorem 1.8. *The relation “to be isomorphic” \simeq is an equivalence on the class of all graphs.*

Proof. We easily show that \simeq is reflexive, symmetric, and transitive. \square \square

The important corollary of this claim is that the class of all graphs is partitioned into the *isomorphism classes*.



Hence when we speak about a “graph”, we (usually) actually mean its whole isomorphism class. A particular presentation of the graph does not matter.

Additional graph notation



Notation: Consider an arbitrary graph G .

- A subgraph $H \subseteq G$ isomorphic to a cycle is called a *cycle in G* .
- Specially, a *triangle* in G is a subgraph isomorphic to the cycle of length 3. \square
- A subgraph $H \subseteq G$ isomorphic to a path is called a *path in G* . \square
- A subgraph $H \subseteq G$ isomorphic to a complete graph is called a *clique in G* .
- A vertex subset $X \subseteq V(G)$ such that no pair from X forms an edge of G is called an *independent set X in G* . \square
- An induced subgraph $H \subseteq G$ isomorphic to a cycle is called an *induced cycle in G* . Analogously an induced path...

What subgraphs and induced subgraphs can you find in the following graphs?



Example 1.9. *Are the following two graphs isomorphic?*



We start analogously to previous Example 1.7. Both these graphs have the same numbers of vertices and edges, and the same degree sequence $2, 2, 2, 2, 3, 3$. However, if one tries to find an isomorphism, even exhaustively, he **fails**. What is going on? □

Which property of the graphs prevents us from finding an isomorphism? Since there are (especially for larger graphs) too many potential bijective mappings to try them exhaustively in practice, we shall look for some other, **ad-hoc**, approaches. □


This time, the first graph has no triangle and the second one has one! Hence they cannot be isomorphic. □

Fact: No universal and efficient way of deciding an isomorphism between two graphs is currently known (the problem is **not known in P**). On the other hand, however, the problem is **neither NP-hard** to our knowledge, and so a general polynomial algorithm might emerge in the future. . .

1.4 Directed graphs and Multigraphs

On some occasions, we need to express a “direction” of a graph edge. □

We then speak about *directed graphs* in which edges (*directed arcs*) are ordered pairs of vertices (and they are drawn as *arrows*).




Definition 1.10. **Directed graph** (digraph) is a pair $D = (V, A)$ where $A \subseteq V \times V$. The notions of subgraphs and isomorphism naturally extend to digraphs.


Fact: Digraphs correspond to binary relations which need not be symmetric. □

Notation: An edge–arc $e = (u, v)$ in a digraph D has its *tail* in u and the *head* in v , or e joins “from u to v ”. The opposite arrow (v, u) is distinct from (u, v) !

For example, a *directed path* of length $n \geq 0$ looks as follows



and a *directed cycle* of length $n \geq 1$ is



Definition: The number of arcs from u in a digraph D is called the *outdegree* $d_D^+(u)$, and the number of those to u the *indegree* $d_D^-(u)$.

Generalizing even more...

On some other occasions one may want to speak about structures in which **more than one edge** exist between one pair of vertices, and the edges might have mixed types (undirected or directed, loops).

This leads to so called *incidence model* of a (multi)graph in which edges are elements on their own, along with the vertices; as opposed to our default *adjacency model* where only vertices are considered as core entities. □

Definition: A *mixed multigraph* is a triple $M = (V, F, \varepsilon)$ where $V \cap F = \emptyset$ and $\varepsilon : F \rightarrow \binom{V}{2} \cup V \cup (V \times V)$ is an *incidence mapping* of the (multi)edges.

In the definition,

- $\binom{V}{2}$ represents unoriented edges,
- V unoriented loops, and
- $V \times V$ oriented edges and loops.

