

Statistics for Computer Sciences

Lecture 10 to Lecture 12 Testing of Statistical Hypotheses

Stanislav Katina¹

¹Institute of Mathematics and Statistics, Masaryk University
Honorary Research Fellow, The University of Glasgow

December 2, 2015



Testing of Statistical Hypotheses

Null and alternative hypothesis

- ▶ a '**hypothesis**' is a theory which is assumed to be true unless evidence is obtained which indicates otherwise
- ▶ '**null**' means 'nothing' and the term '**null hypothesis**' (H_0) means a 'theory of no change' – that is 'no change' from what would be expected from past experience
- ▶ '**alternative hypothesis**' (H_1) means a 'theory of change' – that is 'change' from what would be expected from past experience
- ▶ the procedure which is used to decide between these two opposite theories is called '**hypothesis test**' or sometimes '**significance test**'
- ▶ **one-tail test** – test in which thy alternative hypothesis proposes a change in parameter in only one direction – increase or decrease
- ▶ **two-tail test** – test in which the alternative hypothesis suggests a difference in parameter in either direction



Testing of Statistical Hypotheses

Test statistic, rejection and acceptance region, critical value and quantile

- ▶ the **test statistic** is calculated from the sample – its value is used to decide whether the null hypothesis should be rejected
- ▶ the **rejection** (or **critical**) **region** gives the values of the test statistic for which the null hypothesis is rejected
- ▶ the **acceptance region** gives the values of the test statistic for which the null hypothesis is not rejected
- ▶ the boundary value(s) of the rejection region is (are) called the **critical value(s)** or **quantile(s)**
- ▶ the **significance level** α of a test gives the probability of the test statistic falling in the rejection region when null hypothesis is true



Testing of Statistical Hypotheses

Hypothesis testing procedure

- ▶ a **hypothesis** is a statement about a population parameter base on a sample from this population
- ▶ H_0 and H_1 are two complementary hypotheses in a hypothesis testing problem
- ▶ a **hypothesis testing procedure** or **hypothesis test** is a rule that specifies – for which sample values the decision is made to accept null hypothesis as true – and for which sample values H_0 is rejected
- ▶ the subset of sample space for which H_0 will be rejected is called **rejection region (critical region)**
- ▶ the complement of the rejection region is called the **acceptance region**



Testing of Statistical Hypotheses

Four possibilities

Four choices:

- A H_0 is true – our decision is to reject H_0
- B H_0 is true – our decision is not to reject H_0
- C H_1 is true – our decision is not to reject H_0
- D H_1 is true – our decision is to reject H_0

Decision-reality table:

decision/reality	H_0 is true	H_0 is not true
to reject H_0	Type I error	true decision
not to reject H_0	true decision	Type II error

◀ ▶ ⏪ ⏩ ⏴ ⏵ ⏶ ⏷ ⏸ ⏹ ⏺ ⏻ ⏼ ⏽ ⏾ ⏿ 🔍 ↺

Testing of Statistical Hypotheses

Empirical $100 \times (1 - \alpha)\%$ confidence intervals for parameter θ

Relationship of confidence interval and statistical test

- ▶ Empirical $100(1 - \alpha)\%$ confidence interval (CI) for parameter θ
- ▶ α -level hypothesis test about θ

Three types of intervals:

- ▶ **two-tailed** CI – $\Pr(LB(X) < \theta < UB(X)) = 1 - \alpha$
- ▶ **one-tailed (right-tailed)** CI – $\Pr(\theta < UB^*(X)) = 1 - \alpha$
- ▶ **one-tailed (left-tailed)** CI – $\Pr(LB_*(X) < \theta) = 1 - \alpha$

◀ ▶ ⏪ ⏩ ⏴ ⏵ ⏶ ⏷ ⏸ ⏹ ⏺ ⏻ ⏼ ⏽ ⏾ ⏿ 🔍 ↺

Testing of Statistical Hypotheses

Four possibilities

Four choices:

- A) $\Pr(A) = \Pr(\text{Type I error}) \leq \alpha$ [significance level]
- B) $\Pr(B) \geq 1 - \alpha$ [coverage probability, confidence coefficient (level)]
- C) $\Pr(C) = \Pr(\text{Type II error}) \leq \beta$
- D) $\Pr(D) \geq 1 - \beta$ [power]

Four choices (formalised):

- A) $1 - \alpha \leq \Pr(\text{don't reject } H_0 | H_0 \text{ is true})$
- B) $\alpha \geq \Pr(\text{CHPD}) = \Pr(\text{reject } H_0 | H_0 \text{ is true})$
- C) $\beta = \Pr(\text{CHDD}) = \Pr(\text{don't reject } H_0 | H_0 \text{ isn't true})$
- D) $1 - \beta = \Pr(\text{reject } H_0 | H_0 \text{ isn't true})$

◀ ▶ ⏪ ⏩ ⏴ ⏵ ⏶ ⏷ ⏸ ⏹ ⏺ ⏻ ⏼ ⏽ ⏾ ⏿ 🔍 ↺

Testing of Statistical Hypotheses

Acceptance region

Definition (Acceptance region of H_0)

Let X be a random variable with certain distribution (probabilistic model) dependent on parameter $\theta \in \Theta$, $g(\theta)$ is parametric function. We are testing null hypothesis $H_{01} : g(\theta) = g(\theta_0)$ against two-sided alternative $H_{11} : g(\theta) \neq g(\theta_0)$. Let (LB, UB) be interval estimate of parametric function $g(\theta)$ with coverage probability $1 - \alpha$. Then

$$\mathcal{A}_{1S,1} = \{LB, UB; g(\theta_0) \in (LB, UB)\}$$

is **acceptance region of a test H_{01} against H_{11} on significance level α** . If we are testing $H_{02} : g(\theta) \leq g(\theta_0)$ against one-sided (right) alternative $H_{12} : g(\theta) > g(\theta_0)$ and if LB_* be lower estimate of $g(\theta)$ with coverage probability $1 - \alpha$, then

$$\mathcal{A}_{1S,2} = \{LB_*; LB_* < g(\theta_0)\}$$

is **acceptance region of a test H_{02} against H_{12} on significance level α** . If we are testing $H_{03} : g(\theta) \geq g(\theta_0)$ against one-sided (left) alternative $H_{13} : g(\theta) < g(\theta_0)$ and if UB^* is upper estimate of $g(\theta)$ with coverage probability $1 - \alpha$, then

$$\mathcal{A}_{1S,3} = \{UB^*; UB^* > g(\theta_0)\}$$

is **acceptance region of a test H_{03} against H_{13} on significance level α** .

◀ ▶ ⏪ ⏩ ⏴ ⏵ ⏶ ⏷ ⏸ ⏹ ⏺ ⏻ ⏼ ⏽ ⏾ ⏿ 🔍 ↺

Testing of Statistical Hypotheses

Rejection region

Definition (Rejection (critical) region of H_0)

Let X be a random variable with certain distribution (probabilistic model) dependent on parameter $\theta \in \Theta$, $g(\theta)$ is parametric function. We are testing null hypothesis $H_{01} : g(\theta) = g(\theta_0)$ against two-sided alternative $H_{11} : g(\theta) \neq g(\theta_0)$. Let (LB, UB) be interval estimate of parametric function $g(\theta)$ with coverage probability $1 - \alpha$. Then

$$\mathcal{W}_{IS,1} = \{LB, UB; g(\theta_0) \notin (LB, UB)\}$$

is **critical region of a test H_{01} against H_{11} on significance level α** . If we are testing $H_{02} : g(\theta) \leq g(\theta_0)$ against one-sided (right) alternative $H_{12} : g(\theta) > g(\theta_0)$ and if LB_* be lower estimate of $g(\theta)$ with coverage probability $1 - \alpha$, then

$$\mathcal{W}_{IS,2} = \{LB_*; LB_* \geq g(\theta_0)\}$$

is **critical region of a test H_{02} against H_{12} on significance level α** . If we are testing $H_{03} : g(\theta) \geq g(\theta_0)$ against one-sided (left) alternative $H_{13} : g(\theta) < g(\theta_0)$ and if UB^* is upper estimate of $g(\theta)$ with coverage probability $1 - \alpha$, then

$$\mathcal{W}_{IS,3} = \{UB^*; UB^* \leq g(\theta_0)\}$$

is **critical region of a test H_{03} against H_{13} on significance level α** .



Testing of Statistical Hypotheses

Test criterion

Definition (Test criterion)

A **test criterion** is a test statistic $T = T_0 = T_0(X_1, X_2, \dots, X_n)$, with known asymptotic distribution if H_0 is known. The set of possible values of T_0 is divided to two subsets, i.e. **acceptance region H_0** (notation \mathcal{A}) and **critical region H_0** (notation \mathcal{W}). These two regions are divided by **critical values** $t_{\alpha/2}$ and $t_{1-\alpha/2}$, resp. t_α and $t_{1-\alpha}$ (for particular H_0 and H_1) of the distribution of test statistics T_0 (if H_0 is true).

Definition (Confidence interval)

A **confidence interval (CI)** is a type of interval estimate of a population parameter θ . It is an observed, often called **empirical**, interval (i.e., it is calculated from the observations) that includes the value of an unobservable parameter θ if the experiment is repeated. The frequency that observed interval contains the parameter is determined by the **confidence coefficient** $1 - \alpha$ (i.e. **confidence level, coverage probability**).



Testing of Statistical Hypotheses

To carry out a hypothesis test

- Step 1 define the null and alternative hypothesis (H_0 and H_1)
- Step 2 decide on a significance level $\alpha = 0.1, 0.05, 0.01$
- Step 3 calculate the test statistic (test criterion) T_0
- Step 3 determine the critical value(s)
- Step 5 decide on the outcome of the test (reject/don't reject H_0) depending on one of the following ways:
 - ▶ base on critical region $\mathcal{W} = \mathcal{W}_T$ (observed test statistic $t_0 = t_{\text{obs}}$ and critical values $t_{\alpha/2}$ and $t_{1-\alpha/2}$, resp. t_α and $t_{1-\alpha}$),
 - ▶ base on critical region \mathcal{W}_{IS} , t.j. empirical confidence interval (and $g(\theta_0)$),
 - ▶ base on p-value.
- Step 6 state the conclusion in words



Testing of Statistical Hypotheses

To carry out a hypothesis test – based on test statistic and critical value

Definition (Testing based on critical region \mathcal{W})

Rejecting H_0 . If observed test statistic (realisation of test statistic) t_0 of test statistic T_0 is within a critical region \mathcal{W} (equivalently is not from an acceptance region \mathcal{A}), H_0 is rejected at a significance level α , i.e. we do have sufficiently enough evidence to reject H_0 .

Not rejecting H_0 . If observed test statistic t_0 of test statistic T_0 is within an acceptance region \mathcal{A} (equivalently, it is not from a critical region \mathcal{W}), H_0 is not rejected at a significance level α , i.e. we don't have sufficiently enough evidence to reject H_0 .

Let t_{\min} be the smallest possible value of a test criteria T_0 and t_{\max} be the highest possible value of a test criteria T_0 , then

1. **two-sided alternative** – critical region $\mathcal{W}_1 = (t_{\min}, t_{1-\alpha/2}) \cup (t_{\alpha/2}, t_{\max})$,
2. **one-sided (right) alternative** – critical region $\mathcal{W}_2 = (t_\alpha, t_{\max})$,
3. **one-sided (left) alternative** – critical region $\mathcal{W}_3 = (t_{\min}, t_{1-\alpha})$.



Testing of Statistical Hypotheses

To carry out a hypothesis test – based on CI

Definition (Testing based on CI)

Rejecting H_0 : If $g(\theta) = g(\theta_0)$ is within CI (H_0 is valid), H_0 is rejected at the significance level α , i.e. we do have sufficiently enough evidence to reject H_0 .

Not rejecting H_0 : If $g(\theta) = g(\theta_0)$ is not within CI (H_0 is valid), H_0 isn't rejected at a significance level α , i.e. we don't have sufficiently enough evidence to reject H_0 .

Relationship of confidence interval and statistical test

- ▶ hypothesis testing \equiv CIs
- ▶ α -level hypothesis test \equiv $100(1 - \alpha)\%$ CI
- ▶ **one-tail test** \equiv one-sided CI (**left-sided CI** \equiv **right-sided alternative**, **right-sided CI** \equiv **left-sided alternative**)
- ▶ **two-tail test** \equiv two-sided CI
- ▶ parameter(s) \in CI \equiv not reject H_0
- ▶ parameter(s) \notin CI \equiv reject H_0



Testing of Statistical Hypotheses

To carry out a hypothesis test – based on p-value (observed significance level)

Definition (Testing based on p-value)

Minimal significance level α (for some test statistic T_0), base on which $H_{02} : g(\theta) \leq g(\theta_0)$ is rejected (tested against $H_{12} : g(\theta) > g(\theta_0)$), is called **observed significance level** or **p-value**, i.e.

$$\text{p-value} = \alpha_{\text{obs}} = \sup_{\theta \in \Theta_0} \Pr(T(X_1, X_2, \dots, X_n) \geq T(x_1, x_2, \dots, x_n); \theta).$$

This could be written less formally as **p-value = Pr(any test statistics equal or greater than observed | H_0 is true)**.

The closer α_{obs} is to zero, the smaller is the probability that any test statistic $T(X_1, X_2, \dots, X_n)$ produces a p-value (under H_0) equal to or smaller than that observed, while the probability is higher under H_1 . Therefore, p-value could be understood as **an indicator of credibility of H_0** .



Testing of Statistical Hypotheses

To carry out a hypothesis test – based on p-value (observed significance level)

- ▶ Usually, if $\alpha_{\text{obs}} < \alpha = 0.05$, there is sufficiently enough evidence to reject H_0 and the result of a test **is statistically significant**.
- ▶ While $\alpha_{\text{obs}} > \alpha = 0.1$, there is sufficiently enough evidence to reject H_0 and the result of a test **is not statistically significant**.
- ▶ The values between 0.05 and 0.1 should be taken as reference points in a broad sense. As α_{obs} gets closer to either boundary point of the interval $(0.05, 0.1)$, so this is taken as increasing evidence for one or other alternative.
- ▶ Situation with $\alpha_{\text{obs}} \in (0.05, 0.1)$ are usually most difficult to handle and the result is here **marginally statistically significant**.



Testing of Statistical Hypotheses

To carry out a hypothesis test – based on p-value (observed significance level)

Wording of the results of a statistical test:

range for p-value	stars of significance	wording of the result
(0, 0.001)	***	extremely highly statistically significant
(0.001, 0.01)	**	high statistically significant
(0.01, 0.05)	*	statistically significant
(0.05, 0.1)	.	marginally statistically significant
(0.1, 1)		non-significant



Testing of Statistical Hypotheses

To carry out a hypothesis test – based on p-value (observed significance level)

Interpretation of p-values:

- ▶ p-value < 0.001: the **prevalence** of an estimated effect is smaller than one to one thousand (the **odds** of estimated effect is smaller than 1 : 999), if an effect is not present in a population (the presence of such an effect is **highly improbable**, if an effect is not present in a population – and – the presence of such an effect is **highly probable**, if an effect is present in a population)
- ▶ p-value < 0.01: the **prevalence** of an estimated effect is smaller than one to one hundred (the **odds** of estimated effect is smaller than 1 : 99), if an effect is not present in a population (the presence of such an effect is **very improbable**, if an effect is not present in a population – and – the presence of such an effect is **very probable**, if an effect is present in a population)
- ▶ p-value < 0.05: the **prevalence** of an estimated effect is smaller than one to one hundred (the **odds** of estimated effect is smaller than 5 : 95 or 1 : 19), if an effect is not present in a population (the presence of such an effect is **sufficiently improbable**, if an effect is not present in a population – and – the presence of such an effect is **sufficiently probable**, if an effect is present in a population)
- ▶ p-value \geq 0.05: the prevalence of an estimated effect is five to one hundred or greater (5 % or more);
- ▶ p-value = k , $k \in (0.05, 1)$: the prevalence of an estimated effect is $100 \times k$ to one hundred ($100 \times k$ % or more).



Testing of Statistical Hypotheses

To carry out a hypothesis test – based on p-value (observed significance level)

How is the p-value (mostly) calculated?

1. **two-sided alternative** –
p-value = $2 \min(\Pr(T_0 \leq t_0|H_0), \Pr(T_0 \geq t_0|H_0))$, e.g. for normal and Student distribution of test statistic (symmetric distributions) and for χ_{df}^2 and F_{df_1, df_2} distribution of test statistic (asymmetric distributions) or
p-hodnota = $\min(\Pr(T_0 \leq t_0|H_0), \Pr(T_0 \geq t_0|H_0))$, e.g. for χ_{df}^2 and F_{df_1, df_2} distribution of test statistic (asymmetric distributions)
2. **one-sided (right) alternative** – p-value = $\Pr(T_0 \geq t_0|H_0)$
3. **one-sided (left) alternative** – p-value = $\Pr(T_0 \leq t_0|H_0)$



Testing of Statistical Hypotheses

On a philosophical level

- ▶ distinction between 'rejecting H_0 ' and 'accepting H_1 '
- ▶ 'rejecting H_0 ' – nothing implies about what state the experimenter is accepting, only that the state defined by H_0 is being rejected
- ▶ distinction between 'accepting H_0 ' and 'not rejecting H_0 '
- ▶ 'accepting H_0 ' – the experimenter is willing to assert the state of nature specified by H_0
- ▶ 'not rejecting H_0 ' – the experimenter really does not believe H_0 but does not have the evidence to reject it



Testing of Statistical Hypotheses

Conservative and liberal test and CI

Definition (Conservative and liberal test)

A test with **actual/observed significance level** smaller than **nominal significance level** α , is called **conservative** (the test should theoretically be "rejecting quickly" H_0 , but, in reality, it is the opposite, i.e. the test is "rejecting slowly").

A test with **actual/observed significance level** greater than **nominal significance level** α , is called **liberal** (the test should theoretically be "rejecting slowly" H_0 , but, in reality, it is the opposite, i.e. the test "rejecting quickly").

Definition (Conservative and liberal CI)

CI with **actual/real coverage probability** greater than **nominal coverage probability** $1 - \alpha$, is called **conservative** (i.e. the probability that θ_0 is within CI is greater than expected).

CI with **actual/real coverage probability** smaller than **nominal coverage probability** $1 - \alpha$, is called **liberal** (i.e. the probability that θ_0 is within CI is smaller than expected).



Testing of Statistical Hypotheses

Likelihood ratio – generalised relative likelihood

Two types of hypotheses:

1. **simple hypothesis** – $H_0 : \theta = \theta_0$ against $H_1 : \theta \neq \theta_0$, then **simple likelihood ratio** is equal to

$$\lambda(\mathbf{x}) = \lambda = \frac{L(\theta_0|\mathbf{x})}{\sup_{\theta \in \Theta} L(\theta|\mathbf{x})} = \frac{L(\theta_0|\mathbf{x})}{L(\hat{\theta}|\mathbf{x})},$$

where $\lambda(\mathbf{x}) = \mathcal{L}(\theta_0|\mathbf{x})$ is test statistic and $L(\theta|\mathbf{x})$ is continuous for all \mathbf{x} .

2. **composite hypothesis** – $H_0 : \theta \in \Theta_0$ against $H_1 : \theta \in \Theta_1$, then **generalised likelihood ratio** is equal to

$$\lambda(\mathbf{x}) = \frac{\sup_{\theta \in \Theta_0} L(\theta|\mathbf{x})}{\sup_{\theta \in \Theta} L(\theta|\mathbf{x})}.$$

Navigation icons

Testing of Statistical Hypotheses

Three test statistics

Geometrical interpretation:

1. U_{LR} – is measuring properly standardised difference between log-likelihoods in $\hat{\theta}$ and θ_0 (i.e. in direction of y axis)
2. U_W – is measuring properly standardised absolute value of a difference of $\hat{\theta}$ a θ_0 (in direction of x axis)
3. U_S – is measuring properly standardised slope of log-ratio in θ_0

Example (normal distribution)

Let $X \sim N(\mu, \sigma^2)$, where σ^2 is known, $H_0 : \theta = \theta_0$ against $H_1 : \theta \neq \theta_0$, where $\theta = \mu$. Then

1. $U_{LR} = -2(l(\theta_0|\mathbf{X}) - l(\hat{\theta}|\mathbf{X})) = -\sum_{i=1}^n (X_i - \bar{X})^2 / \sigma^2 + \sum_{i=1}^n (X_i - \mu_0)^2 / \sigma^2 = n \frac{(\bar{X} - \mu_0)^2}{\sigma^2},$
2. $U_W = (\bar{X} - \mu_0)^2 \mathcal{I}(\bar{X}) = n \frac{(\bar{X} - \mu_0)^2}{\sigma^2},$
3. $U_S = \frac{(\mathcal{S}(\mu_0))^2}{\mathcal{I}(\mu_0)} = \frac{(n(\bar{X} - \mu_0) / \sigma^2)^2}{n / \sigma^2} = n \frac{(\bar{X} - \mu_0)^2}{\sigma^2}.$

All three test statistics are equal, i.e. $U_{LR} = U_W = U_S$.

Navigation icons

Testing of Statistical Hypotheses

Likelihood ratio test statistic

Subsets of Θ , Θ_0 and Θ_1 , remain the same after monotone transformation of $\lambda(\mathbf{x})$, i.e. the statistical tests before and after transformation are equivalent. Therefore, **likelihood ratio test statistic** is equal to

$$U_{LR} = -2 \ln \lambda(\mathbf{X}).$$

Its realisation, **observed likelihood ratio test statistic**, is equal to $u_{LR} = -2 \ln \lambda(\mathbf{x})$, where $u_{LR} \in (0, \infty)$.

Navigation icons

Testing of Statistical Hypotheses

Three test statistics

If θ is a scalar, three test statistics are defined as:

1. $U_{LR} = -2(l(\theta_0|\mathbf{X}) - l(\hat{\theta}|\mathbf{X})) \stackrel{\mathcal{D}}{\sim} \chi_1^2,$
2. $U_W = (\hat{\theta} - \theta_0)^2 \mathcal{I}(\hat{\theta}) \stackrel{\mathcal{D}}{\sim} \chi_1^2$ and equivalently $U_W^{1/2} = Z_W \stackrel{\mathcal{D}}{\sim} N(0, 1),$
3. $U_S = \frac{(\mathcal{S}(\theta_0))^2}{\mathcal{I}(\theta_0)} \stackrel{\mathcal{D}}{\sim} \chi_1^2$ and equivalently $U_S^{1/2} = Z_S \stackrel{\mathcal{D}}{\sim} N(0, 1),$

If θ is a vector, three test statistics are defined as:

1. $U_{LR} = -2(l(\theta_0|\mathbf{X}) - l(\hat{\theta}|\mathbf{X})) \stackrel{\mathcal{D}}{\sim} \chi_k^2,$
2. $U_W = (\hat{\theta} - \theta_0)^T \mathcal{I}(\hat{\theta})(\hat{\theta} - \theta_0) \stackrel{\mathcal{D}}{\sim} \chi_k^2,$
3. $U_S = (\mathcal{S}(\theta_0))^T (\mathcal{I}(\theta_0))^{-1} \mathcal{S}(\theta_0) \stackrel{\mathcal{D}}{\sim} \chi_k^2.$

Navigation icons

Testing of Statistical Hypotheses

Three test statistics and related confidence intervals

If θ is a scalar, three confidence intervals are defined as follows:

1. **likelihood ratio empirical** $(1 - \alpha) \times 100\%$ **CI for θ** is defined as

$$CS_{1-\alpha} = \{\theta : U_{LR}(\theta) < \chi_1^2(\alpha)\},$$

$$\text{where } U_{LR}(\theta) = -2 \ln \frac{L(\theta|\mathbf{x})}{L(\hat{\theta}|\mathbf{x})}.$$

2. **Wald empirical** $(1 - \alpha) \times 100\%$ **CI for θ** is defined based on a pivot (pivotal statistics) $T_{\text{piv}} = U_W(\theta)$
3. **Score empirical** $(1 - \alpha) \times 100\%$ **CI for θ** is defined based on a pivot $T_{\text{piv}} = U_S(\theta)$

If θ is a vector, CIs can be generalized to **confidence set** $CS_{1-\alpha}$.

- ▶ If $k = 2$, $CS_{1-\alpha}$ is an **confidence ellipse**.
- ▶ If $k > 2$, $CS_{1-\alpha}$ is an **confidence ellipsoid**.

Additionally, if $k = 1$, $CS_{1-\alpha}$ is an **confidence interval**.



Testing of Statistical Hypotheses

Confidence intervals

Wald empirical $(1 - \alpha) \times 100\%$ **CI for θ** is defined as

$$(d, h) = \left(\hat{\theta} - t_{\alpha/2} \widehat{SE}[\hat{\theta}], \hat{\theta} + t_{\alpha/2} \widehat{SE}[\hat{\theta}] \right),$$

where the critical value $t_{\alpha/2}$ depends on the choice of $\hat{\theta}$.

Likelihood ratio empirical $(1 - \alpha) \times 100\%$ **CI for θ** is defined by its lower and upper bounds as $k\%$ cut-offs of standardized relative log-likelihood as follows

$$\Pr \left(\frac{L(\theta|\mathbf{x})}{L(\hat{\theta}|\mathbf{x})} > c_\alpha \right) = \Pr \left(-2 \ln \frac{L(\theta|\mathbf{x})}{L(\hat{\theta}|\mathbf{x})} < -2 \ln c_\alpha \right) = 1 - \alpha,$$

where $c_\alpha = e^{-\frac{1}{2}\chi_1^2(\alpha)}$. Then

- ▶ if $1 - \alpha = 0.95$, then $c_\alpha = 0.1465001 \doteq 0.15$ (15% cut-off),
- ▶ if $1 - \alpha = 0.90$, then $c_\alpha = 0.2585227 \doteq 0.26$ (26% cut-off),
- ▶ if $1 - \alpha = 0.99$, then $c_\alpha = 0.0362452 \doteq 0.04$ (4% cut-off).



Testing of Statistical Hypotheses

Likelihood confidence intervals – bisection method

Bisection method

Let $\theta_{01}, \theta_{02} \in \langle \theta_L, \theta_U \rangle$ and $f(\theta_{01})f(\theta_{02}) < 0$, $f(\cdot)$ is continuous with at least one root within the interval $\langle \theta_{01}, \theta_{02} \rangle$, where

$$f(\theta) = -2 \ln \mathcal{L}(\theta|\mathbf{x}) - \chi_1^2(\alpha) = 0.$$

If the first derivative of $f(\cdot)$ is having constant sign, then exactly one root $\theta^* \in \langle \theta_{01}, \theta_{02} \rangle$ of $f(\theta) = 0$ exists.

The iterative process is defined as follows:

1. **initialisation step** – starting point $\theta^{(0)} = (\theta_{01} + \theta_{02})/2$ and $i = 1$,
2. **updating equations** – substitution of the boundaries θ_{01} and θ_{02} is defined as

$$\langle \theta_{i1}, \theta_{i2} \rangle = \begin{cases} \langle \theta_{i-1,1}, \theta^{(i-1)} \rangle, & \text{if } f(\theta_{i-1,1})f(\theta^{(i-1)}) < 0 \\ \langle \theta^{(i-1)}, \theta_{i-1,2} \rangle, & \text{if } f(\theta_{i-1,1})f(\theta^{(i-1)}) > 0 \end{cases}$$

if $f(\theta^{(i-1)}) = 0$, then **end**, if not,



Testing of Statistical Hypotheses

Likelihood confidence intervals – Brent-Dekker method

Example (Brent-Dekker method)

Let $X \sim \text{Bin}(N, p)$, where $N = 10$ and $n = x = 8$. Estimate the boundaries of empirical $100 \times (1 - \alpha)\%$ CI for (1) p and (2) odds $\frac{p}{1-p}$.

The empirical CI are of the two types (A) likelihood and (B) Wald.

Draw the log-likelihood function and its quadratic approximation with the lower and upper boundary of CI.

Solution (partial)

Wald empirical $100 \times (1 - \alpha)\%$ CI for p :

$$\hat{p} = \frac{8}{10} = 0.8; \widehat{SE}[\hat{p}] = \sqrt{\frac{\hat{p}(1-\hat{p})}{N}} = 0.13.$$

$$(d, h) = \left(\hat{p} - u_{\alpha/2} \widehat{SE}[\hat{p}], \hat{p} + u_{\alpha/2} \widehat{SE}[\hat{p}] \right) = (0.55, 1.05).$$

Likelihood empirical $100 \times (1 - \alpha)\%$ CI for p :

$$CS_{1-\alpha} = \left\{ p : -2 \ln \frac{L(p|\mathbf{x})}{L(\hat{p}|\mathbf{x})} \leq 3.84 \right\}, \text{ where } (d, h) = (0.50, 0.96),$$

Wald empirical $100 \times (1 - \alpha)\%$ CI for $g(p)$:

$$g(\hat{p}) = \ln \frac{\hat{p}}{1-\hat{p}} = \log \frac{0.8}{0.2} = 1.39.$$

$$\frac{\partial}{\partial p} g(p) = \frac{1}{p} + \frac{1}{1-p};$$

$$\widehat{SE}[g(\hat{p})] = \widehat{SE}[\hat{p}] \left(\frac{1}{\hat{p}} + \frac{1}{1-\hat{p}} \right) = \sqrt{\frac{\hat{p}(1-\hat{p})}{N}} \left(\frac{1}{\hat{p}} + \frac{1}{1-\hat{p}} \right) = \sqrt{\frac{1}{n} + \frac{1}{N-n}} = 0.79.$$

Then $(d_g, h_g) = (-0.16, 2.94)$ and back-transformed $(d, h) = (0.46, 0.95)$.



Testing of Statistical Hypotheses

Likelihood confidence intervals – Brent-Dekker method

```
1 | x <- 8; N <- 10
2 | probs <- seq(0.4, .99, length=1000)
3 | like <- dbinom(8,10,probs)
4 | rellike <- like/max(like)
5 | relloglike <- -2*log(rellike)
6 | cutoff <- exp(-1/2*qchisq(0.95,df=1)) #0.1465001
7 | like.CI.p <- range(probs[rellike>cutoff]) #0.5009910 0.9634234
8 | cutoff <- qchisq(0.95,df=1) #3.841459
9 | like.CI.p <- range(probs[relloglike<cutoff]) #0.500991 0.9634234
10 |
11 | p.hat <- x/N
12 | i.hat <- N/p.hat/(1-p.hat)
13 | loglikeapprox <- -i.hat/2*(probs-p.hat)^2
14 | ra <- range(log(rellike))
15 | wald.is.p <- p.hat + c(-1,1)*qnorm(0.975)*sqrt(1/i.hat)
16 | wald.is.p # 0.552082 1.047918
17 |
18 | gprobs <- log(probs) - log(1-probs)
19 | gp.hat <- log(p.hat) - log(1-p.hat)
20 | i.hat <- x*(N-x)/N
21 | lgp <- -i.hat/2*(gprobs-gp.hat)^2
22 | x <- (gp.hat+c(-1,1)*qnorm(0.975)*sqrt(1/i.hat)) # -0.1632 2.9358
23 | wald.is.gp <- exp(x)/(1+exp(x))
24 | wald.is.gp # 0.4592920 0.9495872
```

◀ ▶ ⏪ ⏩ ⏴ ⏵ ⏶ ⏷ ⏸ ⏹ ⏺ ⏻ ⏼ ⏽ ⏾ ⏿ 🔍 ↺

Testing of Statistical Hypotheses

Likelihood confidence intervals – other numerical method




Figure: Log-likelihood of p and its quadratic approximation

◀ ▶ ⏪ ⏩ ⏴ ⏵ ⏶ ⏷ ⏸ ⏹ ⏺ ⏻ ⏼ ⏽ ⏾ ⏿ 🔍 ↺

Testing of Statistical Hypotheses

To carry out a hypothesis test

Number of (in)dependent samples for θ , $g(\theta)$, θ and $g(\theta)$:

- ▶ **one-sample problem** about – mean, variance, probability distribution, correlation coefficient, probability
- ▶ **two-sample problem** about – difference in means, ratio of variances, difference in probability distributions, difference in correlation coefficients, difference in probabilities
- ▶ **multiple sample problem** about – means, variances, probability distributions, correlation coefficients, probabilities
- ▶ **paired problem** – the mean of the differences

Dimension:

- ▶ **univariate problem**
- ▶ **multivariate problem**

◀ ▶ ⏪ ⏩ ⏴ ⏵ ⏶ ⏷ ⏸ ⏹ ⏺ ⏻ ⏼ ⏽ ⏾ ⏿ 🔍 ↺

Testing of Statistical Hypotheses

One-sample problems

- ▶ **one-sample Z-test** for the mean of one population
- ▶ **one-sample Student t -test** for the mean of one population
- ▶ **one-sample χ^2 -test** for the variance of one population
- ▶ **one-sample Kolmogorov-Smirnov test** for the empirical probability distribution function of one population
- ▶ **one-sample Z-test** for the population proportion of one population
- ▶ **one-sample T -test** for the correlation coefficient of one population

◀ ▶ ⏪ ⏩ ⏴ ⏵ ⏶ ⏷ ⏸ ⏹ ⏺ ⏻ ⏼ ⏽ ⏾ ⏿ 🔍 ↺

Testing of Statistical Hypotheses

Two-sample problems

- ▶ **two-sample Z-test** for the difference between the means of two populations
- ▶ **two-sample Student t -test** for the difference between the means of two populations
- ▶ **two-sample F -test** for the ratio of the variances of two populations
- ▶ **two-sample Kolmogorov-Smirnov test** for the difference between two empirical probability distribution functions
- ▶ **two-sample Z-test** for the difference between two population proportions
- ▶ **two-sample T -test** for the difference between correlation coefficients of two populations