

IA008: Computational Logic

4. Deduction

Achim Blumensath

blumens@fi.muni.cz

Faculty of Informatics, Masaryk University, Brno


Tableaux

Tableau Proofs

For simplicity: first-order logic **without equality**

Statements φ true or φ false

Rule



Interpretation

If $\varphi \sigma$ is **possible** then so is $\psi_i \tau_i, \dots, \vartheta_i v_i$, for some i .

Tableaux

Construction

A **tableau** for a formula φ is constructed as follows:

- ▶ start with φ false
- ▶ choose a branch of the tree
- ▶ choose a statement ψ value on the branch
- ▶ choose a rule with head ψ value
- ▶ add it at the bottom of the branch
- ▶ repeat until every branch contains both statements ψ true and ψ false for some formula ψ

$R(\bar{t})$ true

$R(\bar{t})$ false

$\neg\varphi$ true

$\neg\varphi$ false

φ false

φ true

$\varphi \wedge \psi$ true

$\varphi \wedge \psi$ false

$\varphi \vee \psi$ true

$\varphi \vee \psi$ false

φ true

φ false

ψ false

φ true

ψ true

φ false

ψ true

ψ false

$\varphi \rightarrow \psi$ true

$\varphi \rightarrow \psi$ false

$\varphi \leftrightarrow \psi$ true

$\varphi \leftrightarrow \psi$ false

φ false

ψ true

φ true

φ true

φ false

φ true

φ false

ψ false

ψ true

ψ false

ψ false

ψ true

$\forall x\varphi$ true

$\forall x\varphi$ false

$\exists x\varphi$ true

$\exists x\varphi$ false

$\varphi[x \mapsto t]$ true

$\varphi[x \mapsto c]$ false


$\varphi[x \mapsto c]$ true

$\varphi[x \mapsto t]$ false


c a new constant symbol, t an arbitrary term

Example

$(A \vee B) \rightarrow \neg(\neg A \wedge \neg B)$ false



$\neg(\neg A \wedge \neg B) \rightarrow (A \vee B)$ false



Example

$\exists x \forall y R(x, y) \rightarrow \forall y \exists x R(x, y)$ false

$\exists x \forall y R(x, y)$ true

$\forall y \exists x R(x, y)$ false

$\forall y R(c, y)$ true

$\exists x R(x, d)$ false

$R(c, d)$ true

$R(c, d)$ false

$\forall x R(x, x) \rightarrow \forall x \exists y R(f(x), y)$

$\forall x R(x, x)$ true

$\forall x \exists y R(f(x), y)$ false

$\exists y R(f(c), y)$ false

$R(f(c), f(c))$ false

$R(f(c), f(c))$ true

Soundness and Completeness

Theorem

A first-order formula φ is valid if, and only if, there exists a tableau T for φ false where every branch is contradictory.

Terminology

A tableau **for** a statement φ value is a tableau T where the root is labelled with φ value.

A branch β is **contradictory** if it contains both statements ψ true and ψ false, for some formula ψ .

A branch β is **consistent with** a structure \mathfrak{A} if

- ▶ $\mathfrak{A} \models \psi$, for all statements ψ true on β and
- ▶ $\mathfrak{A} \not\models \psi$, for all statements ψ false on β .

A branch β is **complete** if, for every atomic formula ψ , it contains one of the statements ψ true or ψ false.

Proof Sketch: Completeness

Lemma

If β is consistent with \mathfrak{A} and we extend the tableau by applying a rule, the new tableau has a branch β' extending β that is consistent with \mathfrak{A} .

Corollary

If $\mathfrak{A} \not\models \varphi$, then every tableau for φ false has a branch that is not contradictory.

Corollary

If φ is not valid, there is no tableau for φ false where all branches are contradictory.

Proof Sketch: Soundness

Lemma

If every tableau for φ false has a non-contradictory branch, there exists a tableau for φ false with a branch β that is complete and non-contradictory.

Lemma

If a branch β is complete and non-contradictory, there exists a structure \mathfrak{A} such that β is consistent with \mathfrak{A} .

Corollary

If every tableau for φ false has a non-contradictory branch, there exists a structure \mathfrak{A} with $\mathfrak{A} \not\models \varphi$.

Natural Deduction

Proof Calculi

Notation

$\psi_1, \dots, \psi_n \vdash \varphi$ φ is **provable** with **assumptions** ψ_1, \dots, ψ_n

Proof Calculi

Notation

$\psi_1, \dots, \psi_n \vdash \varphi$ φ is **provable** with **assumptions** ψ_1, \dots, ψ_n

φ is **provable** if $\vdash \varphi$.

Proof Calculi

Notation

$\psi_1, \dots, \psi_n \vdash \varphi$ φ is **provable** with **assumptions** ψ_1, \dots, ψ_n

φ is **provable** if $\vdash \varphi$.

Rules

$$\frac{\Gamma_1 \vdash \varphi_1 \dots \Gamma_n \vdash \varphi_n}{\Delta \vdash \psi} \quad \begin{array}{l} \text{premises} \\ \text{conclusion} \end{array} \quad \varphi_1 \wedge \dots \wedge \varphi_n \Rightarrow \psi$$

Proof Calculi

Notation

$\psi_1, \dots, \psi_n \vdash \varphi$ φ is **provable** with **assumptions** ψ_1, \dots, ψ_n

φ is **provable** if $\vdash \varphi$.

Rules

$$\frac{\Gamma_1 \vdash \varphi_1 \dots \Gamma_n \vdash \varphi_n}{\Delta \vdash \psi} \quad \begin{array}{l} \text{premises} \\ \text{conclusion} \end{array} \quad \varphi_1 \wedge \dots \wedge \varphi_n \Rightarrow \psi$$

Axiom

$$\frac{}{\Delta \vdash \psi} \quad \text{rule without premises}$$

Proof Calculi

Notation

$\psi_1, \dots, \psi_n \vdash \varphi$ φ is **provable** with **assumptions** ψ_1, \dots, ψ_n

φ is **provable** if $\vdash \varphi$.

Rules

$$\frac{\Gamma_1 \vdash \varphi_1 \dots \Gamma_n \vdash \varphi_n}{\Delta \vdash \psi} \quad \begin{array}{l} \text{premises} \\ \text{conclusion} \end{array} \quad \varphi_1 \wedge \dots \wedge \varphi_n \Rightarrow \psi$$

Axiom

$$\frac{}{\Delta \vdash \psi} \quad \text{rule without premises}$$

Remark

Tableaux speak about **possibilities** while Natural Deduction proofs speak about **necessities**.

Proof Calculi

Derivation

$$\frac{\frac{\overline{\Gamma \vdash \varphi} \quad \overline{\Delta_0 \vdash \psi_0}}{\Delta_1 \vdash \psi_1} \quad \overline{\Gamma' \vdash \varphi'}}{\Sigma \vdash \vartheta}$$

tree of rules

Natural Deduction (propositional part)

$$(I_{\top}) \frac{}{\Gamma \vdash \top}$$

$$(I_{\wedge}) \frac{\Gamma \vdash \varphi \quad \Delta \vdash \psi}{\Gamma, \Delta \vdash \varphi \wedge \psi}$$

$$(I_{\vee}) \frac{\Gamma \vdash \varphi}{\Gamma \vdash \varphi \vee \psi} \quad \frac{\Gamma \vdash \psi}{\Gamma \vdash \varphi \vee \psi}$$

$$(I_{\neg}) \frac{\Gamma, \varphi \vdash \perp}{\Gamma \vdash \neg \varphi}$$

$$(I_{\perp}) \frac{\Gamma \vdash \varphi \quad \Gamma \vdash \neg \varphi}{\Gamma \vdash \perp}$$

$$(I_{\rightarrow}) \frac{\Gamma, \varphi \vdash \psi}{\Gamma \vdash \varphi \rightarrow \psi}$$

$$(I_{\leftrightarrow}) \frac{\Gamma, \varphi \vdash \psi \quad \Delta, \psi \vdash \varphi}{\Gamma, \Delta \vdash \varphi \leftrightarrow \psi}$$

$$(Ax) \frac{}{\Gamma, \varphi \vdash \varphi}$$

$$(E_{\wedge}) \frac{\Gamma \vdash \varphi \wedge \psi}{\Gamma \vdash \varphi} \quad \frac{\Gamma \vdash \varphi \wedge \psi}{\Gamma \vdash \psi}$$

$$(E_{\vee}) \frac{\Gamma \vdash \varphi \vee \psi \quad \Delta, \varphi \vdash \vartheta \quad \Delta', \psi \vdash \vartheta}{\Gamma, \Delta, \Delta' \vdash \vartheta}$$

$$(E_{\neg}) \frac{\Gamma, \neg \varphi \vdash \perp}{\Gamma \vdash \varphi}$$

$$(E_{\perp}) \frac{\Gamma \vdash \perp}{\Gamma \vdash \varphi}$$

$$(E_{\rightarrow}) \frac{\Gamma \vdash \varphi \quad \Delta \vdash \varphi \rightarrow \psi}{\Gamma, \Delta \vdash \psi}$$

$$(E_{\leftrightarrow}) \frac{\Gamma \vdash \varphi \quad \Delta \vdash \varphi \leftrightarrow \psi}{\Gamma, \Delta \vdash \psi} \quad \frac{\Gamma \vdash \psi \quad \Delta \vdash \psi \leftrightarrow \varphi}{\Gamma, \Delta \vdash \varphi}$$

Examples

$$\frac{\frac{\frac{\varphi \vee \psi, \neg\varphi \wedge \neg\psi \vdash \varphi \vee \psi}{\varphi \vee \psi, \neg\varphi \wedge \neg\psi \vdash \perp} \quad \frac{\frac{\varphi \vdash \varphi}{\neg\varphi \wedge \neg\psi \vdash \neg\varphi} \quad \frac{\frac{\overline{\neg\varphi \wedge \neg\psi \vdash \neg\varphi \wedge \neg\psi}}{\neg\varphi \wedge \neg\psi \vdash \neg\varphi}}{\varphi, \neg\varphi \wedge \neg\psi \vdash \perp} \quad \frac{\dots}{\psi, \neg\varphi \wedge \neg\psi \vdash \perp}}{\varphi \vee \psi, \neg\varphi \wedge \neg\psi \vdash \perp}}{\varphi \vee \psi \vdash \neg\varphi \wedge \neg\psi \rightarrow \perp}}{\varphi \vee \psi \vdash \neg(\neg\varphi \wedge \neg\psi)}{\vdash (\varphi \vee \psi) \rightarrow \neg(\neg\varphi \wedge \neg\psi)}$$

Natural Deduction (quantifiers and equality)

$$(I_{\exists}) \frac{\Gamma \vdash \varphi[x \mapsto t]}{\Gamma \vdash \exists x \varphi}$$

$$(E_{\exists}) \frac{\Gamma \vdash \exists x \varphi \quad \Delta, \varphi[x \mapsto c] \vdash \psi}{\Gamma, \Delta \vdash \psi}$$

$$(I_{\forall}) \frac{\Gamma \vdash \varphi[x \mapsto c]}{\Gamma \vdash \forall x \varphi}$$

$$(E_{\forall}) \frac{\Gamma \vdash \forall x \varphi}{\Gamma \vdash \varphi[x \mapsto t]}$$

$$(I_{=}) \frac{}{\Gamma \vdash t = t}$$

$$(E_{=}) \frac{\Gamma \vdash s = t \quad \Delta \vdash \varphi[x \mapsto s]}{\Gamma, \Delta \vdash \varphi[x \mapsto t]}$$

c a **new** constant symbol, s, t arbitrary terms

Examples

$$s = t \vdash t = s$$

Examples

$$s = t \vdash t = s \quad \frac{\frac{}{s = t \vdash s = t} \quad \frac{}{\vdash s = s}}{s = t \vdash t = s} \quad (\text{E}_=)$$

Examples

$$s = t \vdash t = s \quad \frac{\frac{}{s = t \vdash s = t} \quad \frac{}{\vdash s = s}}{s = t \vdash t = s} \quad (\text{E}_=)$$

$$s = t, t = u \vdash s = u$$

Examples

$$s = t \vdash t = s \quad \frac{\frac{}{s = t \vdash s = t} \quad \frac{}{\vdash s = s}}{s = t \vdash t = s} \quad (\text{E}_=)$$

$$s = t, t = u \vdash s = u \quad \frac{\frac{}{s = t \vdash s = t} \quad \frac{}{t = u \vdash t = u}}{s = t, t = u \vdash s = u} \quad (\text{E}_=)$$

Examples

$$s = t \vdash t = s \quad \frac{\frac{}{s = t \vdash s = t} \quad \frac{}{\vdash s = s}}{s = t \vdash t = s} \quad (\text{E}_=)$$

$$s = t, t = u \vdash s = u \quad \frac{\frac{}{s = t \vdash s = t} \quad \frac{}{t = u \vdash t = u}}{s = t, t = u \vdash s = u} \quad (\text{E}_=)$$

$$\exists x \forall y R(x, y) \vdash \forall y \exists x R(x, y)$$

Examples

$$s = t \vdash t = s \quad \frac{\frac{}{s = t \vdash s = t} \quad \frac{}{\vdash s = s}}{s = t \vdash t = s} \quad (E_=)$$

$$s = t, t = u \vdash s = u \quad \frac{\frac{}{s = t \vdash s = t} \quad \frac{}{t = u \vdash t = u}}{s = t, t = u \vdash s = u} \quad (E_=)$$

$$\begin{array}{l} \exists x \forall y R(x, y) \vdash \forall y \exists x R(x, y) \\ \frac{\forall y R(c, y) \vdash \forall y R(c, y)}{\forall y R(c, y) \vdash R(c, d)} \\ \frac{\forall y R(c, y) \vdash \exists x R(x, d)}{\forall y R(c, y) \vdash \forall y \exists x R(x, y)} \\ \frac{\exists x \forall y R(x, y) \vdash \exists x \forall y R(x, y) \quad \forall y R(c, y) \vdash \forall y \exists x R(x, y)}{\exists x \forall y R(x, y) \vdash \forall y \exists x R(x, y)} \end{array} \begin{array}{l} (E_{\forall}) \\ (I_{\exists}) \\ (I_{\forall}) \\ (E_{\exists}) \end{array}$$

Soundness and Completeness

Theorem

A formula φ is provable using Natural Deduction if, and only if, it is valid.

Corollary

The set of valid first-order formulae is recursively enumerable.

Isabelle/HOL

Isabelle/HOL

Proof assistant designed for software verification.

General structure

```
theory T
imports T1 ... Tn
begin
  declarations, definitions, and proofs
end
```

Syntax

Two levels:

- ▶ the **meta-language** (Isabelle) used to define theories,
- ▶ the **logical language** (HOL) used to write formulae.

To distinguish the levels, one encloses formulae of the logical language in quotes.

```
datatype 'a list = Nil                (" [] ")
                  | Cons 'a "'a list" (infixr "#" 65)
```

```
primrec app :: "'a list => 'a list => 'a list"
          (infixr "@" 65)
```

where

```
"[] @ ys      = ys" |
"(x # xs) @ ys = x # (xs @ ys)"
```

Logical Language

Types

- ▶ **base types:** bool, nat, int,...
- ▶ **type constructors:** α list, α set,...
- ▶ **function types:** $\alpha \Rightarrow \beta$
- ▶ **type variables:** 'a, 'b,...

Terms

- ▶ **application:** $f\ x\ y$, $x + y$,...
- ▶ **abstraction:** $\lambda x.t$
- ▶ **type annotation:** $t :: \alpha$
- ▶ **if** b **then** t **else** u
- ▶ **let** $x = t$ **in** u
- ▶ **case** x **of** $p_0 \Rightarrow t_0 \mid \dots \mid p_n \Rightarrow t_n$

Formulae

- ▶ terms of type bool
- ▶ boolean operations \neg , \wedge , \vee , \rightarrow
- ▶ quantifiers $\forall x$, $\exists x$
- ▶ predicates $=$, $<$,...

Basic Types

```
datatype bool = True | False
```

```
fun conj :: "bool => bool => bool" where  
"conj True True = True" |  
"conj _ _ = False"
```

```
datatype nat = 0 | Suc nat
```

```
fun add :: "nat => nat => nat" where  
"add 0 n = n" |  
"add (Suc m) n = Suc (add m n)"
```

```
lemma add_02: "add m 0 = m"  
apply (induction m)  
apply (auto)  
done
```


Proofs

```
lemma add_02: "add m 0 = m"
```

Proofs

```
lemma add_02: "add m 0 = m"
```

```
apply (induction m)
```

Proofs

```
lemma add_02: "add m 0 = m"
```

```
apply (induction m)
```

```
1. add 0 0 = 0
```

```
2.  $\wedge m. \text{add } m \ 0 = m \implies \text{add } (\text{Suc } m) \ 0 = \text{Suc } m$ 
```

Proofs

```
lemma add_02: "add m 0 = m"
```

```
apply (induction m)
```

```
1. add 0 0 = 0
```

```
2.  $\wedge m. \text{add } m \ 0 = m \implies \text{add } (\text{Suc } m) \ 0 = \text{Suc } m$ 
```

```
apply (auto)
```

```
datatype 'a list = Nil                ("[]")
                  | Cons 'a "'a list" (infixr "#" 65)
```

```
fun app :: "'a list => 'a list => 'a list"
        (infixr "@" 65)
```

where

```
"[] @ ys      = ys" |
"(x # xs) @ ys = x # (xs @ ys)"
```

```
fun rev :: "'a list => 'a list" where
```

```
"rev []      = []" |
"rev (x # xs) = (rev xs) @ (x # [])"
```

```
theorem rev_rev [simp]: "rev (rev xs) = xs"
```

```
theorem rev_rev [simp]: "rev (rev xs) = xs"  
apply(induction xs)
```

```
theorem rev_rev [simp]: "rev (rev xs) = xs"
```

```
apply(induction xs)
```

```
1. rev (rev Nil) = Nil
```

```
2.  $\wedge x1 xs. \text{rev (rev xs) = xs} ==>$ 
```

```
   rev (rev (Cons x1 xs)) = Cons x1 xs
```



```
theorem rev_rev [simp]: "rev (rev xs) = xs"
```

```
apply(induction xs)
```

```
1. rev (rev Nil) = Nil
```

```
2.  $\wedge x1\ xs. \text{rev (rev xs) = xs} \implies$ 
```

```
   rev (rev (Cons x1 xs)) = Cons x1 xs
```

```
apply(auto)
```

```
theorem rev_rev [simp]: "rev (rev xs) = xs"
```

```
apply(induction xs)
```

```
1. rev (rev Nil) = Nil
```

```
2.  $\wedge x1\ xs. \text{rev (rev xs)} = \text{xs} \implies$ 
```

```
   rev (rev (Cons x1 xs)) = Cons x1 xs
```

```
apply(auto)
```

```
1.  $\wedge x1\ xs.$ 
```

```
   rev (rev xs) = xs  $\implies$ 
```

```
   rev (rev xs @ Cons x1 Nil) = Cons x1 xs
```

```
lemma app_Nil2 [simp]: "xs @ Nil = xs"  
apply(induction xs)  
apply(auto)  
done
```

```
lemma app_Nil2 [simp]: "xs @ Nil = xs"  
apply(induction xs)  
apply(auto)  
done
```

```
lemma rev_app [simp]: "rev (xs @ ys) = rev ys @ rev xs"  
apply(induction xs)  
apply(auto)
```

1. $\wedge x1$ xs.

$$\begin{aligned} \text{rev (xs @ ys)} &= \text{rev ys @ rev xs} ==> \\ (\text{rev ys @ rev xs}) @ \text{Cons } x1 \text{ Nil} &= \\ \text{rev ys @ (rev xs @ Cons } x1 \text{ Nil)} \end{aligned}$$

```
lemma app_Nil2 [simp]: "xs @ Nil = xs"
apply(induction xs)
apply(auto)
done
```

```
lemma rev_app [simp]: "rev (xs @ ys) = rev ys @ rev xs"
apply(induction xs)
apply(auto)
```

1. $\wedge x1$ xs.

$$\begin{aligned} \text{rev (xs @ ys)} &= \text{rev ys @ rev xs} ==> \\ (\text{rev ys @ rev xs}) @ \text{Cons } x1 \text{ Nil} &= \\ \text{rev ys @ (rev xs @ Cons } x1 \text{ Nil)} \end{aligned}$$

```
lemma app_assoc [simp]: "(xs @ ys) @ zs = xs @ (ys @ zs)"
apply (induction xs)
apply (auto)
done
```

```
lemma app_Nil2 [simp]: "xs @ [] = xs"  
apply(induct_tac xs)  
apply(auto)  
done
```

```
lemma app_assoc [simp]: "(xs @ ys) @ zs = xs @ (ys @ zs)"  
apply(induct_tac xs)  
apply(auto)  
done
```

```
lemma rev_app [simp]: "rev(xs @ ys) = (rev ys) @ (rev xs)"  
apply(induct_tac xs)  
apply(auto)  
done
```

```
theorem rev_rev [simp]: "rev(rev xs) = xs"  
apply(induct_tac xs)  
apply(auto)  
done
```

Nonmonotonic Logic

Negation as Failure

Goal

Develop a proof calculus supporting Negation as Failure as used in Prolog.

Monotonicity

Ordinary deduction is **monotone**: if we add new assumption, all consequences we have already derived remain. More information does not invalidate already made deductions.

Non-Monotonicity

Negation as Failure is **non-monotone**:

P implies $\neg Q$ but P, Q does not imply $\neg Q$.

Default Logic

Rule

$$\frac{\alpha_0 \dots \alpha_m : \beta_0 \dots \beta_n}{\gamma}$$

α_i	assumptions
β_i	restraints
γ	consequence

Derive γ provided that we can derive $\alpha_0, \dots, \alpha_m$, but none of β_0, \dots, β_n .

Example

$$\frac{\text{bird}(x) : \text{penguin}(x) \text{ ostrich}(x)}{\text{can_fly}(x)}$$

Semantics

Definition

A set Φ of formulae is **consistent** with respect to a set of rules R if, for every rule

$$\frac{\alpha_0 \dots \alpha_m : \beta_0 \dots \beta_n}{\gamma} \in R$$

such that $\alpha_0, \dots, \alpha_m \in \Phi$ and $\beta_0, \dots, \beta_n \notin \Phi$, we have $\gamma \in \Phi$.

Note

If there are no restraints β_i , consistent sets are **closed under intersection**.

\Rightarrow There is a unique smallest such set, that of all **provable** formulae.

If there are restraints, this may not be the case. Formulae that belong to all consistent sets are called **secured consequences**.

Examples

The system

$$\frac{}{\alpha} \quad \frac{\alpha : \beta}{\beta}$$

has a unique consistent set $\{\alpha, \beta\}$.

The system

$$\frac{}{\alpha} \quad \frac{\alpha : \beta}{\gamma} \quad \frac{\alpha : \gamma}{\beta}$$

has consistent sets

$$\{\alpha, \beta\}, \quad \{\alpha, \gamma\}, \quad \{\alpha, \beta, \gamma\}.$$