

**Exercise 1** Let  $f$  be a binary function symbol,  $g, h$  unary, and  $c$  a constant symbol.

(a) Find the most general unifier for the following pairs of terms.

- (i)  $f(g(x), y)$  and  $f(x, h(y))$
- (ii)  $f(h(x), x)$  and  $f(x, h(y))$
- (iii)  $f(x, f(x, g(y)))$  and  $f(y, f(h(c), x))$
- (iv)  $f(f(x, c), g(f(y, x)))$  and  $f(x, g(x))$

(b) Solve the following set of term equations

$$x = f(y, z), \quad y = g(u), \quad z = h(y), \quad u = f(v, w), \quad v = f(c, w).$$

**Exercise 2** Consider the following formulae.

- (a)  $\exists x \exists y \forall z [z = x \vee z = y]$
- (b)  $\forall x [\exists y R(x, y) \rightarrow \exists y R(y, x)]$
- (c)  $\forall x [\forall y \exists z [R(x, f(y, z))] \rightarrow \forall y \forall z [R(f(x, y), f(x, z)) \vee R(y, z)]]$
- (d)  $\exists x \forall y R(x, y) \wedge \forall x \exists y R(x, y) \wedge \forall x \forall y [R(x, y) \rightarrow \exists z [R(x, z) \wedge R(z, x)]]$

For each of them

- (1) transform it into Skolem normal form;
- (2) transform it into a set of clauses.

**Exercise 3** Use the resolution method to check that the following formulae are inconsistent.

- (a)  $\forall x \forall y [x \leq y \rightarrow (P(x) \leftrightarrow P(y))] \wedge \forall x \forall y [x \leq y \vee y \leq x] \wedge \exists x P(x) \wedge \exists x \neg P(x)$
- (b)  $\forall x \exists y [y \leq x \wedge \neg E(x, y)] \wedge \forall x \forall y [x \leq y \wedge y \leq x \rightarrow E(x, y)] \wedge \exists x \forall y [x \leq y]$
- (c)  $\forall x \forall y [R(x, y) \rightarrow (P(x) \leftrightarrow \neg P(y))] \wedge \forall x \forall y [R(x, y) \rightarrow \exists z [R(x, z) \wedge R(z, y)]] \wedge \exists x \exists y R(x, y)$
- (d)  $\forall x R(x, f(x)) \wedge \forall x \forall y \forall z [R(x, y) \wedge R(y, z) \rightarrow R(x, z)] \wedge \forall x \forall y [E(x, y) \rightarrow \neg R(x, y)]$   
 $\wedge \exists x E(x, f(f(x)))$

**Exercise 4** Use SLD resolution to check that the following set of Horn-formulae is inconsistent.

- (a)  $\forall x T(x, x)$ ,  
 $\forall x \forall y \forall z [E(x, y) \wedge T(y, z) \rightarrow T(x, z)]$ ,  
 $E(a, b)$ ,  
 $E(b, c)$ ,  
 $E(c, d)$ ,  
 $\neg T(a, d)$ .
- (b)  $\forall x T(x, x)$ ,  
 $\forall x \forall y \forall z [T(x, y) \wedge E(y, z) \rightarrow T(x, z)]$ ,  
 $E(a, b)$ ,  
 $E(b, c)$ ,  
 $E(c, d)$ ,  
 $\neg T(a, d)$ .
- (c)  $R(c, x, x)$ ,  
 $\forall x \forall y \forall z \forall w [R(x, f(y, z), w) \rightarrow R(f(y, x), z, w)]$ ,  
 $\neg \forall x \forall y [R(f(x, f(y, c)), c, f(y, f(x, c)))]$ .