

## **First Aid**

### **Bleeding – External**


Mgr. Zdeňka Kubíková Bc. Barbora Zuchová and others

## External bleeding


#### TREATMENT OF PROFUSE BLEEDING

- 1. First aid (pressure) bandage *or in extreme situation*
- 2. Apply pressure directly on to the wound
- 3. Elevate the affected part
- 4. Lay the casualty down
- 5. Utmost application of tourniquet

Be aware of your own safety!!!



# First aid (pressure) bandage



### Treatment of superficial wounds – excoriations

#### First aid:

- clean with water, use soap, brush
- disinfection



Kelnarová, 2007. 1.díl

### Beware of disinfectant containing iodine!!

sterile wound dressing

### Foreign bodies in the wounds

- don't pull out the foreign body – it may be acting as a partial plug, the foreign body averting larger loss of blood
- place sterile dressing either side
- still the patient
- eventually set against sliding

### Dialling of Emergency Medical Service 155



FOTO: Turin, 2008


## Nose, mouth and ear bleeding

- nosebleeds the head forward pinch the casualty s nose just beneath the bridge
- mouthbleeds the head forward when tooth is knocked out of socket, bite a tampon

 earbleeds – do no tamponade let the blood leak out loose







### ...and sport sometimes hurts

Treatment of external bleeding...

and always watch for other symptoms!

Consciousness of the patient, pain, pulse...

Give every detail you know to the ambulance crew!



MAFA, 2007