

Review article

Developmental programming by maternal obesity in 2015: Outcomes, mechanisms, and potential interventions



Naomi C. Penfold *, Susan E. Ozanne

University of Cambridge, Metabolic Research Laboratories MRC Metabolic Diseases Unit, Wellcome Trust-MRC Institute of Metabolic Science, Addenbrooke's Hospital, Cambridge CB2 0QQ, United Kingdom

ARTICLE INFO

Available online 2 July 2015

Keywords:

Developmental programming
Maternal obesity
Leptin
Insulin
Ghrelin
Appetite
Reward
Glucose homeostasis
Intervention
Obesity

ABSTRACT

This article is part of a Special Issue "SBN 2014".

Obesity in women of child-bearing age is a growing problem in developed and developing countries. Evidence from human studies indicates that maternal BMI correlates with offspring adiposity from an early age and predisposes to metabolic disease in later life. Thus the early life environment is an attractive target for intervention to improve public health. Animal models have been used to investigate the specific physiological outcomes and mechanisms of developmental programming that result from exposure to maternal obesity in utero. From this research, targeted intervention strategies can be designed. In this review we summarise recent progress in this field, with a focus on cardiometabolic disease and central control of appetite and behaviour. We highlight key factors that may mediate programming by maternal obesity, including leptin, insulin, and ghrelin. Finally, we explore potential lifestyle and pharmacological interventions in humans and the current state of evidence from animal models.

© 2015 Elsevier Inc. All rights reserved.

Contents

Introduction	143
Animal models have revealed mechanisms underlying programming by maternal obesity	144
Insulin and glucose homeostasis	144
Cardiovascular system	145
Ectopic lipid deposition	145
Central control of food intake: programming the hypothalamus	145
Maternal obesity predisposes to diet-induced obesity: the role of the reward system	146
Programming learning and memory: leptin and the hippocampus	146
Candidate programming mechanisms and factors in maternal obesity	147
Interventions to improve outcomes of offspring exposed to maternal obesity	148
Conclusion	148
Acknowledgments	148
References	148

Introduction

The importance of normal foetal growth was first highlighted by associations between low birth weight and the increased risk of heart disease and type 2 diabetes in adulthood (Barker et al., 1989; Hales et al., 1991). Subsequent studies of maternal under-nutrition and,

more recently, maternal over-nutrition have demonstrated that the maternal nutritional environment and foetal and neonatal growth, collectively known as the first 1000 days of life, are key determinants of health in the next generation (de Rooij et al., 2006; Lumey and Stein, 1997; Ravelli et al., 1976, 1999). In humans, maternal obesity is associated with low and high birth weight (Cedergren, 2004; Gaudet et al., 2014) and increased risk of obesity and metabolic dysfunction in the offspring both in childhood (Boney et al., 2005; Whitaker, 2004) as well as in adulthood (Brisbois et al., 2012; Cooper et al., 2010). Maternal

* Corresponding author.
E-mail address: np325@medschl.cam.ac.uk (N.C. Penfold).

obesity is also associated with increased risk of offspring cardiovascular disease (Drake and Reynolds, 2010), type 2 diabetes (Berends and Ozanne, 2012), and neurodevelopmental and psychiatric disorders, including ADHD, autism, schizophrenia, and mood disorders (Mehta et al., 2014; Rodriguez, 2010).

The prevalence of overweight and obesity has soared in the last 30 years globally (Ng et al., 2014). Worryingly, the number of children classified as overweight or obese has increased 150% worldwide in this timeframe (Ng et al., 2014) and the rate of obesity in women of child-bearing age is still rising (Fisher et al., 2013). Whilst genetic factors that predispose to obesity in an obesogenic environment, have likely contributed to the current global obesity epidemic, the short time-scale of this increase implicates non-genetic factors including the impact of the intrauterine and neonatal environment on adult health and disease (McAllister et al., 2009). It is vital that we understand the mechanisms underlying such developmental programming of disease by maternal obesity in order to develop effective interventions to help mitigate the current rise in obesity, cardiovascular and metabolic diseases as well as mental health disorders. Bariatric surgery to induce weight loss lowers the risk of gestational diabetes mellitus (GDM), foetal macrosomia and the rate of obesity in the offspring as well as improving offspring insulin sensitivity, demonstrating that improving the maternal metabolic state prior to pregnancy is an effective intervention that improves the health of both mother and child (Kral et al., 2006; Shai et al., 2014; Smith et al., 2009). However, bariatric surgery is intrusive, high-risk, costly and can cause nutrient deficiency, the latter of which led to severe neural defects in some children conceived very soon after surgery (Pelizzo et al., 2014). A clearer understanding of the mechanisms mediating the increased risk of metabolic disease in offspring of obese women is required in order to develop less intrusive, better targeted interventions. This review will explore recent progress made in the understanding of the developmental programming by maternal obesity and potential avenues for intervention.

Animal models have revealed mechanisms underlying programming by maternal obesity

Animal studies have confirmed that maternal obesity programs metabolic syndrome—like outcomes in the offspring including impaired insulin action and glucose homeostasis (Martin-Gronert et al., 2010; Samuelsson et al., 2008; Shankar et al., 2010; Shelley et al., 2009), hypertension, and cardiovascular dysfunction (Blackmore et al., 2014; Fernandez-Twinn et al., 2012; Samuelsson et al., 2008), as well as increased adiposity (Bayol et al., 2008; Samuelsson et al., 2008; Song et al., 2015), and an increased susceptibility to diet-induced obesity (DIO) (Bayol et al., 2007; Howie et al., 2009; Kirk et al., 2009; Nivoit et al., 2009; Samuelsson et al., 2008; Shankar et al., 2008; Torrens et al., 2012). The choice of animal model is often a compromise between practicality of the research and translatability to humans. Whilst non-human primates (NHPs) share the closest resemblance to human developmental trajectories and pregnancies, they have a long gestation length and time to maturity of the offspring, leading to high research costs. Sheep and pigs are used due to their similarities in placental structure and function to humans, whilst rabbits are a medium-sized mammal with intermediary similarities and differences to humans. These larger mammals are conducive to repeated sampling of blood and tissue, allowing for longitudinal studies and within-subject analysis. Models with larger litter sizes, such as pigs and rodents, allow for the easier investigation of sex differences in programming. Rodent models have been used extensively due to their short gestation (three weeks) and maturity intervals (five weeks to puberty) and the ease with which to generate a well-powered experiment of animals of ages across the lifecourse. Furthermore, they enable genetic engineering to elucidate mechanisms. A disadvantage is that these smaller mammals are limited to one sampling point, precluding true longitudinal analysis. In addition, there are several differences in developmental timings of key tissues

between rodents and humans. An overarching observation is that the third trimester in humans is roughly equivalent to the first postnatal weeks in the rodent. Notably adipose tissue develops from early in gestation in humans whereas subcutaneous and visceral depots develop from late gestation and early postnatal life, respectively, in rodents (Rosen and Spiegelman, 2014). Cardiomyocyte proliferation and growth is mostly complete by birth in the human and sheep (Morrison et al., 2007), whereas cardiomyocyte division ends at postnatal day 3 to 4 in the rat, with growth occurring over the first two weeks of life (Li et al., 1996). In addition, the development of key intra-hypothalamic connections occurs during the second postnatal week in rodents but these connections are established by birth in humans and NHPs (Bouret, 2012; Coupe and Bouret, 2013; Liu et al., 2013). The choice of animal model will affect the translatability of the results, however the outcomes seen in these models often recapitulate phenotypes reported in humans, signifying the validity of the use of a range of animals to investigate the mechanisms underlying developmental programming.

Insulin and glucose homeostasis

Maternal obesity programs offspring adiposity, decreased glucose tolerance, and impaired insulin sensitivity (Fernandez-Twinn et al., 2012; Samuelsson et al., 2008; Yan et al., 2011). The mechanisms underlying programming of insulin resistance and glucose homeostasis by maternal obesity include alterations in peripheral insulin signalling and insulin secretion [reviewed in (Berends and Ozanne, 2012) and (Duque-Guimaraes and Ozanne, 2013)]. Adult offspring exposed to maternal obesity are hyperinsulinaemic and have alterations in the expression of key insulin signalling and glucose handling molecules in skeletal muscle, the liver and adipose tissue that indicate a predisposition for insulin resistance and impaired glucose tolerance (Martin-Gronert et al., 2010; Nicholas et al., 2013a; Rattanaray et al., 2010; Shelley et al., 2009; Yan et al., 2011). At least some of the programming of insulin signalling protein expression appears to occur through post-transcriptional mechanisms via changes in microRNA (miR-) levels. Maternal obesity at conception in sheep increases hepatic miR-29b, miR-130, and miR-107 levels (Nicholas et al., 2013b). Increased miR-126 expression in adipose tissue of mice exposed to maternal obesity is associated with down-regulated expression of target genes involved in insulin signalling including insulin receptor substrate 1 (IRS-1) (Fernandez-Twinn et al., 2014). These programmed changes in IRS-1 and miR-126 were maintained following differentiation of pre-adipocytes in vitro, indicating that maternal obesity programs altered insulin signalling in the offspring adipose tissue in a cell-autonomous fashion.

In addition to peripheral insulin signalling, recent evidence suggests that the central control of glucose homeostasis is vulnerable to the hyperinsulinaemic obese maternal environment. Genetically-induced maternal hyperinsulinaemia and insulin resistance is associated with disrupted glucose homeostasis and hyperinsulinaemia in male wild-type offspring despite normal body weight and glycaemia in the mother (Isganaitis et al., 2014). Furthermore, a recent study demonstrated that genetic abrogation of insulin signalling specifically in pro-opiomelanocortin (POMC) neurons of offspring exposed to a maternal high-fat diet (HFD) restores POMC innervation of pre-autonomic paraventricular nucleus (PVH) neurons and normalises the impaired glucose tolerance otherwise seen (Vogt et al., 2014). This is associated with an improvement in pancreatic beta cell glucose-stimulated insulin secretion and parasympathetic innervation of beta cells.

Maternal hyperinsulinaemia with insulin resistance might program altered offspring development via the concomitant maternal hyperglycaemia, since insulin does not cross the placenta whereas glucose does (Dabelea, 2007). In humans, impaired glucose tolerance during pregnancy is often associated with increased birth weight and increased risk of childhood obesity (Catalano et al., 2003; Cottrell and Ozanne, 2007; Hillier et al., 2007; Liu et al., 2014; Plagemann et al.,

2002). Treating GDM mothers to lower their blood glucose reduces this risk, particularly in male offspring (Bahado-Singh et al., 2012; Gillman et al., 2010). In a recent study in mice, genetically-induced maternal hyperglycaemia is associated with increased body weight and impaired glucose tolerance in wild-type male offspring (Nadif et al., 2015). Therefore control of glycaemia during pregnancy is not only important for maternal health but also for the long term health of the offspring.

Cardiovascular system

Hypertension and cardiac hypertrophy are common phenotypes observed in offspring exposed to maternal obesity (Fernandez-Twinn et al., 2012; Samuelsson et al., 2008). Studies in rabbits and rats have suggested that changes in sympathetic tone may be an important mediator of these effects. Maternal HFD in rabbits increases renal sympathetic nerve activity in the offspring (Prior et al., 2014). Likewise studies suggest that maternal obesity in rats induces hypertension in the offspring via increased sympathetic drive in early development (Samuelsson et al., 2010), which may be mediated by altered early life leptin signalling.

Leptin action in the nucleus of the solitary tract (NTS) and the ventromedial nucleus of the hypothalamus (VMH) increases sympathetic outflow via the renal nerve (Li et al., 2013; Mark et al., 2009; Marsh et al., 2003). Umbilical cord leptin levels are elevated in obese pregnancies (Ferretti et al., 2014; Karakosta et al., 2013; Walsh et al., 2014) and neonatal circulating leptin is elevated in offspring of obese mice (Samuelsson et al., 2008). Therefore, early life hyperleptinaemia may drive sympathetic hyperstimulation in the developing renal-cardiovascular system, leading to hypertension and cardiovascular dysfunction in adulthood (Briffa et al., 2014). Indeed, neonatal leptin administration in rats results in cardiac hypertrophy and dysfunction in adulthood (Marques et al., 2014b). In addition, rat offspring exposed to maternal obesity show an exaggerated hypertensive response to peripheral leptin administration in adulthood (Samuelsson et al., 2010). This is unlikely to be due to impaired central leptin signalling, as maternal obesity-mediated programming of leptin resistance is hypothalamic nuclei-specific (Kirk et al., 2009) and diet-induced obesity in adulthood does not impair central leptin-mediated sympathetic activity via the renal nerve (Rahmouni et al., 2005). Therefore it has been postulated that the hyperleptinaemia seen in adult offspring of maternal obesity animal models drives the accompanying hypertension via the concomitant increase in central activation of the sympathetic nervous system (Samuelsson et al., 2010; Simonds et al., 2014). Notably, it has recently been shown that the increased risk of hypertension in obese individuals is dependent on functional leptin signalling (Simonds et al., 2014).

Our studies in a mouse model of maternal DIO have shown that male offspring of obese dams display cardiac hypertrophy associated with hyperinsulinaemia and increased oxidative stress prior to a change in body weight or adiposity, indicating that the programming of increased risk of cardiovascular disease is independent from mechanisms relating to obesity (Blackmore et al., 2014; Fernandez-Twinn et al., 2012). Furthermore, frank cardiac dysfunction with increased sympathetic dominance akin to the early stages of heart failure is evident in these mice by young adulthood (Blackmore et al., 2014). This dysfunction may relate to pathological cardiac hypertrophy and cardiac stress as early as weaning. Oxidative stress, inflammation and epigenetic mechanisms may all be involved in the programming of cardiovascular dysfunction by maternal obesity (Blackmore and Ozanne, 2013, 2014). Given that obesity itself increases the risk of heart disease, cardiac dysfunction may be exaggerated in high-fat-fed offspring exposed to maternal obesity. Indeed, the combination of maternal HFD and post-weaning exposure to HFD culminates in reduced vasorelaxation in both mice and non-human primates (Fan et al., 2013; Torrens et al., 2012), with increased oxidative stress in the femoral arteries of adult male offspring (Torrens et al., 2012).

In summary, early life exposure to hyperleptinaemia as a consequence of maternal obesity may drive increased sympathetic tone leading to hypertension and accelerate the onset of cardiac hypertrophy and heart failure.

Ectopic lipid deposition

Maternal obesity programs increased adiposity and adipose tissue function in the offspring via alterations in adipocyte morphology and signalling (Alfaradhi and Ozanne, 2011; Benkalfat et al., 2011; Murabayashi et al., 2013) as well as changes in food intake. As well as increased adiposity, ectopic lipid deposition has also been observed in the liver and pancreas of offspring exposed to maternal obesity (Alfaradhi et al., 2014; Oben et al., 2010a,b), in association with altered hepatic mRNA and protein expression profiles indicative of increased lipogenesis (Bruce et al., 2009), elevated markers of oxidative damage (Alfaradhi et al., 2014; Bringham et al., 2014; Torrens et al., 2012), inflammation, fibrosis and increased sympathetic nervous system activation (Oben et al., 2010a). These results provide evidence for an increased risk of non-alcoholic fatty liver and pancreas diseases (NAFLD and NAFPD, respectively) in offspring of obese mothers, a pathology which commonly occurs in obesity when the normal capacity of white adipose tissue for lipid storage has been exceeded. Recent evidence from a mouse model of maternal DIO suggests that the predisposition for NAFPD in high-fat-fed offspring is associated with a programmed shift in the cellular circadian clock (Carter et al., 2014). Perturbation in internal biological rhythms is a recent addition to the list of offspring physiologies affected by maternal nutrition (Martin-Gronert and Ozanne, 2013) and represents an exciting avenue for investigation, given the new understanding of circadian biology in health and disease (Bailey et al., 2014).

Interestingly, recent evidence from a swine model of maternal obesity suggests that increased risk of liver disease can be programmed transgenerationally, since early postnatal increases in adiposity and markers of paediatric liver disease are found in male piglets of obese grandmothers (Gonzalez-Bulnes et al., 2014).

Central control of food intake: programming the hypothalamus

The increased incidence of offspring obesity is frequently associated with hyperphagia in maternal over-nutrition models (Bayol et al., 2007; Kirk et al., 2009; Long et al., 2011; Nivoit et al., 2009; Samuelsson et al., 2008). This increased caloric intake is accompanied by alterations in hypothalamic expression of key neuropeptides, their receptors and molecules involved in signalling by peripheral factors (Chen and Morris, 2009; Chen et al., 2009; Ferezou-Viala et al., 2007; Gupta et al., 2009; Morris and Chen, 2009; Page et al., 2009) as well as altered hypothalamic development (Chang et al., 2008; Kirk et al., 2009). Alterations in gene expression may be due to epigenetic alterations such as changes in DNA methylation within the gene promoters, as observed in offspring exposed to early life over-nutrition (Plagemann et al., 2009, 2010). The impaired development of hypothalamic circuitry in rodents is likely due to alterations in the hormonal environment in early postnatal life. Insulin has been implicated in the programming of hypothalamic circuits in response to maternal diabetes and maternal over-nutrition (Steculorum et al., 2013; Vogt et al., 2014). Maternal hypoinsulinaemic hyperglycaemia increases the ratio of orexigenic neurons to anorexigenic neurons in the neonatal arcuate nucleus (Arc) (Franke et al., 2005; Steculorum and Bouret, 2011b) and impairs Arc-PVH Agouti-related peptide (AgRP) and POMC projections (Steculorum and Bouret, 2011b). These changes are associated with elevated circulating glucose, insulin and leptin in the neonate and central leptin resistance, hyperphagia and obesity in adult life (Steculorum and Bouret, 2011b). In addition, maternal over-nutrition can alter the timing, amplitude of, and response to the postnatal surge in neonatal leptin concentrations that is critical for the development of hypothalamic circuitry (Ahima et al., 1998; Bouret et al., 2004a; Long et al., 2011; Toste et al., 2006).

Leptin promotes neurite outgrowth from the Arc during neonatal life (Bouret et al., 2004b, 2012) and abnormal neonatal leptin signalling impairs the formation of the Arc-derived hypothalamic projections (Attig et al., 2008; Delahaye et al., 2008; Yura et al., 2005). It has recently emerged that ghrelin also contributes to the early life programming of obesity. Neonatal ghrelin administration increases Arc neuronal number and increases the ratio of orexigenic to anorexigenic gene expression (Steculorum and Bouret, 2011a). Chronic postnatal ghrelin impairs the formation of Arc projections in association with metabolic dysfunction and impaired leptin sensitivity in adulthood (Steculorum et al., 2015). Neonatal over-nutrition, by reducing litter size and thus increasing access to the mother's milk, predisposes offspring to hyperphagia and obesity in adulthood (Colliden et al., 2015; Plagemann et al., 1999). This is associated with decreased serum ghrelin in neonates, due to a loss of the normal up-regulation of ghrelin mRNA in the neonatal stomach, and with abrogation of ghrelin-induced gene expression in the Arc, potentially due to impaired transport of ghrelin into the ventromedial hypothalamus (Colliden et al., 2015). Impairment of central ghrelin action in neonates increases Arc projection density and leads to obesity, hyperglycaemia and impaired leptin sensitivity in adulthood (Steculorum et al., 2015).

Thus alteration of central insulin, leptin, and ghrelin signalling in neonates exposed to maternal obesity, with insulin resistance, hyperglycaemia and hyperleptinaemia, may underlie the programming of altered hypothalamic development and subsequent metabolic dysfunction in the adult offspring.

Maternal obesity predisposes to diet-induced obesity: the role of the reward system

Studies in rodents have demonstrated that offspring exposed to maternal obesity and/or HFD during gestation and lactation are predisposed to a greater increase in adiposity and metabolic dysregulation than those from control dams when the offspring themselves are challenged with a HFD after weaning (Benkalfat et al., 2011; Howie et al., 2009; Page et al., 2009; Parente et al., 2008; Rajia et al., 2010). Post-weaning exposure to a HFD further alters hypothalamic mRNA and protein expression (Page et al., 2009; Rajia et al., 2010), which may mediate the increased caloric intake and drive the increased adiposity. Alternatively, the increased propensity for DIO in offspring of obese mothers may be mediated via programmed dysregulation of the central mechanisms involved in palatable food intake: namely the mesocorticolimbic dopamine pathway from the ventral tegmental area (VTA) to the nucleus accumbens (NAcc). Dopaminergic signalling in the NAcc is thought to control incentive salience, or the motivated “wanting” of palatable foods, whilst opioidergic inputs onto the same pathway are thought to signal the pleasure associated with eating tasty foods and so influence food preferences or the “liking” of palatable foods (Blum et al., 2012; Egecioglu et al., 2011; Volkow et al., 2011). Connections between the reward system and the hypothalamus are critical for the regulation of reward-related feeding (Dietrich et al., 2012; Leinninger et al., 2011).

In humans and rodents, reward signalling is altered in obesity (Batterink et al., 2010; Burger and Stice, 2011; Finger et al., 2012; Johnson and Kenny, 2010; Shin and Berthoud, 2011; Stoeckel et al., 2008), due at least in part to chronic HFD-mediated epigenetic dysregulation of key dopaminergic and opioidergic signalling molecules (Vucetic et al., 2011, 2012). In addition, dysregulated reward signalling may predispose to diet-induced obesity (Blum et al., 2014; Volkow et al., 2008). Thus, the central reward system may be vulnerable to early life exposure to maternal obesity and programmed alterations may underlie the increased propensity for obesity when offspring are exposed to a highly palatable diet in adulthood.

Indeed, in animal models of maternal HFD or obesity, offspring consume more high-fat and high-sugar foods than controls (Bayol et al., 2007; Bocarsly et al., 2012; Ong and Muhlhauser, 2011, 2014; Tamashiro et al., 2009; Walker et al., 2008). This may be due to an increased preference for these macronutrients (Vucetic et al., 2010) but

is not associated with altered orosensory stimulation by their taste (Treesukosol et al., 2014). Whilst food preferences can be programmed by maternal nutrition (reviewed in (Gugusheff et al., 2014)), maternal obesity is also associated with altered motivation for palatable foods in multiple rodent models (Grissom et al., 2014b; Naef et al., 2011; Rodriguez et al., 2012). The programmed increases in preference for fat and sugar and altered motivation to work for such foods are associated with changes in dopaminergic tone (Naef et al., 2011, 2013) as well as in expression of key dopaminergic and opioidergic signalling genes (Naef et al., 2011; Ong and Muhlhauser, 2011; Vucetic et al., 2010), with evidence for epigenetic regulation at some loci (Grissom et al., 2014a; Vucetic et al., 2010). In fact, maternal obesity at conception is sufficient to program opioid dysregulation in the offspring (Grissom et al., 2014c). Therefore, maternal obesity may predispose the offspring to DIO via programmed changes in the mesocorticolimbic reward pathway. Importantly, the mesocorticolimbic dopamine pathway develops in utero in rodents with VTA efferents innervating the accumbens and cortex by birth (Hu et al., 2004). Therefore, investigations into the in utero programming of the reward system may more readily translate from mouse to man than for some other systems.

Programming learning and memory: leptin and the hippocampus

Offspring exposed to maternal obesity are slower to acquire an executive function task, in which they demonstrate greater impulsivity but no difference in attention (Grissom et al., 2014b). The hippocampus mediates learning and develops perinatally in both humans and rodents (Semple et al., 2013). In rodents, an important period of synaptogenesis and dendritic spine formation in the developing hippocampus coincides with the peak of the postnatal leptin surge in rodents, which is significant as leptin induces excitatory synaptogenesis and promotes dendritic spine formation in the adult hippocampus (Dhar et al., 2014a,b). Leptin also potentiates GABAergic transmission in postsynaptic CA3 pyramidal cells from the hippocampi of newborn rats (Guimond et al., 2014). The basal activity of these cells is reduced in leptin-deficient mice, as is a marker of presynaptic GABA synthesis, indicating that leptin signalling is critical for GABAergic transmission in the developing hippocampus (Guimond et al., 2014). In addition, chronic leptin treatment during the first two postnatal weeks alters the expression of genes involved in NMDA signalling and synaptic machinery and reduces long-term potentiation in pre-weaning rats (Walker et al., 2007). A similar phenotype is observed in hyperleptinaemic neonates exposed to maternal HFD from late gestation through lactation (Walker et al., 2008). As such, altered leptin signalling in early life may impair the formation of synapses and dendritic spines and thus the maturation of the hippocampus, which may underpin the reported impaired cognition, learning and memory in later life and predisposition for psychopathologies and obesity (Valleau and Sullivan, 2014).

In addition, the programming of obesity and psychiatric disorders by maternal obesity has been attributed to increased maternal-foetal inflammatory signalling (Bolton and Bilbo, 2014; Marques et al., 2014a). It has recently been shown that the impairment in Arc-PVH neuropeptide Y (NPY) projections seen in mice exposed to maternal DIO may be due to increased foetal exposure to the inflammatory cytokine interleukin-6 (IL-6) (Sanders et al., 2014). Maternal IL-6 is also increased mid-gestation in mothers with GDM and inversely correlates with birth weight and glucose tolerance (Hassiakos et al., 2015). In fact, the correlation between GDM and IL-6 levels is so strong that circulating IL-6 alone can predict GDM status. In addition, maternal obesity is associated with increased levels of inflammatory cytokines (Challier et al., 2008; Kępczynska et al., 2013; Kim et al., 2014), of which IL-6 is associated with increased risk of obesity in the offspring (Dahlgren et al., 2001; Smith et al., 2007). Therefore, inflammatory cytokines are also candidate programming mediators in the early life programming of central dysfunction by maternal obesity.

Candidate programming mechanisms and factors in maternal obesity

Potential molecular mediators of the programming of cardiometabolic disease and central neuroendocrine pathways by maternal obesity have been highlighted by recent mechanistic studies. Identification of the key programming factors is vital for the development of rational intervention strategies. It is also important to understand the key windows for intervention: do we aim to intervene before or during pregnancy and/or during early postnatal life? Should interventions target maternal diet, maternal obesity or both?

In utero exposure to maternal obesity is an important target for intervention. It is important to note here the differences in placental biology and developmental timings between rodents, the key model for mechanistic studies, and humans. Rodent placentae structure and blood flow differ from human placentae, however mice have been used successfully to model intra-uterine growth restriction (Gonzalez-Bulnes and Astiz, 2015). Sheep and pig models are more common in investigations into placental biology and intrauterine development, due to their closer resemblance to human placental morphology but also the ability to insert

catheters into the maternal and foetal circulation in order to monitor placental transfer over time in vivo (Barry and Anthony, 2008).

Maternal obesity during pregnancy may impair foetal nutrition via placental adaptations (Tarrade et al., 2015). Indeed placentae from obese women transport less maternal taurine, a critical beta-amino acid involved in placental development and foetal growth (Ditchfield et al., 2014) and have higher levels of oxidative stress and impaired mitochondrial respiration (Hastie and Lappas, 2014; Mele et al., 2014). In addition, maternal DIO in mice is associated with decreased placental mTOR signalling, which may contribute to the decreased foetal:placental weight ratio in late gestation via altered amino acid transport (Lager et al., 2014). Conversely maternal high-fat feeding, whether accompanied by obesity or not, is associated with foetal overgrowth and up-regulation of glucose and amino acid transport across the placenta (Jones et al., 2009; Sferruzzi-Perri et al., 2013). Thus, foetal growth may be altered in maternal obesity due to alterations in placental function.

In addition, altered maternal intake of vital micronutrients in maternal obesity may contribute to offspring epigenetic programming. Dietary intake of key methyl donors varies seasonally in certain populations such




Fig. 1. Maternal obesity programs obesity, cardiometabolic disease and neuropsychiatric disorders in the offspring. Maternal factors involved include hyperinsulinaemia, hyperglycaemia, hyperleptinaemia, hyperlipidaemia, and impaired placental function. Common programming mechanisms in offspring tissues include oxidative stress, epigenetics, and inflammation. Inflammation, insulin, leptin and ghrelin have all been implicated in brain development. The early life programming of brain circuits [HIP – hippocampus, HYP – hypothalamus, ML – mesolimbic pathway] may contribute to altered energy balance, motivated and other behaviours. Altered central control of the autonomic nervous system (ANS) may underlie cardiac and pancreatic phenotypes in the offspring. Programmed changes in adipose tissue, the liver, pancreas, and skeletal muscle function contribute to impaired glucose homeostasis. Overall, alterations in individual tissue function contribute to the increased risk of obesity, cardiometabolic disease, and neuropsychiatric disorder in the offspring. Current strategies aim to ameliorate metabolic status of the obese mother via lifestyle or pharmacological interventions before conception or during pregnancy in order to normalise offspring phenotype.

as those in the Gambia where the timing of pregnancy in relation to the seasons is associated with permanent alterations in DNA methylation at key loci in the offspring (Dominguez-Salas et al., 2014). This provides some of the earliest evidence for the impact of human maternal methyl donor dietary intake during pregnancy on life-long epigenetic programming in the offspring. In rodents, maternal dietary supplementation with methyl donors ameliorates the increased body weight gain in offspring of obese dams (Carlin et al., 2013; Cordero et al., 2014) and restores fat preference to control levels in association with normalisation of the methylation status at promoter regions of key genes involved in the central reward system (Carlin et al., 2013).

The early postnatal life and the lactation period is another target for intervention. Rodents experience fluctuations in hormonal levels during the first three weeks of life that have been implicated in the development and maturation of key hypothalamic circuitry (Bouret, 2013). Whilst this is different to human development, early postnatal life in humans is also considered to be a vital time for the maturation of the brain and adipose tissue. As such, exposure to maternal obesity during lactation is a factor in offspring health, with one potential mediator being alterations in breast milk lipid content. In both humans and rodents, over-nutrition and accelerated growth during the neonatal period is associated with increased adiposity in later life (Plagemann et al., 2012). The combination of maternal obesity and HFD consumption reduces breast milk lipids, whilst HFD consumption during lactation alone increases them (Rolls et al., 1986). Breast milk lipid content is decreased in HFD-fed obese dams during lactation compared to HFD-fed control dams, due to impaired mammary fatty acid synthesis (Saben et al., 2014). In a maternal DIO rat model, breast milk levels of triglycerides are elevated but free fatty acids are decreased early in lactation and increased in the latter stages (Kirk et al., 2009).

As discussed above, maternal obesity during pregnancy and lactation is associated with elevated maternal circulating leptin, insulin, glucose and inflammatory cytokines, all of which have been linked to cardiometabolic dysfunction in the offspring (Fig. 1). Exposure to these maternal factors both in utero and during early postnatal life can alter offspring development. As such, interventions should aim to target women planning to conceive or soon after pregnancy is confirmed. Ensuring appropriate maternal dietary nutrition, improving the metabolic status of obese women in order to normalise hormonal levels, ameliorate inflammation and improve placental sufficiency, and optimizing infant growth and nutrition in the neonatal period are key aims of intervention.

Interventions to improve outcomes of offspring exposed to maternal obesity

Improving women's metabolic health at the time when they are trying to reproduce is an attractive target, since it would benefit the health of both mother and child and only a temporary improvement in maternal health could improve public health for generations. Notably, dietary and lifestyle advice has been shown to be effective in overweight and obese pregnant women (Dodd et al., 2014).

Rodent models of maternal obesity have been used to study the effectiveness of dietary and exercise interventions in the mother on offspring metabolic and behavioural phenotype, due to the ability to enforce exercise and easily control diets in these species. Dietary intervention from before pregnancy or during lactation normalises the increased adiposity and circulating leptin, insulin and triglycerides in weanling offspring, rescues the altered motivation and hyperphagia and partially normalises glucose homeostasis and adipocyte morphology in adulthood (Bayol et al., 2007; Rodriguez et al., 2012; Zambrano et al., 2010). In addition, maternal dietary intervention rescues the increased anxiety and altered social behaviours in female offspring of maternal DIO mice in association with amelioration of central inflammation in these offspring (Kang et al., 2014). However, the same reversal is not seen in male offspring. Voluntary exercise before and during pregnancy in lean dams improves glucose homeostasis in the offspring (Carter et al., 2012, 2013) and prevents hyperleptinaemia (Laker et al., 2014;

Vega et al., 2013). This may be due to the reduction in levels of maternal circulating triglycerides, glucose, insulin, cholesterol, oxidative stress and corticosterone (Vega et al., 2013).

Randomised controlled trials (RCTs) are now being used to investigate whether the same improvements can be seen in obese human pregnancies. A low glycaemic index (GI) diet during pregnancy has been shown to increase weight loss from pre-pregnancy to three months after birth in overweight women and thus may minimise gestational weight gain (Horan et al., 2014). Current RCTs are addressing the effect of exercise alone (Sagedal et al., 2013; Seneviratne et al., 2014) or in combination with dietary intervention (Briley et al., 2014) to improve health outcomes in overweight and obese mothers and their children.

In addition, pharmacological studies are addressing the possibility of normalising the maternal metabolic and hormonal state with a view to improving offspring health. Metformin, an insulin sensitiser, has been trialled as an alternative to insulin treatment for gestational diabetes, with initial results indicating no effect on offspring blood pressure at 2 years of age in comparison to insulin treatment nor in maternal postpartum weight loss when compared to placebo (Battin et al., 2015; Refuerzo et al., 2015). There is currently a trial underway to test whether metformin administration during pregnancy in obese women will prevent macrosomia and this study will include investigation of maternal factors, including insulin resistance, inflammation and adiposity, as well as foetal adiposity (Chiswick et al., 2015). However, concerns have been raised as to the lack of long-term safety data in offspring exposed to metformin during gestation (Fantus, 2015). Animal models are invaluable to help address this issue. An initial study into the effects of metformin administration during pregnancy in a maternal obesity mouse model found that offspring from metformin-treated dams were protected from glucose intolerance and key gene expression changes in skeletal muscle (Tong et al., 2011). A more recently published study suggests that offspring from high-fat fed dams treated with metformin during pregnancy are protected against the exacerbated body weight gain upon exposure to a high fat diet in adulthood (Salomaki et al., 2014). However, this was not a model of maternal obesity and the number of litters studied was low. Additional investigations in rodent models are needed to understand the long-term effects of gestational exposure to metformin and to complement the human trials on short-term outcomes in obese pregnancies.

Further study in animal models is therefore required to inform the most effective timing and intensity of specific dietary and pharmacological interventions, including in altricial species to model intervention periods in line with human developmental timings (Nathanielsz et al., 2013).

Conclusion

In summary, recent animal studies of developmental programming by maternal obesity have advanced our understanding of the underlying mechanisms as well as further elucidated aspects of offspring physiology that contribute to their increased risk of obesity, cardiometabolic disease and mental health disorders. As the focus shifts towards designing interventions to curtail the developmental programming by maternal obesity, studies in both animals and humans are necessary to ensure safety, effectiveness, and specificity.

Acknowledgments

NCP holds a Wellcome Trust PhD studentship; SEO is a member of the MRC Metabolic Diseases Unit and is funded by MRC grant MC_UU_12012/4. We thank Laura Dearden for her helpful discussions.

References

- Ahima, R.S., Prabakaran, D., Flier, J.S., 1998. Postnatal leptin surge and regulation of circadian rhythm of leptin by feeding. Implications for energy homeostasis and neuroendocrine function. *J. Clin. Invest.* 101, 1020–1027.

- Alfaradhi, M.Z., Ozanne, S.E., 2011. Developmental programming in response to maternal overnutrition. *Front. Genet.* 2, 27.
- Alfaradhi, M.Z., Fernandez-Twinn, D.S., Martin-Gronert, M.S., Musial, B., Fowden, A., Ozanne, S.E., 2014. Oxidative stress and altered lipid homeostasis in the programming of offspring fatty liver by maternal obesity. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 307, R26–R34.
- Attig, L., Solomon, G., Ferezou, J., Abdennebi-Najar, L., Taouis, M., Gertler, A., Djiane, J., 2008. Early postnatal leptin blockage leads to a long-term leptin resistance and susceptibility to diet-induced obesity in rats. *Int. J. Obes. (Lond)* 32, 1153–1160.
- Bahado-Singh, R.O., Mele, L., Landon, M.B., Ramin, S.M., Carpenter, M.W., Casey, B., Wapner, R.J., Varner, M.W., Rouse, D.J., Thorp Jr, J.M., Sciscione, A., Catalano, P., Harper, M., Saade, G., Caritis, S.N., Peaceman, A.M., Tolosa, J.E., 2012. Eunice Kennedy Shriver National Institute of Child Health and Human Development Maternal-fetal Medicine Units Network, Fetal male gender and the benefits of treatment of mild gestational diabetes mellitus. *Am. J. Obstet. Gynecol.* 206 (422), e421–e425.
- Bailey, S.M., Udoh, U.S., Young, M.E., 2014. Circadian regulation of metabolism. *J. Endocrinol.* 222, R75–R96.
- Barker, D.J., Winter, P.D., Osmond, C., Margetts, B., Simmonds, S.J., 1989. Weight in infancy and death from ischaemic heart disease. *Lancet* 2, 577–580.
- Barry, J.S., Anthony, R.V., 2008. The pregnant sheep as a model for human pregnancy. *Theriogenology* 69, 55–67.
- Batterink, L., Yokum, S., Stice, E., 2010. Body mass correlates inversely with inhibitory control in response to food among adolescent girls: an fMRI study. *Neuroimage* 52, 1696–1703.
- Battin, M.R., Obolonkin, V., Rush, E., Hague, W., Coat, S., Rowan, J., 2015. Blood pressure measurement at two years in offspring of women randomized to a trial of metformin for GDM: follow up data from the MiG trial. *BMC Pediatr.* 15, 54.
- Bayol, S.A., Farrington, S.J., Stickland, N.C., 2007. A maternal 'junk food' diet in pregnancy and lactation promotes an exacerbated taste for 'junk food' and a greater propensity for obesity in rat offspring. *Br. J. Nutr.* 98, 843–851.
- Bayol, S.A., Simbi, B.H., Bertrand, J.A., Stickland, N.C., 2008. Offspring from mothers fed a 'junk food' diet in pregnancy and lactation exhibit exacerbated adiposity that is more pronounced in females. *J. Physiol. Lond.* 586, 3219–3230.
- Benkalfat, N.B., Merzouk, H., Bouanane, S., Merzouk, S.A., Bellenger, J., Gresti, J., Tessier, C., Narce, M., 2011. Altered adipose tissue metabolism in offspring of dietary obese rat dams. *Clin. Sci. (Lond.)* 121, 19–28.
- Berends, L.M., Ozanne, S.E., 2012. Early determinants of type-2 diabetes. *Best practice & research. Clin. Endocrinol. Metab.* 26, 569–580.
- Blackmore, H.L., Ozanne, S.E., 2013. Maternal diet-induced obesity and offspring cardiovascular health. *J. Dev. Orig. Health Dis.* 4, 338–347.
- Blackmore, H.L., Ozanne, S.E., 2014. Programming of cardiovascular disease across the life-course. *J. Mol. Cell. Cardiol.* 83, 122–130.
- Blackmore, H.L., Niu, Y., Fernandez-Twinn, D.S., Tarry-Adkins, J.L., Giussani, D.A., Ozanne, S.E., 2014. Maternal diet-induced obesity programs cardiovascular dysfunction in adult male mouse offspring independent of current body weight. *Endocrinology* 155, 3970–3980.
- Blum, K., Gardner, E., Oscar-Berman, M., Gold, M., 2012. "Liking" and "wanting" linked to Reward Deficiency Syndrome (RDS): hypothesizing differential responsivity in brain reward circuitry. *Curr. Pharm. Des.* 18, 113–118.
- Blum, K., Thanos, P.K., Gold, M.S., 2014. Dopamine and glucose, obesity, and reward deficiency syndrome. *Front. Psychol.* 5, 919.
- Bocarsly, M.E., Barson, J.R., Hauca, J.M., Hoebel, B.G., Leibowitz, S.F., Avena, N.M., 2012. Effects of perinatal exposure to palatable diets on body weight and sensitivity to drugs of abuse in rats. *Physiol. Behav.* 107, 568–575.
- Bolton, J.L., Bilbo, S.D., 2014. Developmental programming of brain and behavior by perinatal diet: focus on inflammatory mechanisms. *Dialogues Clin. Neurosci.* 16, 307–320.
- Boney, C.M., Verma, A., Tucker, R., Vohr, B.R., 2005. Metabolic syndrome in childhood: association with birth weight, maternal obesity, and gestational diabetes mellitus. *Pediatrics* 115, e290–e296.
- Bouret, S.G., 2012. Nutritional programming of hypothalamic development: critical periods and windows of opportunity. *Int. J. Obes. Suppl.* 2, S19–S24.
- Bouret, S.G., 2013. Organizational actions of metabolic hormones. *Front. Neuroendocrinol.* 34, 18–26.
- Bouret, S.G., Draper, S.J., Simerly, R.B., 2004a. Formation of projection pathways from the arcuate nucleus of the hypothalamus to hypothalamic regions implicated in the neural control of feeding behavior in mice. *J. Neurosci.* 24, 2797–2805.
- Bouret, S.G., Draper, S.J., Simerly, R.B., 2004b. Trophic action of leptin on hypothalamic neurons that regulate feeding. *Science* 304, 108–110.
- Bouret, S.G., Bates, S.H., Chen, S., Myers Jr., M.G., Simerly, R.B., 2012. Distinct roles for specific leptin receptor signals in the development of hypothalamic feeding circuits. *J. Neurosci.* 32, 1244–1252.
- Briffa, J.F., McAinch, A.J., Romano, T., Wlodek, M.E., Hryciw, D.H., 2014. Leptin in pregnancy and development: a contributor to adulthood disease? *Am. J. Physiol. Endocrinol. Metab.* 308 (5), E335–350.
- Briley, A.L., Barr, S., Badger, S., Bell, R., Croker, H., Godfrey, K.M., Holmes, B., Kinnunen, T.I., Nelson, S.M., Oteng-Ntim, E., Patel, N., Robson, S.C., Sandall, J., Sanders, T., Sattar, N., Seed, P.T., Wardle, J., Poston, L., 2014. A complex intervention to improve pregnancy outcome in obese women; the UPBEAT randomised controlled trial. *BMC Pregnancy Childbirth* 14, 74.
- Bringhenti, I., Ormellas, F., Martins, M.A., Mandarim-de-Lacerda, C.A., Aguila, M.B., 2014. Early hepatic insult in the offspring of obese maternal mice. *Nutr. Res.* 35 (2), 136–145.
- Brisbois, T.D., Farmer, A.P., McCargar, L.J., 2012. Early markers of adult obesity: a review. *Obes. Rev.* 13, 347–367.
- Bruce, K.D., Cagampang, F.R., Argenton, M., Zhang, J., Ethirajan, P.L., Burdge, G.C., Bateman, A.C., Clough, G.F., Poston, L., Hanson, M.A., McConnell, J.M., Byrne, C.D., 2009. Maternal high-fat feeding primes steatohepatitis in adult mice offspring, involving mitochondrial dysfunction and altered lipogenesis gene expression. *Hepatology* 50, 1796–1808.
- Burger, K.S., Stice, E., 2011. Variability in reward responsivity and obesity: evidence from brain imaging studies. *Curr. Drug Abuse Rev.* 4, 182–189.
- Carlin, J., George, R., Reyes, T.M., 2013. Methyl donor supplementation blocks the adverse effects of maternal high fat diet on offspring physiology. *PLoS One* 8.
- Carter, L.G., Lewis, K.N., Wilkerson, D.C., Tobia, C.M., Ngo Tenlep, S.Y., Shridas, P., Garcia-Cazarin, M.L., Wolff, G., Andrade, F.H., Charnigo, R.J., Esser, K.A., Egan, J.M., de Cabo, R., Pearson, K.J., 2012. Perinatal exercise improves glucose homeostasis in adult offspring. *Am. J. Physiol. Endocrinol. Metab.* 303, E1061–E1068.
- Carter, L.G., Qi, N.R., De Cabo, R., Pearson, K.J., 2013. Maternal exercise improves insulin sensitivity in mature rat offspring. *Med. Sci. Sports Exerc.* 45, 832–840.
- Carter, R., Mouralidarane, A., Soeda, J., Ray, S., Pombo, J., Saraswati, R., Novelli, M., Fusai, G., Rappa, F., Saracino, C., Paziienza, V., Poston, L., Taylor, P.D., Vinciguerra, M., Oben, J.A., 2014. Non-alcoholic fatty pancreas disease pathogenesis: a role for developmental programming and altered circadian rhythms. *PLoS One* 9, e89505.
- Catalano, P.M., Thomas, A., Huston-Presley, L., Amini, S.B., 2003. Increased fetal adiposity: a very sensitive marker of abnormal in utero development. *Am. J. Obstet. Gynecol.* 189, 1698–1704.
- Cedergren, M.I., 2004. Maternal morbid obesity and the risk of adverse pregnancy outcome. *Obstet. Gynecol.* 103, 219–224.
- Challier, J.C., Basu, S., Bintein, T., Minium, J., Hotmire, K., Catalano, P.M., Hauguel-de Mouzon, S., 2008. Obesity in pregnancy stimulates macrophage accumulation and inflammation in the placenta. *Placenta* 29, 274–281.
- Chang, G.Q., Gaysinskaya, V., Karatayev, O., Leibowitz, S.F., 2008. Maternal high-fat diet and fetal programming: increased proliferation of hypothalamic peptide-producing neurons that increase risk for overeating and obesity. *J. Neurosci.* 28, 12107–12119.
- Chen, H., Morris, M.J., 2009. Differential responses of orexigenic neuropeptides to fasting in offspring of obese mothers. *Obesity* 17, 1356–1362.
- Chen, H., Simar, D., Morris, M.J., 2009. Hypothalamic neuroendocrine circuitry is programmed by maternal obesity: interaction with postnatal nutritional environment. *PLoS One* 4, e6259.
- Chiswick, C.A., Reynolds, R.M., Denison, F.C., Whyte, S.A., Drake, A.J., Newby, D.E., Walker, B.R., Forbes, S., Murray, G.D., Quenby, S., Wray, S., Norman, J.E., 2015. Efficacy of metformin in pregnant obese women: a randomised controlled trial. *BMJ Open* 5, e006854.
- Collden, G., Ballard, E., Parkash, J., Caron, E., Langlet, F., Prevot, V., Bouret, S.G., 2015. Neonatal overnutrition causes early alterations in the central response to peripheral ghrelin. *Mol. Metab.* 4, 15–24.
- Cooper, R., Hypponen, E., Berry, D., Power, C., 2010. Associations between parental and offspring adiposity up to midlife: the contribution of adult lifestyle factors in the 1958 British Birth Cohort Study. *Am. J. Clin. Nutr.* 92, 946–953.
- Cordero, P., Milagro, F.I., Campion, J., Martinez, J.A., 2014. Supplementation with methyl donors during lactation to high-fat-sucrose-fed dams protects offspring against liver fat accumulation when consuming an obesogenic diet. *J. Dev. Orig. Health Dis.* 5, 385–395.
- Cottrell, E.C., Ozanne, S.E., 2007. Developmental programming of energy balance and the metabolic syndrome. *Proc. Nutr. Soc.* 66, 198–206.
- Coupe, B., Bouret, S.G., 2013. Development of the hypothalamic melanocortin system. *Front. Endocrinol.* 4, 38.
- Dabelea, D., 2007. The predisposition to obesity and diabetes in offspring of diabetic mothers. *Diabetes Care* 30, S169–S174.
- Dahlgren, J., Nilsson, C., Jennische, E., Ho, H.P., Eriksson, E., Niklasson, A., Bjornorp, P., Albertsson Wikland, K., Holmang, A., 2001. Prenatal cytokine exposure results in obesity and gender-specific programming. *Am. J. Physiol. Endocrinol. Metab.* 281, E326–E334.
- de Rooij, S.R., Painter, R.C., Phillips, D.I.W., Osmond, C., Michels, R.P.J., Godland, I.F., Bossuyt, P.M.M., Bleker, O.P., Roseboom, T.J., 2006. Impaired insulin secretion after prenatal exposure to the Dutch famine. *Diabetes Care* 29, 1897–1901.
- Delahaye, F., Breton, C., Risold, P.Y., Enache, M., Dutriez-Casteloot, I., Laborie, C., Lesage, J., Vieau, D., 2008. Maternal perinatal undernutrition drastically reduces postnatal leptin surge and affects the development of arcuate nucleus proopiomelanocortin neurons in neonatal male rat pups. *Endocrinology* 149, 470–475.
- Dhar, M., Wayman, G.A., Zhu, M., Lambert, T.J., Davare, M.A., Appleyard, S.M., 2014a. Leptin-induced spine formation requires TrpC channels and the CaM kinase cascade in the hippocampus. *J. Neurosci.* 34, 10022–10033.
- Dhar, M., Zhu, M., Impy, S., Lambert, T.J., Bland, T., Karatsoreos, I.N., Nakazawa, T., Appleyard, S.M., Wayman, G.A., 2014b. Leptin induces hippocampal synaptogenesis via CREB-regulated microRNA-132 suppression of p250GAP. *Mol. Endocrinol.* 28, 1073–1087.
- Dietrich, M.O., Bober, J., Ferreira, J.G., Tellez, L.A., Mineur, Y.S., Souza, D.O., Gao, X.B., Picciotto, M.R., Araujo, I., Liu, Z.W., Horvath, T.L., 2012. AgRP neurons regulate development of dopamine neuronal plasticity and nonfood-associated behaviors. *Nat. Neurosci.* 15, 1108–1110.
- Ditchfield, A.M., Desforges, M., Mills, T.A., Glazier, J.D., Wareing, M., Mynett, K., Sibley, C.P., Greenwood, S.L., 2014. Maternal obesity is associated with a reduction in placental taurine transporter activity. *Int. J. Obes. (Lond)* 39 (4), 557–564.
- Dodd, J.M., Cramp, C., Sui, Z., Yelland, L.N., Deussen, A.R., Grivell, R.M., Moran, L.J., Crowther, C.A., Turnbull, D., McPhee, A.J., Wittert, G., Owens, J.A., Robinson, J.S., Limit Randomised Trial Group, 2014. The effects of antenatal dietary and lifestyle advice for women who are overweight or obese on maternal diet and physical activity: the LIMIT randomised trial. *BMC Med.* 12, 161.
- Dominguez-Salas, P., Moore, S.E., Baker, M.S., Bergen, A.W., Cox, S.E., Dyer, R.A., Waterland, A.J., Guan, Y.T., Laritsky, E., Silver, M.J., Swan, G.E., Zeisel, S.H., Innis, S.M., Wulford, R.A., Prentice, A.M., Hennig, B.J., 2014. Maternal nutrition at conception modulates DNA methylation of human metastable epialleles. *Nat. Commun.* 5.

- Drake, A.J., Reynolds, R.M., 2010. Impact of maternal obesity on offspring obesity and cardiometabolic disease risk. *Reproduction* 140, 387–398.
- Duque-Guimaraes, D.E., Ozanne, S.E., 2013. Nutritional programming of insulin resistance: causes and consequences. *Trends Endocrin. Metab.* 24, 525–535.
- Egecioglu, E., Skibicka, K.P., Hansson, C., Alvarez-Crespo, M., Friberg, P.A., Jerlhag, E., Engel, J.A., Dickson, S.L., 2011. Hedonic and incentive signals for body weight control. *Rev. Endocr. Metab. Disord.* 12, 141–151.
- Fan, L., Lindsley, S.R., Comstock, S.M., Takahashi, D.L., Evans, A.E., He, G.W., Thornburg, K.L., Grove, K.L., 2013. Maternal high-fat diet impacts endothelial function in nonhuman primate offspring. *Int. J. Obes. (Lond)* 37, 254–262.
- Fantus, I.G., 2015. Is metformin ready for prime time in pregnancy? Probably not yet. *Diabetes Metab. Res. Rev.* 31, 36–38.
- Ferezou-Viala, J., Roy, A.F., Serougne, C., Grippo, D., Parquet, M., Bailleux, V., Gertler, A., Delplanque, B., Djiane, J., Riottot, M., Taouis, M., 2007. Long-term consequences of maternal high-fat feeding on hypothalamic leptin sensitivity and diet-induced obesity in the offspring. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 293, R1056–R1062.
- Fernandez-Twinn, D.S., Blackmore, H.L., Siggins, L., Giussani, D.A., Cross, C.M., Foo, R., Ozanne, S.E., 2012. The programming of cardiac hypertrophy in the offspring by maternal obesity is associated with hyperinsulinemia, AKT, ERK, and mTOR activation. *Endocrinology* 153, 5961–5971.
- Fernandez-Twinn, D.S., Alfaradhi, M.Z., Martin-Gronert, M.S., Duque-Guimaraes, D.E., Piekarz, A., Ferland-McCollough, D., Bushell, M., Ozanne, S.E., 2014. Downregulation of IRS-1 in adipose tissue of offspring of obese mice is programmed cell-autonomously through post-transcriptional mechanisms. *Mol. Metab.* 3, 325–333.
- Ferretti, G., Cester, A.M., Bacchetti, T., Raffaelli, F., Vignini, A., Orici, F., Martino, C., Tranquilli, A., 2014. Leptin and paraoxonase activity in cord blood from obese mothers. *J. Matern. Fetal. Neonatal Med.* 27, 1353–1356.
- Finger, B.C., Dinan, T.G., Cryan, J.F., 2012. Diet-induced obesity blunts the behavioural effects of ghrelin: studies in a mouse-progressive ratio task. *Psychopharmacology (Berl)* 220, 173–181.
- Fisher, S.C., Kim, S.Y., Sharma, A.J., Rochat, R., Morrow, B., 2013. Is obesity still increasing among pregnant women? Prepregnancy obesity trends in 20 states, 2003–2009. *Prev. Med.* 56, 372–378.
- Franke, K., Harder, T., Aerts, L., Melchior, K., Fahrenkrog, S., Rodekamp, E., Ziska, T., Van Assche, F.A., Dudenhausen, J.W., Plagemann, A., 2005. 'Programming' of orexigenic and anorexigenic hypothalamic neurons in offspring of treated and untreated diabetic mother rats. *Brain Res.* 1031, 276–283.
- Gaudet, L., Ferraro, Z.M., Wen, S.W., Walker, M., 2014. Maternal obesity and occurrence of fetal macrosomia: a systematic review and meta-analysis. *BioMed. Res. Int.* 2014, 640291.
- Gillman, M.W., Oakey, H., Baghurst, P.A., Volkmer, R.E., Robinson, J.S., Crowther, C.A., 2010. Effect of treatment of gestational diabetes mellitus on obesity in the next generation. *Diabetes Care* 33, 964–968.
- Gonzalez-Bulnes, A., Astiz, S., 2015. Effects of lifestyle on female reproductive features and success: lessons from animal models. *Handbook of Fertility: Nutrition, Diet, Lifestyle and Reproductive Health*. Academic Press, pp. 191–202.
- Gonzalez-Bulnes, A., Astiz, S., Ovilo, C., Lopez-Bote, C.J., Sanchez-Sanchez, R., Perez-Solana, M.L., Torres-Rovira, L., Ayuso, M., Gonzalez, J., 2014. Early-postnatal changes in adiposity and lipids profile by transgenerational developmental programming in swine with obesity/leptin resistance. *J. Endocrinol.* 223, M17–M29.
- Grissom, N., Bowman, N., Reyes, T.M., 2014a. Epigenetic programming of reward function in offspring: a role for maternal diet. *Mamm. Genome* 25, 41–48.
- Grissom, N.M., Lyde, R., Christ, L., Sasson, I.E., Carlin, J., Vitins, A.P., Simmons, R.A., Reyes, T.M., 2014b. Obesity at conception programs the opioid system in the offspring brain. *Neuropsychopharmacology* 39, 801–810.
- Grissom, N.M., Herdt, C.T., Desilets, J., Lidsky-Everson, J., Reyes, T.M., 2014c. Dissociable deficits of executive function caused by gestational adversity are linked to specific transcriptional changes in the prefrontal cortex. *Neuropsychopharmacology* 40 (6), 1353–1363.
- Gugusheff, J.R., Ong, Z.Y., Muhlhauser, B.S., 2014. The early origins of food preferences: targeting the critical windows of development. *FASEB J.* 29 (2), 365–373.
- Guimond, D., Diabira, D., Porcher, C., Bader, F., Ferrand, N., Zhu, M., Appleyard, S.M., Wayman, G.A., Gaiarsa, J.L., 2014. Leptin potentiates GABAergic synaptic transmission in the developing rodent hippocampus. *Front. Cell. Neurosci.* 8, 235.
- Gupta, A., Srinivasan, M., Thamadiok, S., Patel, M.S., 2009. Hypothalamic alterations in fetuses of high fat diet-fed obese female rats. *J. Endocrinol.* 200, 293–300.
- Hales, C.N., Barker, D.J.P., Clark, P.M.S., Cox, L.J., Fall, C., Osmond, C., Winter, P.D., 1991. Fetal and infant growth and impaired glucose-tolerance at age 64. *Brit. Med. J.* 303, 1019–1022.
- Hassiakos, D., Eleftheriades, M., Papastefanou, I., Lambrinou, I., Kappou, D., Lavranos, D., Akalestos, A., Aravantinos, L., Pervanidou, P., Chrousos, G., 2015. Increased maternal serum interleukin-6 concentrations at 11 to 14 weeks of gestation in low risk pregnancies complicated with gestational diabetes mellitus: development of a prediction model. *Horm. Metab. Res.* (pub. ahead of print).
- Hastie, R., Lappas, M., 2014. The effect of pre-existing maternal obesity and diabetes on placental mitochondrial content and electron transport chain activity. *Placenta* 35, 673–683.
- Hillier, T.A., Pedula, K.L., Schmidt, M.M., Mullen, J.A., Charles, M.A., Pettitt, D.J., 2007. Childhood obesity and metabolic imprinting – The ongoing effects of maternal hyperglycemia. *Diabetes Care* 30, 2287–2292.
- Horan, M.K., McGowan, C.A., Gibney, E.R., Donnelly, J.M., McAuliffe, F.M., 2014. Maternal diet and weight at 3 months postpartum following a pregnancy intervention with a low glycaemic index diet: results from the ROLO randomised control trial. *Nutrients* 6, 2946–2955.
- Howie, G.J., Sloboda, D.M., Kamal, T., Vickers, M.H., 2009. Maternal nutritional history predicts obesity in adult offspring independent of postnatal diet. *J. Physiol. Lond.* 587, 905–915.
- Hu, Z., Cooper, M., Crockett, D.P., Zhou, R., 2004. Differentiation of the midbrain dopaminergic pathways during mouse development. *J. Comp. Neurol.* 476, 301–311.
- Isganaitis, E., Woo, M., Ma, H., Chen, M., Kong, W., Lytras, A., Sales, V., Decoste-Lopez, J., Lee, K.J., Leatherwood, C., Lee, D., Fitzpatrick, C., Gall, W., Watkins, S., Patti, M.E., 2014. Developmental programming by maternal insulin resistance: hyperinsulinemia, glucose intolerance, and dysregulated lipid metabolism in male offspring of insulin-resistant mice. *Diabetes* 63, 688–700.
- Johnson, P.M., Kenny, P.J., 2010. Dopamine D2 receptors in addiction-like reward dysfunction and compulsive eating in obese rats. *Nat. Neurosci.* 13, 635–641.
- Jones, H.N., Woollett, L.A., Barbour, N., Prasad, P.D., Powell, T.L., Jansson, T., 2009. High-fat diet before and during pregnancy causes marked up-regulation of placental nutrient transport and fetal overgrowth in C57/BL6 mice. *FASEB J.* 23, 271–278.
- Kang, S.S., Kurti, A., Fair, D.A., Fryer, J.D., 2014. Dietary intervention rescues maternal obesity induced behavior deficits and neuroinflammation in offspring. *J. Neuroinflammation* 11, 156.
- Karakosta, P., Georgiou, V., Fthenou, E., Papadopoulou, E., Roumeliotaki, T., Margioris, A., Castanas, E., Kampa, M., Kogevas, M., Chatzi, L., 2013. Maternal weight status, cord blood leptin and fetal growth: a prospective mother-child cohort study (Rhea study). *Paediatr. Perinat. Epidemiol.* 27, 461–471.
- Kepeczynska, M.A., Wargent, E.T., Cawthorne, M.A., Arch, J.R., O'Dowd, J.F., Stocker, C.J., 2013. Circulating levels of the cytokines IL10, IFN γ and resistin in an obese mouse model of developmental programming. *J. Dev. Orig. Health Dis.* 4, 491–498.
- Kim, D.W., Young, S.L., Grattan, D.R., Jasoni, C.L., 2014. Obesity during pregnancy disrupts placental morphology, cell proliferation, and inflammation in a sex-specific manner across gestation in the mouse. *Biol. Reprod.* 90, 130.
- Kirk, S.L., Samuelsson, A.M., Argenton, M., Dhonye, H., Kalamatianos, T., Poston, L., Taylor, P.D., Coen, C.W., 2009. Maternal obesity induced by diet in rats permanently influences central processes regulating food intake in offspring. *PLoS One* 4, e5870.
- Kral, J.G., Biron, S., Simard, S., Hould, F.S., Lebel, S., Marceau, S., Marceau, P., 2006. Large maternal weight loss from obesity surgery prevents transmission of obesity to children who were followed for 2 to 18 years. *Pediatrics* 118, e1644–e1649.
- Lager, S., Samuelsson, A.M., Taylor, P.D., Poston, L., Powell, T.L., Jansson, T., 2014. Diet-induced obesity in mice reduces placental efficiency and inhibits placental mTOR signaling. *Physiol. Rep.* 2, e00242.
- Laker, R.C., Lillard, T.S., Okutsu, M., Zhang, M., Hoehn, K.L., Connelly, J.J., Yan, Z., 2014. Exercise prevents maternal high-fat diet-induced hypermethylation of the Pgc- α gene and age-dependent metabolic dysfunction in the offspring. *Diabetes* 63, 1605–1611.
- Leininger, G.M., Opland, D.M., Jo, Y.H., Faouzi, M., Christensen, L., Cappellucci, L.A., Rhodes, C.J., Gnegy, M.E., Becker, J.B., Pothos, E.N., Seasholtz, A.F., Thompson, R.C., Myers Jr., M.G., 2011. Leptin action via neurotensin neurons controls orexin, the mesolimbic dopamine system and energy balance. *Cell Metab.* 14, 313–323.
- Li, F., Wang, X., Capasso, J.M., Gerdes, A.M., 1996. Rapid transition of cardiac myocytes from hyperplasia to hypertrophy during postnatal development. *J. Mol. Cell. Cardiol.* 28, 1737–1746.
- Li, B., Shi, Z., Cassaglia, P.A., Brooks, V.L., 2013. Leptin acts in the forebrain to differentially influence baroreflex control of lumbar, renal, and splanchnic sympathetic nerve activity and heart rate. *Hypertension* 61, 812–819.
- Liu, K.Y., Chow, J.M., Sherry, C., 2013. Early life obesity and diabetes: origins in pregnancy. *Open J. Endocr. Metab. Dis.* 3, 1–12.
- Liu, G.S., Li, N., Sun, S.R., Wen, J., Lyu, F.J., Gao, W., Li, L.L., Chen, F., Baccarelli, A.A., Hou, L.F., Hu, G., 2014. Maternal OGTT glucose levels at 26–30 gestational weeks with offspring growth and development in early infancy. *BioMed. Res. Int.* 516980. <http://dx.doi.org/10.1155/2014/516980>.
- Long, N.M., Ford, S.P., Nathanielsz, P.W., 2011. Maternal obesity eliminates the neonatal lamb plasma leptin peak. *J. Physiol.* 589, 1455–1462.
- Lumey, L.H., Stein, A.D., 1997. Offspring birth weights after maternal intrauterine under-nutrition: a comparison within sibships. *Am. J. Epidemiol.* 146, 810–819.
- Mark, A.L., Agassandian, K., Morgan, D.A., Liu, X., Cassell, M.D., Rahmouni, K., 2009. Leptin signaling in the nucleus tractus solitarius increases sympathetic nerve activity to the kidney. *Hypertension* 53, 375–380.
- Marques, E.B., Rocha, N.N., Dos Santos, M.C., Nascimento, J.H., Scaramello, C.B., 2014a. Cardiac programming in rats submitted to leptin treatment during lactation. *Int. J. Cardiol.* 181C, 141–143.
- Marques, A.H., Bjorke-Monsen, A.L., Teixeira, A., Silverman, M.N., 2014b. Maternal stress, nutrition and physical activity: impact on immune function, CNS development and psychopathology. *Brain Res.* 1617, 28–46.
- Marsh, A.J., Fontes, M.A., Killinger, S., Pawlak, D.B., Polson, J.W., Dampney, R.A., 2003. Cardiovascular responses evoked by leptin acting on neurons in the ventromedial and dorsomedial hypothalamus. *Hypertension* 42, 488–493.
- Martin-Gronert, M.S., Ozanne, S.E., 2013. Early life programming of obesity. *Med. Wieku Rozwoj.* 17, 7–12.
- Martin-Gronert, M.S., Fernandez-Twinn, D.S., Poston, L., Ozanne, S.E., 2010. Altered hepatic insulin signalling in male offspring of obese mice. *J. Dev. Orig. Health Dis.* 1, 184–191.
- McAllister, E.J., Dhurandhar, N.V., Keith, S.W., Aronne, L.J., Barger, J., Baskin, M., Benca, R.M., Biggio, J., Boggiano, M.M., Eisenmann, J.C., Elobeid, M., Fontaine, K.R., Gluckman, P., Hanlon, E.C., Katzmarzyk, P., Pietrobelli, A., Redden, D.T., Ruden, D.M., Wang, C., Waterland, R.A., Wright, S.M., Allison, D.B., 2009. Ten putative contributors to the obesity epidemic. *Crit. Rev. Food Sci. Nutr.* 49, 868–913.
- Mehta, S.H., Kerver, J.M., Sokol, R.J., Keating, D.P., Paneth, N., 2014. The association between maternal obesity and neurodevelopmental outcomes of offspring. *J. Pediatr.* 165, 891–896.

- Mele, J., Muralimanoharan, S., Maloyan, A., Myatt, L., 2014. Impaired mitochondrial function in human placenta with increased maternal adiposity. *Am. J. Physiol. Endocrinol. Metab.* 307, E419–E425.
- Morris, M.J., Chen, H., 2009. Established maternal obesity in the rat reprograms hypothalamic appetite regulators and leptin signaling at birth. *Int. J. Obes.* 33, 115–122.
- Morrison, J.L., Botting, K.J., Dyer, J.L., Williams, S.J., Thornburg, K.L., McMillen, I.C., 2007. Restriction of placental function alters heart development in the sheep fetus. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 293, R306–R313.
- Murabayashi, N., Sugiyama, T., Zhang, L., Kamimoto, Y., Umekawa, T., Ma, N., Sagawa, N., 2013. Maternal high-fat diets cause insulin resistance through inflammatory changes in fetal adipose tissue. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 169, 39–44.
- Nadif, R., Dilworth, M.R., Sibley, C.P., Baker, P.N., Davidge, S.T., Gibson, J.M., Aplin, J.D., Westwood, M., 2015. The maternal environment programs postnatal weight gain and glucose tolerance of male offspring, but placental and fetal growth are determined by fetal genotype in the *leprdb/+* model of gestational diabetes. *Endocrinology* 1, 360–366.
- Naef, L., Moquin, L., Dal Bo, G., Giros, B., Gratton, A., Walker, C.D., 2011. Maternal high-fat intake alters presynaptic regulation of dopamine in the nucleus accumbens and increases motivation for fat rewards in the offspring. *Neuroscience* 176, 225–236.
- Naef, L., Moquin, L., Gratton, A., Walker, C.D., 2013. Reduced anticipatory dopamine responses to food in rats exposed to high fat during early development. *Int. J. Obes. (Lond)* 37, 885–888.
- Nathanielsz, P.W., Ford, S.P., Long, N.M., Vega, C.C., Reyes-Castro, L.A., Zambrano, E., 2013. Interventions to prevent adverse fetal programming due to maternal obesity during pregnancy. *Nutr. Rev.* 71 (Suppl. 1), S78–S87.
- Ng, M., Fleming, T., Robinson, M., Thomson, B., Graetz, N., Margono, C., Mullany, E.C., Biryukov, S., Abbafati, C., Abera, S.F., Abraham, J.P., Abu-Rmeileh, N.M., Achoki, T., AlBuhairan, F.S., Alemu, Z.A., Alfonso, R., Ali, M.K., Ali, R., Guzman, N.A., Ammar, W., Anvari, P., Banerjee, A., Barquera, S., Basu, S., Bennett, D.A., Bhutta, Z., Blore, J., Cabral, N., Nonato, I.C., Chang, J.C., Chowdhury, R., Courville, K.J., Criqui, M.H., Cundiff, D.K., Dabhadkar, K.C., Dandona, L., Davis, A., Dayama, A., Dharmaratne, S.D., Ding, E.L., Durrani, A.M., Esteghamati, A., Farzadfar, F., Fay, D.F., Feigin, V.L., Flaxman, A., Forouzanfar, M.H., Goto, A., Green, M.A., Gupta, R., Hafezi-Nejad, N., Hankey, G.J., Harewood, H.C., Havmoeller, R., Hay, S., Hernandez, L., Husseini, A., Idrisov, B.T., Ikeda, N., Islami, F., Jahangir, E., Jassal, S.K., Jee, S.H., Jeffreys, M., Jonas, J.B., Kabagambe, E.K., Khalifa, S.E., Kengne, A.P., Khader, Y.S., Khang, Y.H., Kim, D., Kimokoti, R.W., Kinge, J.M., Kokubo, Y., Kosen, S., Kwan, G., Lai, T., Leinsalu, M., Li, Y., Liang, X., Liu, S., Logroscino, G., Lotufo, P.A., Lu, Y., Ma, J., Mainoo, N.K., Mensah, G.A., Merriman, T.R., Mokdad, A.H., Moschandreas, J., Naghavi, M., Naheed, A., Nand, D., Narayan, K.M., Nelson, E.L., Neuhouser, M.L., Nisar, M.I., Ohkubo, T., Oti, S.O., Pedroza, A., Prabhakaran, D., Roy, N., Sampson, U., Seo, H., Sepanlou, S.G., Shibuya, K., Shiri, R., Shiue, I., Singh, G.M., Singh, J.A., Skirbekk, V., Stapelberg, N.J., Sturua, L., Sykes, B.L., Tobias, M., Tran, B.X., Trasande, L., Toyoshima, H., van de Vijver, S., Vasankari, T.J., Veerman, J.L., Velasquez-Melendez, G., Vlassov, V.V., Vollset, S.E., Vos, T., Wang, C., Wang, X., Weiderpass, E., Werdecker, A., Wright, J.L., Yang, Y.C., Yatsuya, H., Yoon, J., Yoon, S.J., Zhao, Y., Zhou, M., Zhu, S., Lopez, A.D., Murray, C.J., Gakidou, E., 2014. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 384, 766–781.
- Nicholas, L.M., Morrison, J.L., Rattanatr, L., Ozanne, S.E., Kleemann, D.O., Walker, S.K., MacLaughlin, S.M., Zhang, S., Martin-Gronert, M.S., McMillen, I.C., 2013a. Differential effects of exposure to maternal obesity or maternal weight loss during the periconceptual period in the sheep on insulin signalling molecules in skeletal muscle of the offspring at 4 months of age. *PLoS One* 8.
- Nicholas, L.M., Rattanatr, L., MacLaughlin, S.M., Ozanne, S.E., Kleemann, D.O., Walker, S.K., Morrison, J.L., Zhang, S., Muhlhauser, B.S., Martin-Gronert, M.S., McMillen, I.C., 2013b. Differential effects of maternal obesity and weight loss in the periconceptual period on the epigenetic regulation of hepatic insulin-signaling pathways in the offspring. *FASEB J.* 27, 3786–3796.
- Nivoit, P., Morens, C., Van Assche, F.A., Jansen, E., Poston, L., Remacle, C., Reusens, B., 2009. Established diet-induced obesity in female rats leads to offspring hyperphagia, adiposity and insulin resistance. *Diabetologia* 52, 1133–1142.
- Oben, J.A., Mouralidarane, A., Samuelsson, A.M., Matthews, P.J., Morgan, M.L., McKee, C., Soeda, J., Fernandez-Twinn, D.S., Martin-Gronert, M.S., Ozanne, S.E., Sigala, B., Novelli, M., Poston, L., Taylor, P.D., 2010a. Maternal obesity during pregnancy and lactation programs the development of offspring non-alcoholic fatty liver disease in mice. *J. Hepatol.* 52, 913–920.
- Oben, J.A., Patel, T., Mouralidarane, A., Samuelsson, A.M., Matthews, P., Pombo, J., Morgan, M., McKee, C., Soeda, J., Novelli, M., Poston, L., Taylor, P., 2010b. Maternal obesity programmes offspring development of non-alcoholic fatty pancreas disease. *Biochem. Biophys. Res. Commun.* 394, 24–28.
- Ong, Z.Y., Muhlhauser, B.S., 2011. Maternal “junk-food” feeding of rat dams alters food choices and development of the mesolimbic reward pathway in the offspring. *FASEB J.* 25, 2167–2179.
- Ong, Z.Y., Muhlhauser, B.S., 2014. Consuming a low-fat diet from weaning to adulthood reverses the programming of food preferences in male, but not in female, offspring of ‘junk food’-fed rat dams. *Acta Physiol.* 210, 127–141.
- Page, K.C., Malik, R.E., Ripple, J.A., Anday, E.K., 2009. Maternal and postweaning diet interaction alters hypothalamic gene expression and modulates response to a high-fat diet in male offspring. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 297, R1049–R1057.
- Parente, L.B., Aguila, M.B., Mandarim-de-Lacerda, C.A., 2008. Deleterious effects of high-fat diet on perinatal and postweaning periods in adult rat offspring. *Clin. Nutr.* 27, 623–634.
- Pelizzo, G., Calcaterra, V., Fusillo, M., Nakib, G., Ierullo, A.M., Alfei, A., Spinillo, A., Stronati, M., Cena, H., 2014. Malnutrition in pregnancy following bariatric surgery: three clinical cases of fetal neural defects. *Nutr. J.* 13.
- Plagemann, A., Harder, T., Rake, A., Voits, M., Fink, H., Rohde, W., Dorner, G., 1999. Perinatal elevation of hypothalamic insulin, acquired malformation of hypothalamic galaninergic neurons, and syndrome x-like alterations in adulthood of neonatally overfed rats. *Brain Res.* 836, 146–155.
- Plagemann, A., Harder, T., Franke, K., Kohlhoff, R., 2002. Long-term impact of neonatal breast-feeding on body weight and glucose tolerance in children of diabetic mothers. *Diabetes Care* 25, 16–22.
- Plagemann, A., Harder, T., Brunn, M., Harder, A., Roepke, K., Wittrock-Staar, M., Ziska, T., Schellong, K., Rodekamp, E., Melchior, K., Dudenhausen, J.W., 2009. Hypothalamic proopiomelanocortin promoter methylation becomes altered by early overfeeding: an epigenetic model of obesity and the metabolic syndrome. *J. Physiol. Lond.* 587, 4963–4976.
- Plagemann, A., Roepke, K., Harder, T., Brunn, M., Harder, A., Wittrock-Staar, M., Ziska, T., Schellong, K., Rodekamp, E., Melchior, K., Dudenhausen, J.W., 2010. Epigenetic malprogramming of the insulin receptor promoter due to developmental overfeeding. *J. Perinat. Med.* 38, 393–400.
- Plagemann, A., Harder, T., Schellong, K., Schulz, S., Stupin, J.H., 2012. Early postnatal life as a critical time window for determination of long-term metabolic health. *Best Pract. Res. Clin. Endocrinol. Metab.* 26, 641–653.
- Prior, L.J., Davern, P.J., Burke, S.L., Lim, K., Armitage, J.A., Head, G.A., 2014. Exposure to a high-fat diet during development alters leptin and ghrelin sensitivity and elevates renal sympathetic nerve activity and arterial pressure in rabbits. *Hypertension* 63, 338–345.
- Rahmouni, K., Morgan, D.A., Morgan, G.M., Mark, A.L., Haynes, W.G., 2005. Role of selective leptin resistance in diet-induced obesity hypertension. *Diabetes* 54, 2012–2018.
- Rajia, S., Chen, H., Morris, M.J., 2010. Maternal overnutrition impacts offspring adiposity and brain appetite markers—modulation by postweaning diet. *J. Neuroendocrinol.* 22, 905–914.
- Rattanatr, L., MacLaughlin, S.M., Kleemann, D.O., Walker, S.K., Muhlhauser, B.S., McMillen, I.C., 2010. Impact of maternal periconceptual overnutrition on fat mass and expression of adipogenic and lipogenic genes in visceral and subcutaneous fat depots in the postnatal lamb. *Endocrinology* 151, 5195–5205.
- Ravelli, G.P., Stein, Z.A., Susser, M.W., 1976. Obesity in young men after famine exposure in utero and early infancy. *New Engl. J. Med.* 295, 349–353.
- Ravelli, A.C.J., van der Meulen, J.H.P., Osmond, C., Barker, D.J.P., Bleker, O.P., 1999. Obesity at the age of 50 y in men and women exposed to famine prenatally. *Am. J. Clin. Nutr.* 70, 811–816.
- Refuerzo, J.S., Viteri, O.A., Hutchinson, M., Pedroza, C., Blackwell, S.C., Tyson, J.E., Ramin, S.M., 2015. The effects of metformin on weight loss in women with gestational diabetes: a pilot randomized, placebo-controlled trial. *Am. J. Obstet. Gynecol.* 212 (389), e381–e389.
- Rodriguez, A., 2010. Maternal pre-pregnancy obesity and risk for inattention and negative emotionality in children. *J. Child Psychol. Psychiatry* 51, 134–143.
- Rodriguez, J.S., Rodriguez-Gonzalez, G.L., Reyes-Castro, L.A., Ibanez, C., Ramirez, A., Chavira, R., Larrea, F., Nathanielsz, P.W., Zambrano, E., 2012. Maternal obesity in the rat programs male offspring exploratory, learning and motivation behavior: prevention by dietary intervention pre-gestation or in gestation. *Int. J. Dev. Neurosci.* 30, 75–81.
- Rolls, B.A., Gurr, M.I., Vanduijvenvoorde, P.M., Rolls, B.J., Rowe, E.A., 1986. Lactation in lean and obese rats — effect of cafeteria feeding and of dietary obesity on milk-composition. *Physiol. Behav.* 38, 185–190.
- Rosen, E.D., Spiegelman, B.M., 2014. What we talk about when we talk about fat. *Cell* 156, 20–44.
- Saben, J.L., Bales, E.S., Jackman, M.R., Orlicky, D., MacLean, P.S., McManaman, J.L., 2014. Maternal obesity reduces milk lipid production in lactating mice by inhibiting acetyl-CoA carboxylase and impairing fatty acid synthesis. *PLoS One* 9.
- Sagedal, L.R., Overby, N.C., Lohne-Seiler, H., Bere, E., Torstveit, M.K., Henriksen, T., Vistad, L., 2013. Study protocol: fit for delivery — can a lifestyle intervention in pregnancy result in measurable health benefits for mothers and newborns? A randomized controlled trial. *BMC Public Health* 13, 132.
- Salomaki, H., Heinaniemi, M., Vahatalo, L.H., Ailanen, L., Eerola, K., Ruohonen, S.T., Pesonen, U., Koulu, M., 2014. Prenatal metformin exposure in a maternal high fat diet mouse model alters the transcriptome and modifies the metabolic responses of the offspring. *PLoS One* 9, e115778.
- Samuelsson, A.M., Matthews, P.A., Argenton, M., Christie, M.R., McConnell, J.M., Jansen, E.H.J.M., Piersma, A.H., Ozanne, S.E., Twinn, D.F., Remacle, C., Rowlerson, A., Poston, L., Taylor, P.D., 2008. Diet-induced obesity in female mice leads to offspring hyperphagia, adiposity, hypertension, and insulin resistance — A novel murine model of developmental programming. *Hypertension* 51, 383–392.
- Samuelsson, A.M., Morris, A., Igosheva, N., Kirk, S.L., Pombo, J.M., Coen, C.W., Poston, L., Taylor, P.D., 2010. Evidence for sympathetic origins of hypertension in juvenile offspring of obese rats. *Hypertension* 55, 76–82.
- Sanders, T.R., Kim, D.W., Glendinning, K.A., Jasoni, C.L., 2014. Maternal obesity and IL-6 lead to aberrant developmental gene expression and deregulated neurite growth in the fetal arcuate nucleus. *Endocrinology* 155, 2566–2577.
- Seiple, B.D., Blomgren, K., Gimlin, K., Ferriero, D.M., Noble-Haesslein, L.J., 2013. Brain development in rodents and humans: identifying benchmarks of maturation and vulnerability to injury across species. *Prog. Neurobiol.* 106–107, 1–16.
- Seneviratne, S.N., Parry, G.K., McCowan, L.M.E., Ekeroma, A., Jiang, Y.N., Gusso, S., Peres, G., Rodrigues, R.O., Craigie, S., Cutfield, W.S., Hofman, P.L., 2014. Antenatal exercise in overweight and obese women and its effects on offspring and maternal health: design and rationale of the IMPROVE (Improving Maternal and Progeny Obesity Via Exercise) randomised controlled trial. *BMC Pregnancy Childbirth* 14.
- Sferruzzi-Perri, A.N., Vaughan, O.R., Haro, M., Cooper, W.N., Musial, B., Charalambous, M., Pestana, D., Ayyar, S., Ferguson-Smith, A.C., Burton, G.J., Constancia, M., Fowden, A.L., 2013. An obesogenic diet during mouse pregnancy modifies maternal nutrient partitioning and the fetal growth trajectory. *FASEB J.* 27, 3928–3937.

- Shai, D., Shoham-Vardi, I., Amsalem, D., Silverberg, D., Levi, I., Sheiner, E., 2014. Pregnancy outcome of patients following bariatric surgery as compared with obese women: a population-based study. *J. Matern. Fetal Neonatal Med.* 27, 275–278.
- Shankar, K., Harrell, A., Liu, X.L., Gilchrist, J.M., Ronis, M.J.J., Badger, T.M., 2008. Maternal obesity at conception programs obesity in the offspring. *Am. J. Physiol. Reg. Integr.* 294, R528–R538.
- Shankar, K., Kang, P., Harrell, A., Zhong, Y., Marecki, J.C., Ronis, M.J.J., Badger, T.M., 2010. Maternal overweight programs insulin and adiponectin signaling in the offspring. *Endocrinology* 151, 2577–2589.
- Shelley, P., Martin-Gronert, M.S., Rowleron, A., Poston, L., Heales, S.J.R., Hargreaves, I.P., McConnell, J.M., Ozanne, S.E., Fernandez-Twinn, D.S., 2009. Altered skeletal muscle insulin signaling and mitochondrial complex II–III linked activity in adult offspring of obese mice. *Am. J. Physiol. Reg. Integr.* 297, R675–R681.
- Shin, A.C., Berthoud, H.R., 2011. Food reward functions as affected by obesity and bariatric surgery. *Int. J. Obes. (Lond)* 35 (Suppl. 3), S40–S44.
- Simonds, S.E., Pryor, J.T., Ravussin, E., Greenway, F.L., Dileone, R., Allen, A.M., Bassi, J., Elmquist, J.K., Keogh, J.M., Henning, E., Myers Jr., M.G., Licinio, J., Brown, R.D., Enriori, P.J., O'Rahilly, S., Sternson, S.M., Grove, K.L., Spanswick, D.C., Farooqi, I.S., Cowley, M.A., 2014. Leptin mediates the increase in blood pressure associated with obesity. *Cell* 159, 1404–1416.
- Smith, S.E.P., Li, J., Garbett, K., Mirnics, K., Patterson, P.H., 2007. Maternal immune activation alters fetal brain development through interleukin-6. *J. Neurosci.* 27, 10695–10702.
- Smith, J., Cianflone, K., Biron, S., Hould, F.S., Lebel, S., Marceau, S., Lescelleur, O., Biertho, L., Simard, S., Kral, J.G., Marceau, P., 2009. Effects of maternal surgical weight loss in mothers on intergenerational transmission of obesity. *J. Clin. Endocrinol. Metab.* 94, 4275–4283.
- Song, Y., Yu, Y., Wang, D., Chai, S., Liu, D., Xiao, X., Huang, Y., 2015. Maternal high-fat diet feeding during pregnancy and lactation augments lung inflammation and remodeling in the offspring. *Respir. Physiol. Neurobiol.* 207, 1–6.
- Steculorum, S.M., Bouret, S.G., 2011a. Developmental effects of ghrelin. *Peptides* 32, 2362–2366.
- Steculorum, S.M., Bouret, S.G., 2011b. Maternal diabetes compromises the organization of hypothalamic feeding circuits and impairs leptin sensitivity in offspring. *Endocrinology* 152, 4171–4179.
- Steculorum, S.M., Vogt, M.C., Bruning, J.C., 2013. Perinatal programming of metabolic diseases: role of insulin in the development of hypothalamic neurocircuits. *Endocrinol. Metab. Clin. N. Am.* 42, 149–164.
- Steculorum, S.M., Collden, G., Coupe, B., Croizier, S., Lockie, S., Andrews, Z.B., Jarosch, F., Klusmann, S., Bouret, S.G., 2015. Neonatal ghrelin programs development of hypothalamic feeding circuits. *J. Clin. Invest.* 125 (2), 846–858.
- Stoeckel, L.E., Weller, R.E., Cook III, E.W., Twieg, D.B., Knowlton, R.C., Cox, J.E., 2008. Widespread reward-system activation in obese women in response to pictures of high-calorie foods. *Neuroimage* 41, 636–647.
- Tamashiro, K.L., Terrillion, C.E., Hyun, J., Koenig, J.L., Moran, T.H., 2009. Prenatal stress or high-fat diet increases susceptibility to diet-induced obesity in rat offspring. *Diabetes* 58, 1116–1125.
- Tarrade, A., Panchenko, P., Junien, C., Gabory, A., 2015. Placental contribution to nutritional programming of health and diseases: epigenetics and sexual dimorphism. *J. Exp. Biol.* 218, 50–58.
- Tong, J.F., Yan, X., Zhao, J.X., Zhu, M.J., Nathanielsz, P.W., Du, M., 2011. Metformin mitigates the impaired development of skeletal muscle in the offspring of obese mice. *Nutr. Diabetes* 1, e7.
- Torrens, C., Ethirajan, P., Bruce, K.D., Cagampang, F.R.A., Siow, R.C.M., Hanson, M.A., Byrne, C.D., Mann, G.E., Clough, G.F., 2012. Interaction between maternal and offspring diet to impair vascular function and oxidative balance in high fat fed male mice. *PLoS One* 7.
- Toste, F.P., de Moura, E.G., Lisboa, P.C., Fagundes, A.T., de Oliveira, E., Passos, M.C., 2006. Neonatal leptin treatment programmes leptin hypothalamic resistance and intermediary metabolic parameters in adult rats. *Br. J. Nutr.* 95, 830–837.
- Treesukosol, Y., Sun, B., Moghadam, A.A., Liang, N.C., Tamashiro, K.L., Moran, T.H., 2014. Maternal high-fat diet during pregnancy and lactation reduces the appetitive behavioral component in female offspring tested in a brief-access taste procedure. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 306, R499–R509.
- Valleau, J.C., Sullivan, E.L., 2014. The impact of leptin on perinatal development and psychopathology. *J. Chem. Neuroanat.* 61–62, 221–232.
- Vega, C.C., Reyes-Castro, L.A., Bautista, C.J., Larrea, F., Nathanielsz, P.W., Zambrano, E., 2013. Exercise in obese female rats has beneficial effects on maternal and male and female offspring metabolism. *Int. J. Obes. (Lond)* 39 (4), 712–719.
- Vogt, M.C., Paeger, L., Hess, S., Steculorum, S.M., Awazawa, M., Hampel, B., Neupert, S., Nicholls, H.T., Mauer, J., Hausen, A.C., Predel, R., Kloppenburg, P., Horvath, T.L., Bruning, J.C., 2014. Neonatal insulin action impairs hypothalamic neurocircuit formation in response to maternal high-fat feeding. *Cell* 156, 495–509.
- Volkow, N.D., Wang, G.J., Fowler, J.S., Telang, F., 2008. Overlapping neuronal circuits in addiction and obesity: evidence of systems pathology. *Philos. Trans. R. Soc. Lond. Ser. B Biol. Sci.* 363, 3191–3200.
- Volkow, N.D., Wang, G.J., Baler, R.D., 2011. Reward, dopamine and the control of food intake: implications for obesity. *Trends Cogn. Sci.* 15, 37–46.
- Vucetic, Z., Kimmel, J., Totoki, K., Hollenbeck, E., Reyes, T.M., 2010. Maternal high-fat diet alters methylation and gene expression of dopamine and opioid-related genes. *Endocrinology* 151, 4756–4764.
- Vucetic, Z., Kimmel, J., Reyes, T.M., 2011. Chronic high-fat diet drives postnatal epigenetic regulation of mu-opioid receptor in the brain. *Neuropsychopharmacology* 36, 1199–1206.
- Vucetic, Z., Carlin, J.L., Totoki, K., Reyes, T.M., 2012. Epigenetic dysregulation of the dopamine system in diet-induced obesity. *J. Neurochem.* 120, 891–898.
- Walker, C.D., Long, H., Williams, S., Richard, D., 2007. Long-lasting effects of elevated neonatal leptin on rat hippocampal function, synaptic proteins and NMDA receptor subunits. *J. Neurosci. Res.* 85, 816–828.
- Walker, C.D., Naef, L., d'Asti, E., Long, H., Xu, Z., Moreau, A., Azeddine, B., 2008. Perinatal maternal fat intake affects metabolism and hippocampal function in the offspring: a potential role for leptin. *Ann. N. Y. Acad. Sci.* 1144, 189–202.
- Walsh, J.M., Byrne, J., Mahony, R.M., Foley, M.E., McAuliffe, F.M., 2014. Leptin, fetal growth and insulin resistance in non-diabetic pregnancies. *Early Hum. Dev.* 90, 271–274.
- Whitaker, R.C., 2004. Predicting preschooler obesity at birth: the role of maternal obesity in early pregnancy. *Pediatrics* 114, e29–e36.
- Yan, X., Huang, Y., Zhao, J.X., Long, N.M., Uthlaut, A.B., Zhu, M.J., Ford, S.P., Nathanielsz, P.W., Du, M., 2011. Maternal obesity-impaired insulin signaling in sheep and induced lipid accumulation and fibrosis in skeletal muscle of offspring. *Biol. Reprod.* 85, 172–178.
- Yura, S., Itoh, H., Sagawa, N., Yamamoto, H., Masuzaki, H., Nakao, K., Kawamura, M., Takemura, M., Kakui, K., Ogawa, Y., Fujii, S., 2005. Role of premature leptin surge in obesity resulting from intrauterine undernutrition. *Cell Metab.* 1, 371–378.
- Zambrano, E., Martinez-Samayo, P.M., Rodriguez-Gonzalez, G.L., Nathanielsz, P.W., 2010. Dietary intervention prior to pregnancy reverses metabolic programming in male offspring of obese rats. *J. Physiol.* 588, 1791–1799.