


Fluid, electrolyte and acide-base disturbances in surgery (not only)



L.Dadak

Dept. of Anesthesia and
Intensive Care

St. Ann's University
Hospital

Top Secret

pH - situation of cells ... in Blood

- hyperkalemia
- hypokalemia

- acidosis acidemia
- alkalosis alkalemia

- acute/chronic ... resp/met ... Ac/Alk

- MD rule = Metabolic Disease

Homeostasis

- tendency to keep stable
- isovolemia
- H^+ = pH, pCO₂,
- Glc, ions
- isohydria, isoionia, isoosmia

Laboratory

- Never completely trust the laboratory
 - errors with blood sample
- Will result change my decision?

Never completely trust the laboratory

Quick	<0.10	(0.70 - 1.34)	<-()	repeated
aPTT	>150	s (20.0 - 40.0)	<=()	repeated
Fibrinogen	4.50	g/l (1.80 - 4.00)	()	->
Antitrombin III	32	% (80 - 120)	<-()	

----- same patient, 30 min later -----

Quick	0.55	(0.70 - 1.34)	<-()	
aPTT	44.7	s (20.0 - 40.0)	()	->
Aptt ratio	1.49			
Fibrinogen	5.40	g/l (1.80 - 4.00)	()	->
INR	1.59	(0.85 - 1.38)	()	->
Antitrombin III	61	% (80 - 120)	<-()	

Blood for analysis

- arterial
- capillary
- venous
 - peripheral
 - central
 - mixed venous (v.cava, a.pulmonalis)

Body Fluid Compartments

Total Body Water (TBW): 50-70% of total body wt.

- Avg. is greater for males.
- Highest in newborn, 75-80%. By first year of life TBW ~ 65%. Decreases with age.
- Most in muscle, less in fat.
- $TBW = ECF + ICF$
- $ICF \sim 2/3$ & $ECF \sim 1/3$
- $ECF = \text{Intravascular } (1/3) + \text{Interstitial } (2/3)$

- Compartment = place of water + ions

Water - compartments:

$$\text{ECF} = \text{IVF} + \text{ISF}$$

ICF

↔	↔	↔	↔
5%		15%	40%
Na		Na + -	Na
K		K + -	K
P		+ -	
		P	

Electrolyte Physiology

- Primary intravascular/ECF cation is Na^+ . Very small contribution of K^+ , Ca^{2+} , and Mg^{2+} .
- Primary intravascular/ECF anion is Cl^- . Smaller contribution from HCO_3^- , SO_4^{2-} & PO_4^{3-} , organic acids, and protein.
- Primary ICF cation is K^+ . Smaller contribution from Mg^{2+} & Na^+ .
- Number of intravasc anions not routinely detected.

ICF (mEq/L)

Cations

K⁺ (150-154)

Na⁺ (6-10)

Mg⁺² (40)

Anions

Organic PO₄⁻³ (100-106)

protein (40-60)

SO₄⁻² (17)

HCO₃⁻ (10-13)

organic acids (4)

ECF (mEq/L)

Na⁺(142)

Ca⁺² (5)

K⁺ (4-5)

Mg⁺² (3)

Cl⁻(103-105)

HCO₃⁻ (24-27)

protein (15)

PO₄⁻³ (3-5), SO₄⁻² (4)

Organic acids (2-5)

Intra Vascular Fluid

- Treatable volume

Provides:

- Nutrition
- Oxygenation
- Waste removal
- Temperature
- Alkalinity

Priorities

1. fluid volume and perfusion deficits
2. correction of pH
3. K, Ca, Mg
4. Na, Cl

IV Fluid/Electrolyte Therapy

Three key concepts in consideration of fluid and electrolyte management:



- cell membrane permeability
- osmolarity
- electroneutrality

Cell membrane permeability refers to the ability of a cell membrane to allow certain substances such as water and urea to pass freely, while charged ions such as sodium cannot cross the membrane and are trapped on one side of it.

Osmolarity

- Osmolarity is a property of particles in solution. If a substance can dissociate in solution, it may contribute more than one equivalent to the osmolarity of the solution. For instance, NaCl will dissociate into two osmotically active ions: Na and Cl. One millimolar NaCl yields a 2 milliosmolar solution.

Osmosis



Osmolarity, osmolality

- Each particle present in the water binds number molecules of water.

Serum osmolarity is **measured** directly by determining the **freezing** point of serum.

normal 275 .. 295 mOsm/l

Calculated osmolarity = $2 * \text{Na} + \text{Glc} + \text{Urea}$ [mOsm/l]
 $2 * 140 + 5 + 3$

Gap > 10 mOsm/l ... another solute (lactate, ethanol)

Gap > 50 mOsm/l ... often fatal

Osmolality [mmol/kg of water]

Electroneutrality

- in every solution
- sum of [mval/l] cations is equal to sum of anions
- Na^+ , K^+ , Mg^{++} , Ca^{++} ...
- Cl^- , HCO_3^- , PO_4^{--} , proteins-

Water

- 55% - 60%, new born 80% of body weight

Table 2-1
Water Exchange (60- to 80-kg Man)

<i>Routes</i>	<i>Average Daily Volume (mL)</i>	<i>Minimal, (mL)</i>	<i>Maximal, (mL)</i>
H₂O gain:			
Sensible:			
Oral fluids	800-1500	0	1500/h
Solid foods	500-700	0	1500
Insensible:			
Water of oxidation	250	125	800
Water of solution	0	0	500
H₂O loss:			
Sensible:			
Urine	800-1500	300	1400/h
Intestinal	0-250	0	2500/h
Sweat	0	0	4000/h
Insensible:			
Lungs and skin	600	600	1500

Table 2 Daily Electrolyte Requirements

	DAILY REQUIREMENT	FOR 70-KG ADULT	FOR 10-KG CHILD
Sodium	1-2 meq/kg	70-140 meq/day	10-20 meq/day
Potassium	0.5-1.0 meq/kg	35-70 meq/day	5-10 meq/day
Calcium	0.2-0.3 meq/kg	1.4-2.1 meq/day	2.0-3.0 meq/day
Magnesium	0.35-0.45 meq/kg	24.5-31.5 meq/day	3.5-4.5 meq/day
Chloride	equal to sodium	equal to sodium	equal to sodium
Bicarbonate/Acetate	use with chloride to balance cations and help pH	use with chloride to balance cations and help pH	use with chloride to balance cations and help pH

Basic Needs (Adult)

- Basic need 2 ml/kg/h
- Current losses
 - $1^\circ\text{C fever} = 500\text{ml/d}$
 - sweating
 - diarrhea ... water with ions [mmol/l]

	Sodium	Potassium	Chloride	Bicarbonate
Saliva	10-60	10-20	15-40	30-15
Stomach	40-100	5-15	15-20	—
Bile	130-140	4-6	95-105	30-40
Pancreas	130-140	4-6	40-60	80-100
Small intestine	130-140	4-6	40-60	80-100
Colon	80-140	25-45	80-100	30-50
Sweat	40-50	5-10	45-60	—

Types of IV Fluid

- Crystalloid
- Colloid

Crystalloid:

- Balanced salt/electrolyte solution; forms a true solution and is capable of passing through semipermeable membranes. May be isotonic, hypertonic, or hypotonic.
- Normal Saline (0.9% NaCl), Lactated Ringer's, Hypertonic saline (3, 5, & 7.5%), Ringer's solution.
- However, hypertonic solutions are considered plasma expanders as they act to increase the circulatory volume via movement of intracellular and interstitial water into the intravascular space.

Colloid:

- Colloid: High-molecular-weight solutions, draw fluid into intravascular compartment via oncotic pressure (pressure exerted by plasma proteins not capable of passing through membranes on capillary walls). Plasma expanders, as they are composed of macromolecules, and are retained in the intravascular space.
- HAES, Gelatina (Dextran);
- Albumin, Plasma

"Free H₂O solutions:"

- Free H₂O solutions: provide water that is not bound by macromolecules or organelles, free to pass through.
- D5W (5% dextrose in water), D10W, D20W, D50W, and Dextrose/crystalloid mixes.

IVF can supply 3 things

- fluid,
- electrolytes,
- calories (150 ml/h D5W)

The most common uses for IVF:

- Acutely expand intravascular volume in hypovolemic states
- Correct electrolyte imbalances
- Maintain basal hydration

infusion:

- Ringer's lactate solution
- NS = Normal Saline = NaCl 0,9%

Normal Saline (0.9% NaCl):

- Isotonic salt water.
154 mEq/L Na⁺; 154 mEq/L Cl⁻; 308mOsm/L.
- (Cheapest), commonly used crystalloid.
- High [Cl⁻] above the normal serum 103 mEq/L imposes on the kidneys an appreciable load of excess Cl⁻ that cannot be rapidly excreted. A **dilutional acidosis** may develop by reducing base bicarb relative to carbonic acid. Thus exist the risk of hyperchloremic acidosis.
- Only solution that may be administered with blood products.
- Does not provide free water or calories. Restores NaCl deficits.

Lactated Ringer's solution

- isotonic, beginning of volume resuscitation

Ingredients:

- * 130 mEq of sodium ion.
- * 109 mEq of chloride ion.
- * 28 mEq of lactate.
- * 4 mEq of potassium ion.
- * 3 mEq of calcium ion.

Lactate is converted readily to bicarb by the liver. Has minimal effects on normal body fluid composition and pH. More closely resembles the electrolyte composition of normal blood serum. Does not provide calories.

Volume

Volume deficits are best estimated by acute changes in weight. Less than 5% loss is very difficult to detect clinically and loss of 15+% will be associated with severe circulatory compromise.

- Mild deficit represents a loss of $\sim 4\%$ body wt.
- Moderate deficit --- a loss of $\sim 6-8\%$ body wt.
- Severe deficit --- a loss of $\sim >10\%$ body wt.

Volume deficit may be a pure water deficit or combined water and electrolyte deficit.

Resuscitative IV Fluids

Principle of trauma & surgery: Crystalloids; isotonic balanced salt solutions (Ringer-Lactate).

Amount given based upon body wt, clinical picture, and vital signs = shock.

Generally a **bolus of 500-2000cc** is given depending on the above, then rates are run at 1.5-2x maintenance or 10-20cc/kg/d on top of maintenance. Continuous clinical reassessment of vitals and response to fluids already given is required for ongoing IVF therapy.

Resuscitative IVF therapy is for initial stabilization and overlaps with further replacement therapy.

Monitoring endpoints for IVF therapy

- Endpoint should be maintenance or reestablishment of homeostasis.
- In order to reestablish homeostasis in a pt, IVF therapy must not only provide a **balance of water and electrolytes**, but must ensure adequate **oxygen delivery** to all organs and renal perfusion as evidenced by urine output.
 - Endpoints: normalization of VS, $UO > 0.5 \text{ ml/kg/hr}$ (1 ml/kg/hr for a child) and restoration of normal mental status and lack of clinical signs of deficit.
 - Other endpoints include normalization of labs, such as U/Cr ratio and electrolyte values.

Hypovolemia

- deficit of water
- estimated from
 - weight loss
 - thirst
 - physical signs (soft eyes, tachycardia, hypotension, oliguria, organ dysfunction – brain)
- hypo, iso, hypertonic
- Treatment: add water (crystaloid, coloid)

Hypervolemia

- hypotonic – excess of water (no ions e.g. 5% Glc)
- isotonic – anuria + intake crystalloids
- hypertonic – intake of concentrated solutions, loss of hypoosmolar fluid. / rare/

Ions in the body

- Sodium Na^+
- Potassium K^+
- Calcium Ca^{++}
- Magnesium Mg^{++}
- Phosphorus PO_4^-
- Chloride Cl^-
-
- Glucose Glc

Sodium Na⁺

- extracellular fluid 140 mmol/l
- intracellular fluid 10 mmol/l

- Hyponatremia
- Hypernatremia

Hyponatremia Na^+ in serum $< 120 \text{ mmol/l}$

- usually due to hemodilution by too much water
- sodium loss
 - vomiting
 - diarrhea
 - sweating,
 - renal / CNS disorders, diuretics
 - third space sequestration (burns, pancreatitis, peritonitis)
- factitious (hyperglycemia, hyperlipidemia, manitol) - osmolality normal / increased

Hyponatremia - symptoms

- Fatigue
 - Apathy, coma, **change in mental status**
 - Headache
 - Muscle **cramps**, weakness
 - Anorexia, nausea, vomiting,
-
- Mild to moderate hyponatremia is usually asymptomatic.

Treatment of hyponatremia

- stable pat. - water restriction (less 1l/d)
- severe, acute hyponatremia (duration < 48 h) ,
symptomatic pt. with brain edema
3% NaCl i.v.
- Rate of correction should not exceed 0.5-1 mEq/L/h, with
a total increase not to exceed 12 mEq/L/d
- Risk of rapid treatment - demyelination

Hypernatremia

- inadequate water intake
- excessive loss of water
 - diarrhea
 - vomiting
 - hyperpyrexia
 - excessive sweating
 - diabetes insipidus (ADH) = loss of hypotonic urine
- increased intake of salt
- coma, no response to thirst

Therapy: Glc 5% i.v.

Potassium K^+

- Major intracellular cation
- serum (2% of total) 3.8 .. 5.6 mmol/l
- electric potential on membrane (Na^+/K^+ ATPase)
- arytmiias

- extremely responsive to changes of pH!!

Acidosis in cell (H^+) banish K^+ out of cell.


delta pH 0,1 0,5..0,6 mmol/l (i.v.)

Hypokalemia $K < 4 \text{ mmol/l}$

- losses in urine
- diuretics, diarrhea, vomiting
- reduced intake
- Alkalosis
- CAVE severe muscle weakness, asystolia


Treatment:

- KCl p.os; max KCl 40 mmol/h i.v.
- ECG monitoring !!!!



Hyperkalemia

- hemolysis
 - muscle damage
 - anuria, renal failure
 - Acidosis
 - CAVE intracardiac block (diastolic arrest) or fibrillation
 - muscle weakness – ventilatory failure
- therapy:
- stop intake
 - Glc + HMR i.v., loop diuretic (furosemide)
 - Calcium i.v., bicarbonate i.v
 - resonium p.os
 - dialysis





Calcium Ca^{++}

- most abundant mineral in the body 2kg
- Parathormone PTH
 - stimulate osteoklast
 - stimulate intestine
 - resorption in kidney
- Calcitonin
 - inhibites osteoklast
- Vitamine D
 - potens saving Ca^{++}

Ionised $\text{Ca} = 1.1 \text{ mmol/l}$ // efect of all Calcium
bound by proteins =ineffective to receptors

Calcium Ca^{++}

- Hypocalcemia
 - Respiratory Alkalosis, hypoPTH,
 - shock, sepsis, pancreatitis
 - together hypomagnesemia
- Hypercalcemia
 - muscle damage
 - malignancy




Chloride Cl⁻

- Major anion in Extracellular fluid
- see ABR

Acide-base

arterial blood:

pH	7,35-7,45
pCO ₂	4,6-6 kPa
pO ₂	10-13 kPa
HCO ₃ ⁻	22-26mmol/L
BE	-2 .. +2 mmol/L
SpO ₂	95-98%



Basic laws

$$\text{pH} = -\log [\text{H}^+] \quad [\text{H}^+] \dots \text{mol/l}$$

$$\text{pH} = \text{pK} + \log (\text{H}^+ \text{ acceptor} / \text{H}^+ \text{ donor})$$

$$[\text{H}^+] = 24 \times \text{pCO}_2 / [\text{HCO}_3^-] \quad \text{Henderson equation}$$


- acidosis $\text{pH} < 7.36$
- alkalosis $\text{pH} > 7.44$

Place of error:

- Respiratory (lung) $\dots \text{pCO}_2$
- Metabolic (kidney,...) $\dots \text{BE}$

$\text{BE} =$ number of acid needed to correct sample to $\text{pH} 7.4$

CO₂



ΔpH

0.1

$\Delta p \text{ CO}_2$

1,6 kPa = 12 mmHg

Kilopascals for PCO₂.

- Many texts and papers express the PCO₂ in kilopascals (kPa). It is useful to remember that this value is almost the same as the percentage of atmospheric pressure. For example, the normal arterial PCO₂ of 40 mmHg is 5.33 kPa or 5.61 %.
- To convert pressure in mmHg to kPa, it is necessary to divide the value in mmHg by 7.5
- [40mmHg /7,5 = 5,33kPa]

Genesis of Acid = donor of H⁺


- lactate - shock
- strong acids intake (HCl, H₂SO₄)
- acetylsalicylic acid (drug overdose)
- ...


Δ pH

0.1

Δ BE

6mmol/l





pH
7.35 - 7.45
Acidosis \leftrightarrow Alkalosis

If PCO_2 has an indirect relationship to the pH the condition is a Respiratory imbalance.

PCO_2
35 - 45

Normal?

HCO_3^-
22 - 26

Does the HCO_3^- have a direct relationship with the pH? \rightarrow then condition is Metabolic Imbalance


Resp. Imbalances
Look @ HCO_3^- to determine state of compensation. If abnormal = partial compensation

Compensation
Has occurred if pH is in normal range

Metabolic Imbalances
Look @ PCO_2 to determine state of compensation. If abnormal = partial compensation

Respiratory Acidosis ($\downarrow \text{pH}_a$ & $\uparrow P_{a\text{CO}_2}$)

Hypoventilation



Causes: Anoxia, COLD, near drowning, strangulation

Fig. 17-9

Easy equation

	Resp. PCO2	Meta HCO3
Alka ↑		
Acid ↓		

ΔpH

0.1

BE

6mmol/l

$\Delta p \text{ CO}_2$

1,6 kPa = 12 mmHg

Easy equation

	Resp. PCO2	Meta HCO3
Alka ↑	↓	↑
Acid ↓	↑	↓

↓ ΔpH
0.1

↓ BE
6mmol/l

↑ $\Delta p \text{ CO}_2$
1,6 kPa = 12 mmHg

Respiratory Acidosis (RAc)

- The **decision to ventilate** a patient to reduce the PCO_2 is a clinical decision and is based on exhaustion, prognosis, prospect of improvement from concurrent therapy, and in part on the PCO_2 level. Once the decision is made, the PCO_2 helps to calculate the appropriate correction.
- The PCO_2 reflects a **balance** between the carbon dioxide production and its elimination. Unless the metabolic rate changes, the amount of carbon dioxide to be eliminated remains constant. It directly determines the amount of ventilation required and the level of PCO_2 . Where V_T equals tidal volume and f equals respiratory rate:
- $PCO_2 \times \text{Ventilation} = \text{Constant}$, i.e.,
 $PCO_2 \times V_T \times f = k$

MAc

- **kidney unable to eliminate H^+ = anuria**
- **big production of acides.**
- The treatment for a metabolic acidosis is, again, judged largely on clinical grounds. Bicarbonate therapy is justified when metabolic acidosis accompanies difficulty in resuscitating an individual or in maintaining cardiovascular stability.
- A typical dose of bicarbonate might be 1 mEq per kilogram of body weight followed by repeat blood gas analysis.
- Calculation is based on BE and the size of the treatable space (0.3 x weight, e.g., 21 liters):
Dose (mEq) = 0.3 x Wt (kg) x BE (mEq/L).

Serum Anion Gap

Cations

Na^+ , K^+ , Ca^+ , Mg^+

Anions

Cl^- , HCO_3^- , PO_4^- , SO_4^- , Albumin,
organic acids (OA)

$$[\text{Na}^+] + \text{UC} = [\text{Cl}^-] + [\text{HCO}_3^-] + \text{UA}$$



$$\text{UA} - \text{UC} = [\text{Na}^+] - ([\text{Cl}^-] + [\text{HCO}_3^-])$$

$$\text{Anion gap} = [\text{UA}] - [\text{UC}] = [\text{Na}^+] - ([\text{Cl}^-] + [\text{HCO}_3^-])$$

UC, unmeasured cations; UA, unmeasured anions


Metabolic Acidosis

- Unmeasured Anions {organic acids - lactate, ...}
- Cl⁻
- loss of HCO₃, Na, K, Cl
 {diarrheal fluid, wrong ratio}



RA1

- hyperventilation
- lost of ionized Calcium / hypocalcemia / tetania



MA1

- increased loss of NH_4 to urine
- saving HCO_3^- by kidney
- loss of Cl^- (vomiting)
-
- $\text{BE} > 0$
- $\text{pH} > 7.44$
- Th: i.v. FR (NaCl)
-

How to

1. what is wrong
2. what the body do
3. what to do

OR / AAA, 5 000ml lost, haemorrh. shock, NA i.v.,
general anesthesia, VCV

pH akt.	7.083	(7.350 - 7.450)	<-()
pCO ₂	6.36 kPa	(4.80 - 5.90)	()->
pO ₂	30.78 kPa	(10.66 - 13.30)	()=>
BE	-15.8 mmol/l	(-2.6 - 2.6)	<=()
BB	32.1 mmol/l	(40.0 - 44.0)	<=()
HCO ₃ akt.	13.9 mmol/l	(22.0 - 26.0)	<=()
O ₂ sat.	99.3	(95.0 - 98.0)	()=>

OR / AAA, 6 500ml loost, haemorh. shock, NA i.v.

pH akt.	7.1	(7.350 - 7.450)	\leq ()
pCO ₂	5.0 kPa	(4.80 - 5.90)	(*)
BE	-18 mmol/l	(-2.6 - 2.6)	\leq ()
lactate	13 mmol/l	(1 - 2.5)	() \Rightarrow

Try it yourself

pH = 7,2

pCO₂ = 14 kPa

BE = 20 mmol/l

pH 7,35-7,45

pCO₂ 4,6-6 kPa

pO₂ 10-13 kPa

HCO₃⁻ 22-26mmol/L

BE -2 .. +2 mmol/L

SpO₂ 95-98%

- polytrauma + sepsis + ARDS

Measured		Calculated	
Na	131 ↓	HCO ₃ ⁻	21 ↓
K	4,2 =	BE	-4 ↓
Mg	3,6 ↑		
Ca	2,2 =		
Cl	86 ↓		
Pi	2,3 ↑		
Alb	8 ↓↓		
pH	7,31 ↓		
PaCO ₂	5,4 =		

- polytrauma + sepsis + ARDS

Henderson-Hasselbach:

- metabolic acidosis

Measured		Calculated	
Na	131 ↓	HCO ₃ ⁻	21 ↓
K	4,2 =	BE	-4 ↓
Mg	3,6 ↑		
Ca	2,2 =		
Cl	86 ↓		
Pi	2,3 ↑		
Alb	8 ↓↓		
pH	7,31 ↓		
PaCO ₂	5,4 =		

- polytrauma, sepsis ARDS

Measured		Calculated	
Na	131 ↓	HCO ₃ ⁻	21 ↓
K	4,2 =	BE	-4 ↓
Mg	3,6 ↑		
Ca	2,2 =		
Cl	86 ↓		
Pi	2,3 ↑		
Alb	8 ↓↓		
pH	7,31 ↓		
PaCO ₂	5,4 =		

Stewart-Fencl:

- lactic acidosis
- dilution acidosis
- hypochloremic alkalosis
- hypoalbuminemic alkalis


#3 example

• pH 7.45 — N

• PCO₂ 43 — N

• HCO₃⁻ 25 — N

• this is the hardest one
for beginners... it's normal



SUMMARY

- Biologic system react primary to rate of change and not to absolute concentrations.
- Abnormalities should be treated at proximately the rate at which they developed.
- DO NOT rapid correction of a chronic asymptomatic abnormality.

When order electrolytes exam:

- poor oral intake
- vomiting
- chronic hypertension
- diuretic use
- recent seizure
- muscle weakness
- age over 65
- alcoholism
- history of electrolyte abnormality

When order blood gasses:

- acid-base problems
- artificial ventilation
-

acute CNS change

immediately look for

- hypoxemia
- hypoglycemia
- hyponatremia
- sepsis

Priorities

1. fluid volume and perfusion deficits
2. correction of pH
3. K, Ca, Mg
4. Na, Cl

Bleeding – transfusion strategy

Indication:

- Transfuse any symptomatic patient (e.g., tachycardia, hypotension, CHF, angina)
- Asymptomatic, presurgical, stable patient
- Hemodynamically stable postsurgical stable patient
- Postsurgical patient at risk for ischemic disease (e.g., cardiac, bowel)
- Hemodynamically stable, nonpregnant, ICU patients >age 16 without ongoing blood loss

Transfuse to Maintain:

- Until no longer symptomatic
- Hb 7-8 g/dl
- Hb 8 g/dl
- Hb 10 g/dl
- Transfuse at 7 g/dl to maintain Hb at 7-9 g/dl