





TELENCEPHALIC CORTEX - NOMENCLATURE



paleocortex, archicortex (allocortex) – neurons in 3-5 layers


neocortex (isocortex) – neurons in 6 layers


peripaleocortex et periarchicortex = mesocortex



















LOCATION OF MAIN FUNCTIONAL REGIONS OF TELENCEPHALIC CORTEX

- somatosenzory and senzory
- motor
- cortex of limbic forebrain
- association cortex



Brodmann's area	cortical location	functional involvement
a 3, 2, 1	postcentral gyrus	analysis of the somatosensory information
a 4, 6	precentral gyrus	primary motor cortex
a 41, 42	gyri temporales transversi	analysis of the hearing
a 17	cortex parallel with the calcarine sulcus	primary visual cortex
a 18, 19	cortex parallel with a 17	secondary visual cortex
a43	caudal part of the postcentral gyrus	analysis of the taste




CORTICAL AREAS FOR SPEECH - I

Broca's (motor) cortical area - g. front. inf. a44, 45


for right-hander in L-hemisphere, for left-hander in R-hemisphere



Wernicke's (sensory) cortical area - a 22,39,40 in dominant hemisphere


CORTICAL AREAS FOR SPEECH - II




A SCHEMATIC DRAWING OF 3D ARRANGEMENT OF SOME TELENCEPHALIC STRUCTURES










Limbic system – classic conception

Papez's circuit (James Papez 1939)




RECENT CONCEPTION OF LIMBIC FOREBRAIN

- basomedial telencephalon, structures of diencephalon and mesencephalon for emotion and motivation of our behavior

Regular structures

- g. cinguli, g. parahippocampalis, hippocampus, *insular cortex*
- neocortical regions of forebrain - basal frontotemporal regions, orbital cortex, ventral striatum (pallidum)
- *area septalis*, amygdalar ncl., hypothalamus (ncl. mammillaris)
- ncl. anterior et *medialis dorsalis* thalami
- *habenulla*

Limbic system – classic conception Papez's circuit (James Papez 1939)




notion of tooth pain

Cingulate gyrus

Anterior nucleus of thalamus



notion of fear

**memory of
listening to
music**



AMYGDALAR NUCLEI




A**B****C**

Septal nuclei

Amygdala

Fornix

Hippocampus



BASAL GANGLIA AND RELATED STRUCTURES

ncl. caudatus, putamen, globus pallidus, claustrum and amygdalar ncl.


functional: + thalamus, substantia nigra and ncl. subthalamicus

ncl. caudatus + putamen = **neostriatum (striatum)**


globus pallidus (ext. + int. segment) = **paleostriatum (pallidum)**

globus pallidus + putamen = **ncl. lentiformis**


BASAL GANGLIA AND RELATED STRUCTURES





VENTRAL PALIDUM AND VENTRAL STRIATUM



SUBSTANTIA INNOMINATA a NCL. ACCUMBENS





WHITE MATTER OF TELENCEPHALON

Pathways - associated, projection and commissural

ASSOCIATED PATHWAYS - interconnections of various cortical regions

fasciculus longitudinalis superior


fasciculus longitudinalis inferior



fasciculus occipitofrontalis superior

fasciculus uncinatus

fasciculi occipitales verticales

cingulum





Projection pathways

Short projection pathways

Long projection pathways - *capsula interna*

crus anterius, genu et crus posterius capsulae internae

CAPSULA INTERNA

crus anterius – anterior tr. thalamo-corticalis and tr. fronto-pontinus

genu - tr. cortico-nuclearis, from area 4 to contralateral motoneurons of cranial nerves

crus posterius - tr. cortico-spinalis (somatotopic arrangement), tr. cortico-reticularis and tr. cortico-rubralis, posterior tr. thalamo-corticalis (somatosenzory information to parietal cortex), tr. parieto- , temporo-, occipito-pontinus, radiatio optica, radiatio acustica

