



Pathophysiology of GIT II

Exocrine pancreas
Liver
Biliary tract



1


Pathophysiology of exocrine pancreas



2

Pancreas - secretion


- endocrine part (2%)
 - insulin, glucagon, somatostatin, gastrin, pancreatic polypeptide, amylin
- exocrine part (85%) - acines
 - pancreatic juice (pH up to 8.3)
 - approx. 1-1.5l day
 - production stimulated by acetylcholine, CCK and secretin produced in duodenum
 - production inhibited by pancreatic polypeptide
 - composition
 - ions and water (← secretin)
 - Na^+ , Cl^- , K^+ and HCO_3^- (up to 150 mmol/l)
 - HCO_3^- necessary to neutralize acid content of stomach, for activation of pancreatic enzymes and formation of micelle
 - enzymes (← CCK)
 - active - lipase, amylase, ribonuclease, deoxyribonuclease
 - inactive (activated by enterokinase in duodenum) - trypsinogen, chymotrypsinogen, procarboxypeptidase, proelastase, phospholipase
 - inhibitory trypsin ($\alpha 1$ -antitrypsin)
- disorder of secretion - insufficiency of exocrine pancreas
 - most often due to chron. pancreatitis
 - carcinoma of pancreas, cystic fibrosis, protein malnutrition



3

Chronic pancreatitis


- chronic inflammation of pancreas leading to progressive dysfunction of pancreatic acins, stenosis and dilation of ducts, fibrosis and atrophy of gland and calcifications in ducts
- etiology
 - hypertriglyceridemia
 - hypocalcaemia
 - chron. malnutrition
 - alcoholism
 - tropical form
 - hereditarily
 - cystic fibrosis
- consequences
 - absence of lipase
 - maldigestion and malabsorption of fats (→ steatorrhea, diarrhea)
 - deficiency of lipid-soluble vitamins
 - absence of amylase and peptidases
 - mostly compensated by stomach and intestinal enzymes, malabsorption of sugars and AA thus clinically insignificant
 - hypocalcaemia and hyperphosphatemia (due to ↓ vit. D) → osteomalacia
 - deficit of vit. B12 (due to deficit of protease its release from dietary sources low) → anemia
 - pain
 - secondary diabetes mellitus (destruction of islets of Langerhans)
- complications
 - cysts, closure of ducts, leak of juice to peritoneal and pleural cavity



4

Cystic fibrosis (mucoviscidosis)


- monogenic (AR) disease due to mutation in gene encoding "cystic fibrosis transmembrane conductance regulator" (CFTR)
 - >600 known mutations in one of the 4 classes
 - I – defective protein (preterm stop of translation of CFTR mRNA)
 - II – increased degradation of protein in endopl. reticulum (incl. the most common mutation $\Delta F508 \sim 70\%$)
 - III – inactivated channel
 - IV – defect of transport
- function of CFTR
 - encodes a complex protein forming chloride channel
 - regulates other channels (e.g. Na)
- CF affects
 - epithelia of respiratory tract
 - viscous secret, limitation of respiration and coughing, terrain for infection (*Pseudomonas aeruginosa*) → chron. bronchitis, bronchiectasis, pneumonia
 - epithelia in pancreatic ducts
 - recycling of Cl involved in secretion of HCO_3^- into pancreatic juice → due to decreased bicarbonate too viscose protein secret blocking ducts(chron. pancreatitis)
 - sweat glands
 - decreased reabsorption of Cl (diagnostic sign - high Cl in sweat)
 - intestine
 - meconic ileus of newborns
 - liver and biliary tract
 - genitals



5

Acute pancreatitis


- acute destruction of pancreatic tissue and neighboring tissue due to autodigestion by pancreatic enzymes activated directly in the gland
- very serious and severe condition associated with high mortality
- symptoms
 - intensive pain
 - nausea and vomiting
 - fever
- etiology
 - biliary
 - blockage by bile stone in common duct
 - alcohol
 - relaxation of sphincter of Oddi
 - reflux of bile into pancreatic duct
 - abdominal trauma
 - infection
 - hypertriglyceridemia
 - hypercalcaemia
 - drugs
- pathogenesis
 - intracellular and extracellular activation of trypsinogen and subsequently of other enzymes
 - cathepsine B in low pH
 - autodigestion of gland
 - elastase digests elastin in vessel walls → hemorrhage into gland, leak of juice into circulation and damage of systemic circulation
 - lipolysis of pancreas by pancreatic lipase and phospholipase A2



6


Tumors of pancreas

- most commonly adenocarcinoma
 - ↑ risk
 - chron. pancreatitis
 - smokers
 - chron. alcoholism
 - typically head and body, less often caudal pancreas
 - signs
 - obstructive icterus (compression of biliary duct)
 - pancreatic insufficiency
 - thrombophlebitis
 - very poor prognosis
- tumors of endocrine pancreas
 - insulinoma (hypoglycemia)
 - gastrinoma (Zollinger-Ellison syndrome)
 - VIPoma (diarrhea, hypokalemia)
 - carcinoid



7


Anatomy and histology of liver




- liver (hepar) ~1.5kg
- 2 lobes (sin. and dx.) divided by ligament
- liver parenchyma has characteristic architecture
 - liver lobule is a basic morphologic unit
 - central vein lobule
 - peripheral portobiliar "trias"
 - liver acinus is basic functional unit
 - part of the tissue supplied by branches of one circumlobular vein
- function of liver
 - complex metabolic function
 - saccharides
 - glycogen synthesis, glycogen lysis, gluconeogenesis
 - lipids
 - clearance of lipoproteins, synthesis of cholesterol, synthesis of TAG
 - proteins
 - trans- and de-amination of AA, protein synthesis (albumin, clotting factors)
 - formation of bile
 - metabolisms of haem
 - biotransformation, detoxification
 - hormones, drugs, toxins, ammoniac from intestine
 - storage of vitamins and trace substances

8

Liver blood supply




- v. portae (80% of blood supply)
 - drainage from splanchnic organs (= functional supply)
 - capillaries from stomach, intestine, pancreas and spleen connect in portal vein
 - its branches encircle liver lobules (v. interlobulares and circumlobulares)
 - they enter them as liver sinusoids
 - sinusoids join to form central vein
- a. hepatica (20% of supply)
 - branch of truncus coeliacus (= nutritional supply)
 - drain to sinusoid and then to the central vein
- v. hepatica
 - drainage from liver
 - central veins connect to right and left liver vein leading to lower vena cava




9

Liver blood supply







10

Morphology of liver - lobule





11

Liver lobule schematically





12

Liver lobule vs. acinus



13


Hepatocytes, liver sinusoids



14


Etiology of liver damage

- infection
 - viral
 - hepatitis viruses (HAV, HBV, HCV, ...)
 - inf. mononucleosis (EBV)
 - bacterial
 - leptospirosis
 - parasite
 - Echinococcus
 - globally, Europe - Mediterranean
 - Schistosomiasis (= bilharzias)
 - Africa, J. America, Caribbean, SE Asia
 - malaria
- toxic
 - alcohol
 - faloidin (Amanita faioides)
 - drugs (e.g. paracetamol)
 - chemicals
- autoimmune
 - autoimmune hepatitis
 - prim. biliary cirrhosis
- metabolic disorders
 - common - NAFLD/NASH
 - rare
 - heredit. hemochromatosis
 - Wilson disease
 - porphyria
 - glycogenosis
- tumors
 - primary (hepatocellular carcinoma)
 - metastases



16



REACTION OF LIVER TO DAMAGE & LIVER DISEASES



15

Liver infection - hepatitis

- time course
 - acute
 - usually without residual damage
 - fulminant form leading to liver failure
 - chronic
 - only persistent infection (carriers)
 - necrosis of parenchyma and progression to cirrhosis
- viral hepatitis
 - hepatitis A (HAV – RNA virus)
 - only acute time course
 - virus directly cytotoxic
 - epidemic
 - fecal-oral transmission (vaccination)
 - hepatitis B (HBV – DNA virus)
 - blood borne (parenteral) and STD
 - time course
 - virus is not directly cytotoxic, damage is the results of the reaction of immune system
 - mostly acutely without residual damage
 - in 10% of cases progresses to chronicity
 - either solely HBsAg positive carriers
 - or active process leading to fibrosis and cirrhosis
 - hepatitis C (HCV – RNA virus)
 - blood born (parenteral) and STD
 - acute phase typically asymptomatic
 - more than 80% cases progress to chronicity – can lead to cirrhosis



17

Hepatitis Viruses

Virus Family	Hepatitis A Picornavirus	Hepatitis E Calicivirus	Hepatitis B Hepadnavirus	Hepatitis C Flavivirus	Delta virus Satellite virus (only in combination with HBV)
Commonality	All generate conditions of illness in the liver				
Symptoms (acute)	All the same – malaise, dark urine, anorexia, nausea, vomiting, jaundice				
Transmission	Enteric (food and water)		Sex, blood and close contact		
Chronic condition	No	No	Yes	Yes	Yes
Virus genome	+ss RNA	+ss RNA	DNA with reverse transcriptase activity	+ss RNA	-ss RNA
Virus antigens	HA Ag	HEV ORF2 proteins	HBsAg HBcAg HBeAg	Many – core E1 E2 NS3	Delta antigen
Incubation	1 month (15 – 50 d)		4 months (45 – 160 d)	2 months (15 – 150 d)	1 – 2 months
Current therapeutics	No specific treatment	No specific treatment	Interferon alpha, Lamivudine, Adefovir, Etecovir	Interferon alpha + ribavirin, Pegylated Interferon	Follow HBV therapy
Vaccines available?	Yes Havix (GSK) Vacta (Merck)	No	Yes Engerix-B (HBsAg) GSK Recombivax B (Merck)	No	Can be prevented by vaccination against HBV

18

Reaction of liver to damage





(Painting: Christian Griepenkerl's Prometheus Tortured by the Eagle, 1800's)

19

Reaction of liver to damage

- liver can react the same way to various etiologies of damage
 - mild damage change metabolic activity of hepatocyte, which become to cumulate fat (= **steatosis**)
 - steatosis with lab. signs of inflammation is called **steatohepatitis**
 - more severe damage leads to cell death, however liver has a considerable ability to regenerate
 - long-term damage leads to production of connective tissue in periportal areas (= **fibrosis**)
 - combination of intensive necrosis, fibrosis and regeneration significantly altering lobular architecture is called **cirrhosis**




20

Initial (reversible) liver damage


steatosis (S)

- normally fat content (TAG) in hepatocytes <5%
 - histologically microvesicular or macrovesicular
- causes
 - excessive dietary intake or lipolysis in adipose tissue
 - increased endogenous synthesis
 - decreased catabolism in liver
 - combination
- steatosis itself is not harmful for liver (sometimes is even considered protective mechanisms), however it represents substrate for increased lipid peroxidation



steatohepatitis (SH)

- together with S also necrosis, inflammation and fibrosis
- more serious than simple S (which is reversible when causing factor ceases)
- it can reverse to normal or progress to fibrosis or cirrhosis
- transition of S to SH enhanced by other factors such as oxidative stress, endotoxin, immune system, nutrition etc.




etiology S a SH

- alcoholic**
 - energetic content of alcohol
 - alteration of intermediary metabolism
 - inhibition of β -oxidation
 - \uparrow NADH and acetyl-CoA (\uparrow synthesis FFA)
- non-alcoholic fatty liver disease (NAFLD) and steatohepatitis (NASH)**
 - component of insulin resistance syndrome
 - \uparrow lipolysis in adipose tissue - \uparrow uptake of FFA by liver
 - \uparrow peroxidation of lipids and ox. stress for hepatocytes
 - hyperinsulinemia stimulates synthesis of FFA and TAG

21

NAFLD and NASH


- prevalence ~20 - 30% in industrialised countries (associated with **OBESITY!!!**)
- can be difficult to dissect non-alcoholic from alcoholic damage in countries where alcohol consumption is socially accepted and common
 - definition of non-alcoholic etiology: daily intake <10g/day in men (i.e. ~140g ethanol per week) and (~70g ethanol in women)
- pathogenesis of NAFLD/NASH = metabolic alterations resulting in hepatic triglyceride accumulation in **insulin-resistant states**
 - insulin resistance is manifested by **hyperinsulinemia**, increased hepatic glucose production, and decreased glucose disposal
 - in adipocytes, hyperinsulinemia increases hormone-sensitive lipase (HSL) activity, resulting in elevated rates of triglyceride lipolysis and **enhanced FFA flux to the liver**
 - FFAs can either be oxidized in the mitochondria to form ATP or esterified to produce triglycerides for storage or incorporation into VLDL particles
 - in liver, hyperinsulinemia induces SREBP-1c and ChREBP expression, leading to the transcriptional activation of all lipogenic genes and the enzymatic machinery necessary for the **conversion of excess glucose to fatty acids**
 - a consequence of increased fatty acid synthesis is increased production of malonyl-CoA, which inhibits CPT-1, the protein responsible for fatty acid transport into the mitochondria
 - thus, in the setting of insulin resistance, FFAs entering the liver from the periphery, as well as those derived from de novo lipogenesis, will be preferentially esterified to triglycerides.
 - ACL, ATP citrate lyase; CPT-1, carnitine palmitoyl transferase-1; FAS, fatty acid synthase; LCE, long-chain fatty acyl elongase
- NAFLD represent good terrain for lipid peroxidation due to oxidative stress
 - \uparrow ox. stress in ins. resistance (\uparrow resistin, TNF α , IL-6 and other pro-inflammatory adipokines)
- products of lipid peroxidation - malondialdehyd (MDA) or 4-hydroxynonenal (HNE) - stimulate Kupffer and HSC to fibroproduction and chemotaxis of neutrophils



22


More advanced (reversible?) liver damage

- result of chronic damage of hepatocytes
 - infection, alcohol, toxic substances, accumulation of metals (Cu, Fe), drugs, ...
- collagen in normal liver
 - I and III in periportal areas
 - IV in Disse space
- fibrosis (F)** = increased content of connective tissue
 - damaged hepatocyte activate Kupffer cells which release paracrine factors (PDGF and TGF- β)
 - activation of hepatic stellate cells (HSC)
 - regulation of blood flow through sinusoids (\uparrow resistance)
 - synthesis of connective tissue (collagen, laminin, ...)
 - release of photolytic enzymes (matrix-metalloproteinases)
 - alteration of morphology of sinusoids (loss of fenestrations of endothelia), accumulation of extracel. matrix




23

Activation of HSC in fibrosis /cirrhosis



- HCS activated by growth factors from damaged hepatocytes and Kupffer cells
- synthesis of collagen I and III in Disse space
- loss of microvilli of hepatocytes
- loss of fenestration of sinusoids (= capillarisation of sinusoids)



Role of HSC in I. fibrosis on hepatic sinusoidal cells



25


Advanced (irreversible) liver damage

- **cirrhosis (C)**
 - histologically micronodular or macronodular
 - irreversible change of architecture (lobules, vessels, collagen)
 - parallel processes:
 - fibrosis + necrosis + nodular regeneration
 - loss of functional parenchyma
 - portal hypertension and liver failure
 - ↑ risk of carcinoma
- general symptoms of advanced liver diseases
 - weakness, weight loss
 - jaundice
 - bleeding
 - due to deficit of clotting factors
 - edema, ascites
 - due to hypoalbuminemia
 - prolonged action of hormones
 - gynecomastia in men
 - spider nevi
 - liver encephalopathy
 - due to hyperammonemia




26

Alcohol and liver - endotoxin




- alcohol increases permeability for endotoxin from intestine to circulation
 - endotoxin is a part of the G-negative bacteria wall
- endotoxin (via receptors CD14 and TLR4) activates Kupffer cells (specialized macrophages along liver sinusoids) to production of cytokines (NFκB) and superoxide (NADPH oxidase)

27



28

Pathophysiological differences between normal (A) and fibrotic/cirrhotic (B) liver




- In normal liver (A), normal fenestrae along the hepatic sinusoids allow free passage of blood (arrows) into the Space of Disse, in which, stellate cells (green) are found. In liver cirrhosis (B), there is an increase in the number of stellate cells, associated with deposition of collagenous fibers in the Space of Disse, and loss of fenestrae as the sinusoids become more capillary-like. As a result, transfer of low-molecular-weight compounds (e.g. contrast medium) from the sinusoids into the Space of Disse becomes more impeded (small arrows).

29


Consequences of liver cirrhosis

- portal hypertension
- hypoalbuminemia
- disorder of hemostasis
 - vitamin K deficit and thus inadequate formation of clotting factors
- suppression of bone marrow
 - due to bleeding, hypersplenism and low K vitamin resorption
- hyperbilirubinemia or icterus
- decreased degradation of circulating hormones
 - aldosterone
 - loss of K by urine, intracel. acidosis, **metabolic alkalosis**
 - decreased ionization of NH_3 !!!!
 - androgens – increased conversion to estrogens in periphery
 - gynecomasty in men
 - pavlovčkové névy
- metabolic consequences
 - abnormal metabolism of AA (\uparrow conc. of aromatic AA – atyp. neurotransmitters in CNS)
 - disorder of glucoregulation
 - impaired urea cycle
- intrahepatic cholestasis




30

Hyperbilirubinemia/icterus



31


Portal hypertension



- normal pressure in portal circulation 5 – 15 mmHg
- localization of portal hypertension
 - pre-hepatic
 - thrombosis v. portae, malformation, compression
 - intra-hepatic
 - due to cirrhosis, parasites
 - post-hepatic
 - right heart failure (hepatosplenomegaly), thrombosis of liver veins (Budd-Chiari syndrome), compression by tumour
- increased pressure before liver sinusoids does not create pressure overload for liver, after sinusoids it does, therefore damage is greater

32


Portal hypertension



- 1) congestion of blood in the v. portae and stasis of blood in splanchnic organs
 - stomach and intestine
 - malnutrition and maldigestion
 - erosion and ulcers
 - increased permeability for bacteria
 - spleen
 - hypersplenism → destruction of Ery and platelets
- 2) blood flow through portocaval anastomoses directly to systemic circulation
 - normally there are small veins
 - under the high pressure risk of mechanical damage and bleeding
 - vv. oesophageae (esoph. varices)
 - vv. rectales (hemoroids)
 - vv. paraumbilicales (caput Medusae)
- 3) ascites and edemas
 - fluid in peritoneal cavity due to portal hypertension + hypoalbuminemia + retention of Na (aldosterone)
 - increased permeability for bacteria = spontaneous bact. peritonitis
- 5) hepatorenal syndrome

33


Esophageal varices



34


Liver encephalopathy

- abnormalities of conscience (quantitative and qualitative), behavior and neuromuscular functions
 - reversible only in initial stages
- impaired detoxification of ammonia in urea cycle
 - sources of ammoniac
 - oxidative de-amination by glutamatdehydrogenase from Glu
 - glutaminase from Gln to Glu
 - degradation of purines and pyrimidines
 - de-amination by monoaminoxidase
 - synthesis of hem
 - bacteria in large intestine
 - ammoniac >50μmol/l toxic for CNS
 - in blood as NH₃/NH₄⁺
 - balance depends on pH (normally 99% ionised)
 - alcalosis increases free ammoniac and thus toxicity
 - urea (= ornithin) cycle in liver daily produces 20 – 40 g urea
 - CO₂ + NH₄⁺ → CO(NH₂)₂ + H₂O + 2H⁺
 - 5 enzymes – mitochondria and cytosol
 - urea excreted by kidney
- blood from splanchnic contains not only nutrients by also toxins (ammoniac, mercaptans, phenols etc. produced by bacteria)
- if not properly detoxified in liver
 - formation of "false" neurotransmitters in brain
 - change of behavior and conscience, "flapping" tremor, apraxia




35

Intestine and liver - ammonia



36


Impaired balance of excitatory and inhibitory AA in the brain



37

Hepatorenal syndrome

- kidney failure accompanying liver disease without pre-existing kidney pathology
- etiology
 - Na and water retention
 - hyperaldosteronemia
 - however, effective circulating volume is decreased due to escape to the third space (ascites)
 - hypoalbuminemia
 - decrease of renal perfusion and GFR
 - systemic vasodilation but intrarenal vasoconstriction
 - contraction of afferent arterioles (RAS)



38

Liver tumors


- benign
 - hemangioma
 - hematoma
- malign
 - hepatocellular carcinoma
 - in 70% consequence of cirrhosis
 - prevalence increases
 - poor prognosis
- metastases
 - colorectal carcinoma, ...



39

Pathophysiology of biliary tract

- cholecystolithiasis (gallstones)
 - typically 55-65 yrs ~10% men and ~20% women
 - causes – alteration of the ration between bile components
 - type of stones
 - cholesterol (70-90%)
 - pigmented (calcium + bilirubin)
 - mixed
 - increased concentration of cholesterol
 - diet, obesity
 - decrease of bile acids and phospholipids
 - malnutrition, Crohn disease, resection of ileum
 - cholecystitis
 - stagnation of bile
 - diet, starvation
- complications of cholecystolithiasis
 - biliary colic (blockade of d. cysticus)
 - extrahetal cholestasis (blockade of d. choledochus)
 - inflammation (cholecystitis, cholangitis)
 - acute pancreatitis



40



Cirrhosis of the river.