

Pathophysiology of the respiratory system I

- Structural properties of airways and lungs
- Defense mechanisms of the respiratory system
- Respiration and gas exchange
 - ventilation & diffusion & perfusion
- Pulmonary mechanics
- Ventilation – perfusion (in)equality
- Control of ventilation




Figure 17.1 Diagrammatic view of human respiratory system (Sectional view of the left lung is also shown)

The delicate structure-function coupling of lungs

- The main role of the respiratory system is to **extract oxygen from the external environment and dispose of waste gases**, principally carbon dioxide
 - at the end of deep breath 80% of lung volume is air, 10% blood and 10% tissue
 - lung tissue spreads over an enormous area !
- The lungs have to provide
 - a **large surface area** accessible to the environment (~tennis court area) for gas exchange
 - alveoli walls have to present **minimal resistance to gas diffusion**
- Close contact with the external environment means lungs can be damaged by dusts, gases and infective agents
 - **host defense** is therefore a key priority for the lung and is achieved by a combination of structural and immunological means




Structure of airways

		Generation		
Conducting airways	Trachea	0		
	Bronchi	Cartilage, bronchial glands	1	
			2	
			7	
			8	
		Bronchioles	Nonrespiratory	9
				15
				16
	17			
Terminal respiratory units	Respiratory	18		
		19		
		20		
		21		
		22		
		Alveolar ducts	23	

- There are about 23 (18-30) divisions (2^{23} i.e. approx. 8 millions of sacs) between the trachea and the alveoli
 - the first seven divisions, the bronchi have:
 - walls consisting of cartilage and smooth muscle
 - epithelial lining with cilia and goblet cells
 - submucosal mucus-secreting glands
 - endocrine cells - Kulchitsky or APUD (amine precursor and uptake decarboxylation) containing 5-hydroxytryptamine
 - the next 16-18 divisions the bronchioles have:
 - no cartilage
 - muscular layer progressively becomes thinner
 - a single layer of ciliated cells but very few goblet cells
 - granulated Clara cells that produce a surfactant-like substance


Source: McPhee SJ, Ganong WF: *Pathophysiology of Disease: An Introduction to Clinical Medicine*, 5th Edition: <http://www.accessmedicine.com>
Copyright © The McGraw-Hill Companies, Inc. All rights reserved.

Wall structure of conducting airways and alveolar region




Alveoli

- There are approximately 300-400 million alveoli in each lung with the total surface area is 40 - 80m²
- Cell types of the epithelial lining
 - type I pneumocytes
 - an extremely thin cytoplasm, and thus provide only a thin barrier to gas exchange, derived from type II pneumocytes
 - connected to each other by tight junctions that limit the fluid movements in and out of the alveoli
 - easily damageable, but cannot divide!
 - type II pneumocytes
 - slightly more numerous than type I cells but cover less of the epithelial lining
 - the source of type I cells and **surfactant**
 - macrophages




Alveolo - capillary barrier




- Alveolar epithelia
 - type I and II cells
- Capillary endothelium
 - non-fenestrated
- Interstitium
 - cells (very few!)
 - fibroblasts
 - contractile cells
 - immune cells (interstitial macrophages, mast cells, ...)
 - ECM
 - elastin and collagen fibrils

Pulmonary surfactant

- Complex mixture of lipids and proteins at the alveolar cell surface (liquid – gas interface) reducing surface tension
 - superficial layer made of phospholipids (dipalmitoyl lecithin)
 - deeper layer (hypophase) made of proteins (SP-A, -B, -C, -D)
- Surfactant maintains lung volume at the end of expiration
- Continually recycles
 - influenced by many hormones incl. glucocorticoids
 - lung maturation in pre-term newborns



Pulmonary surfactant adsorption to the interface and surface film formation : Processes that may contribute to transport of surface active surfactant species to the interface include 1) direct cooperative transfer of surfactant from secreted lamellar body-like particles (LB) touching the interface, 2) unravelling of secreted LB to form intermediate structures such as tubular myelin (TM) or large surfactant layers that have the potential to move and transfer large amounts of material to the interface, and 3) rapid movement of surface active species through a continuous network of surfactant membranes (so-called surface phase) connecting secreting cells with the interface.




©2010 by American Physiological Society

Perez-Gil J, Weaver T E *Physiology* 2010;25:132-141

Pulmonary vasculature and lymphatics


- Lungs are the only organ through which **all the blood (CO)** has to pass!!!
- Lungs have a **dual blood supply**
 - deoxygenated blood from the right ventricle via the pulmonary artery
 - systemic (nutritional) supply throughout the bronchial circulation
 - arises from the descending aorta
 - bronchial arteries supply tissues down to the level of the respiratory bronchiole
 - bronchial veins drain into the pulmonary vein, forming part of the **physiological shunt** observed in normal individuals
- Drainage is provided by the **four main pulmonary veins** (into the left atrium)
- Lymphatics start in the interstitial space between the alveolar cells and the capillary endothelium of the pulmonary arterioles
 - the tracheobronchial lymph nodes arranged in five main groups:
 - paratracheal, superior tracheobronchial, subcarinal, bronchopulmonary and pulmonary



10

Defense mechanisms of the resp. tract


- These can be divided into two kinds of mechanisms:
 - physical
 - humidification
 - particle removal
 - over 90% of particles greater than 10 μm diameter are removed in the nostril or nasopharynx (incl. most pollen grains which are typically >20 microns in diameter)
 - particles between 5 - 10 microns become impacted in the carina
 - particles smaller than 1 micron tend to remain
 - mucus
 - particle expulsion
 - by coughing, sneezing or gagging
 - immunological
 - humoral
 - cellular
- Pulmonary disease often results from a failure of the many defense mechanisms that usually protect the lung in a healthy individual



11

The ciliated epithelium

- Very important defense mechanism
- Each cell contains approx. 200 cilia beating at 1000 beats per minute in organized waves of contraction
- Each cilium consists of nine peripheral pairs and two inner longitudinal fibrils in a cytoplasmic matrix
 - nexin links join the peripheral pairs
 - dynein arms consisting of ATPase protein project towards the adjacent pairs.
- Bending of the cilia results from a sliding movement between adjacent fibrils powered by an ATP-dependent shearing force developed by the dynein arms
 - congenital absence of dynein arms leads to immotile cilia syndrome
- Mucus, which contains macrophages, cell debris, inhaled particles and bacteria, is moved by the cilia towards the larynx at about 1.5 cm/min (the „**mucociliary escalator**“)



© Elsevier Science Ltd

12

Respiratory tract secretions - mucus

- gelatinous substance (~5 mm thick) consisting of acid and neutral polysaccharides
- relatively impermeable to water
 - mucus floats on a liquid or sol layer that is present around the cilia of the epithelial cells
- secreted from **goblet cells** and **mucous glands** as distinct globules that coalesce increasingly in the central airways to form a more or less continuous mucus blanket
- under normal conditions cilia are in contact with the under surface of the gel phase and coordinate their movement to push the mucus blanket upwards
 - it may only take 30-60 minutes for mucus to be cleared from the large bronchi
 - there may be a delay of several days before clearance is achieved from respiratory bronchioles
- reduction in mucociliary transport
 - one of the major long-term effects of **cigarette smoking**
 - contributes to recurrent infection and in the larger airways it prolongs contact with carcinogens
 - air pollutants, local and general anaesthetics
 - bacterial and viral **infections**
 - congenital defects in mucociliary transport (characterized by recurrent infections and eventually with the development of bronchiectasis)
 - the 'immotile cilia' syndrome and cystic fibrosis: an abnormal mucus composition is associated with ciliary dyskinesia

13

Cellular defense mechanisms

- Pulmonary alveolar macrophages
 - derived from precursors in the bone marrow and migrate to the lungs via the bloodstream
 - phagocytose particles, including bacteria, and are removed by the mucociliary escalator, lymphatics and bloodstream
 - dominant cell in the airways at the level of the alveoli
 - comprise 90% of all cells obtained by bronchoalveolar lavage
 - work principally as scavengers and are not particularly good at presenting antigens to the immune system
- Dendritic cells
 - form a network throughout the airways and are thought to be the key antigen-presenting cell in the airway
- Lymphoid tissue
 - the lung contains large numbers of lymphocytes which are scattered throughout the airways. Sensitized lymphocytes contribute to local immunity through differentiation into IgA-secreting plasma cells. IgG and IgE are found in low concentrations in airway secretions from a combination of local and systemic production.
 - In addition to these resident cells, the lung has the usual range of acute inflammatory responses and can mobilize neutrophils promptly in response to injury or infection and play a major part in inflammatory conditions such as asthma.


15

Humoral defense mechanisms

- Non-specific soluble factors
 - characteristic for lungs
 - **α -Antitrypsin (α -antiprotease)**
 - present in lung secretions derived from plasma
 - inhibits chymotrypsin and trypsin and neutralizes proteases and elastase
 - **Surfactant protein A (SPA)**
 - one of four species of surfactant proteins which opsonizes bacteria/particles, enhancing phagocytosis by macrophages
 - generally found on biological barriers
 - **Lysozyme**
 - an enzyme found in granulocytes that has bactericidal properties
 - **Lactoferrin**
 - synthesized from epithelial cells and neutrophil granulocytes and has bactericidal properties.
 - **Interferon (produced by most cells in response to viral infection)**
 - a potent modulator of lymphocyte function. It renders other cells resistant to infection by any other virus.
 - **Complement**
 - present in secretions and is derived by diffusion from plasma
 - in association with antibodies, it plays an important cytotoxic role
 - **Defensins**
 - bactericidal peptides present in the azurophil granules of neutrophils

14


Summary – lung defense



16


Respiration and gas exchange in the lungs

- **ventilation** = mechanical process
 - breathing in narrower meaning
- **diffusion** = chemical process
 - through alveolo-cappillary barrier
- **perfusion** = circulatory process
 - circulation of blood in lungs



Copyright © 2004 Pearson Education, Inc., publishing as Benjamin Cummings.


17



VENTILATION & PULMONARY MECHANICS

18

Mechanika dýchání




Copyright © 2006 by The McGraw-Hill Companies, Inc. All rights reserved.

- tlaky a tlakové gradienty
 - tlak na povrchu těla (P_{bs}), většinou totožný s atmosferickým (P_{ao})
 - tlak v alveolu (P_{alv})
 - „elastický“ tlak vyvíjený parenchymem plic a povrch. napětím (P_e)
 - tlak v pleurální dutině (P_{pl})
 - transpulmonální tlak = tlakový rozdíl mezi alveolem a pleurální dutinou (P_L)
 - $P_L = P_{alv} - P_{pl}$
 - transtorakální tlak = rozdíl mezi alveoly a tělesným povrchem (P_{rs}), určuje zda probíhá inspirium nebo expirium
 - $P_{rs} = P_{alv} - P_{bs}$

19

Ventilation




1. Barometric air pressure (P_B) is equal to alveolar pressure (P_{alv}) and there is no air movement.
2. Increased thoracic volume results in increased alveolar volume and decreased alveolar pressure. Barometric air pressure is greater than alveolar pressure, and air moves into the lungs.

- pressure necessary to distend lungs has to overcome two kinds of resistances
 - **DYNAMIC** = airway resistance (in the convection part of airways)
 - **STATIC** = elastic recoil (in the respiratory part of airways and lung parenchyma)
- energy requirements for respiratory muscles to overcome these two resistances is normally quite low (2-5% of a total O_2 consumption)
 - but increases dramatically when resistance increases (up to 30%) → subjective perception as a **dyspnea**

20


Ventilation (breathing) as a mechanical process



Copyright © 2001 Benjamin Cummings, an imprint of Addison Wesley Longman, Inc.

- Inspiration
 - an active process that results from the descent of the **diaphragm** and movement of the ribs upwards and outwards under the influence of the **intercostal muscles**
 - in resting healthy individuals, contraction of the diaphragm is responsible for most inspiration
 - respiratory muscles are similar to other skeletal muscles but are less prone to fatigue
 - weakness may play a part in respiratory failure resulting from neurological and muscle disorders and possibly with severe chronic airflow limitation
 - inspiration against increased resistance may require the use of the **accessory muscles of ventilation**
 - sternocleidomastoid and scalene muscles
- Expiration
 - follows passively as a result of gradual lessening of contraction of the intercostal muscles, allowing the lungs to collapse under the influence of their own elastic forces (**elastic recoil**)
 - forced expiration is also accomplished with the aid of accessory muscles
 - abdominal wall



21



23



Elastic properties of the lung

- lungs have an inherent elastic property that causes them to tend to collapse generating a negative pressure within the pleural space
 - the strength of this retractive force relates to the volume of the lung; for example, at higher lung volumes the lung is stretched more, and a greater negative intrapleural pressure is generated
 - at the end of a quiet expiration, the retractive force exerted by the lungs is balanced by the tendency of the thoracic wall to spring outwards
 - at this point, respiratory muscles are resting and the volume of the lung is known as the **functional residual capacity (FRC)**




Elastic recoil is determined by two kinds of forces

- **lung compliance** ("distensibility")
 - a measure of the relationship between this retractive force and lung volume
 - defined as the change in lung volume brought about by unit change in transpulmonary (intrapleural) pressure (L/kPa)
- **surface tension** produced by the layer of fluid that lines the alveoli
 - determined by the cohesive (binding together) forces between molecules of the same type
 - on the inner surface of the alveoli is fluid that can resist lung expansion
 - there would be a lot of surface tension because there is an air-water interface in every alveolus
 - if surface tension remained constant, decreasing r during expiration would increase P and smaller alveolus would empty into large one (A)
 - this collapsing tendency is offset by pulmonary surfactant which significantly lowers surface tension (B)





24




25

Abnormalities of elastic properties

- change of lung compliance (\downarrow TLC, \downarrow FRC, \downarrow RV)
 - \uparrow pulmonary **emphysema**, aging (\uparrow TLC, \uparrow FRC, \uparrow RV)
 - \downarrow **interstitial disease** (\downarrow TLC, \downarrow FRC, \downarrow RV), e.g. pulmonary fibrosis or bronchopneumonia
- lack of surfactant (\downarrow TLC, \downarrow FRC, \downarrow RV)
 - infant or adult **respiratory distress syndromes** (IRE or ARDS, resp.), i.e. lung collapse
 - lung **edema** (damages surfactant)
- diseases that affect the movement of the thoracic cage and diaphragm
 - marked obesity
 - diseases of the thoracic spine
 - ankylosing spondylitis and kyphoscoliosis
 - neuropathies
 - e.g. the Guillain-Barré syndrome)
 - injury to the phrenic nerves
 - myasthenia gravis



26




Airflow

- From the trachea to the periphery, the airways become smaller in size (although greater in number)
 - the cross-sectional area available for airflow increases as the total number of airways increases
 - the flow of air is greatest in the trachea and slows progressively towards the periphery (as the velocity of airflow depends on the ratio of flow to cross-sectional area)
 - in the terminal airways, gas flow occurs solely by diffusion
- The **resistance to airflow** is very low (0.1-0.2 kPa/L in a normal tracheobronchial tree), **steadily increasing from the small to the large airways**
- Airway tone is under the control of the autonomic nervous system
 - bronchomotor tone is maintained by **vagal efferent nerves**
 - many **adrenoceptors** on the surface of bronchial muscles respond to **circulating catecholamines**
 - sympathetic nerves do not directly innervate them!

27

Airflow

- Movement of air through the airways results from a **difference between the pressure in the alveoli and the atmospheric pressure**
 - alveolar pressure (P_{ALV}) is equal to the elastic recoil pressure (P_{EL}) of the lung plus the pleural pressure (P_{PL})
 - positive P_{ALV} occurs in expiration and a negative pressure occurs in inspiration
- During quiet breathing the sub-atmospheric pleural pressure throughout the breathing cycle slightly distends the airways
 - during vigorous expiratory efforts (e.g. cough) the central airways are compressed by positive pleural pressures exceeding 10 kPa
 - the airways do not close completely because the driving pressure for expiratory flow (alveolar pressure) is also increased
- When there is no airflow (i.e. during a pause in breathing) the tendency of the lungs to collapse (the positive P_{EL}) is exactly balanced by an equivalent negative P_{PL}




The relationship between maximal flow rates on expiration and inspiration is demonstrated by the maximal flow-volume (MFV) loops

28


Airflow obstruction

- In patients with severe COPD, limitation of expiratory flow occurs even during tidal breathing at rest
- To increase ventilation these patients have to breathe at higher lung volumes and also allow more time for expiration by increasing flow rates during inspiration, where there is relatively less flow limitation
- Thus patients with severe airflow limitation have a prolonged expiratory phase to their respiration



29


Other examples of flow-volume loops



30

Airway resistance

- Poiseuille's law for pressure states that pressure is directly proportional to flow, tube length, and viscosity, and it is inversely proportional to tube radius
- Overcoming increased resistance requires **forced expiration**




Q	Flow rate
P	Pressure
r	Radius
η	Fluid viscosity
l	Length of tubing

$$Q = \frac{\pi Pr^4}{8\eta l}$$

31

Airflow resistance - bronchoconstriction




- theoretical amplifying effect of luminal mucus on airflow resistance in asthma. (a) According to Poiseuille's law, resistance to flow (R) is proportional to the reciprocal of the radius (r) raised to the fourth power. (b) Without luminal mucus, bronchoconstriction to reduce the airway radius by half increases airflow resistance 16-fold. (c) A small increase in mucus thickness (t_M), which reduces the radius of the airway by only one-tenth, has a negligible effect on airflow in the unstricted airway (compare with panel a). (d) With bronchoconstriction, the same amount of luminal mucus markedly amplifies the airflow resistance of this airway


32

Dynamic compression


- In forced expiration, the driving pressure raises both the P_{ALV} and the P_{PL}
 - between the alveolus and the mouth, a point will occur (C) where the airway pressure will equal the intrapleural pressure, and airway compression will occur
 - however, this compression of the airway is temporary, as the transient occlusion of the airway results in an increase in pressure behind it (i.e. upstream) and this raises the intra-airway pressure so that the airways open and flow is restored
 - the airways thus tend to vibrate at this point of 'dynamic compression'




33



Dynamic compression in various situations



- The respiratory system is represented as a piston with a single alveolus and the collapsible part of the airways within the piston
 - C, compression point; PALV, alveolar pressure; PEL, elastic recoil pressure; PPL, pleural pressure.
 - (a) at rest at functional residual capacity
 - (b) forced expiration in normal subjects
 - (c) forced expiration in a patient with COPD



VÝMĚNA PLYNŮ V PLICÍCH & POMĚR VENTILACE-PERFÚZE

35

36


Funkční klasifikace dýchacích cest

	Name	Division	Diameter (mm)	How many?	Cross-sectional area (cm)
Conducting system	Trachea	0	15-22	1	2.5
	Primary bronchi	1	10-15	2	↓
	Smaller bronchi	2	1-10	4	
		3			
		4			
		5			
6-11	1×10^4				
Exchange surface	Bronchioles	12-23	0.5-1	2×10^4	100
	Alveoli	24	0.3	8×10^7	5×10^3
				$3-6 \times 10^8$	$>1 \times 10^6$

37


Functional classification of airways

- **Conducting airways (= anatomical dead space)**
 - nose (mouth)
 - larynx
 - trachea
 - main bronchi & bronchioles
 - gas conduction, warming
- **Acinar airways (= respiratory space)**
 - respiratory bronchioles
 - alv. ducts & sacs
 - alveoli
 - gas exchange
- **The concept of acinus**
 - the functional 3-D unit - part of parenchyma - in which all airways have alveoli attached to their wall and thus participating in gas exchange




38

Koncept acinu



- 3-D struktura následující po terminálním bronchiolu
 - 3 úrovně větvení resp. bronchiolu a násl. cca 8 větvení alveolárních duktů,
 - každý plicní lalůček (anatomický pojem) tak vyplňuje 10 - 30 acinů


Výměna plynů v plicích



- hlavní funkce dýchacího systému – **výměna plynů mezi okolím a krví** – je podřízena časově variabilním nárokům organismu na O_2
 - udržovány v optimálním rozmezí zejm. regulací intenzity ventilace
- nároky jsou určeny spotřebou **ATP** a jeho nahrazováním **mitochondriemi** (ox. fosforylací)
- alveolo-kapilární výměna plynů probíhá z alv. prostoru do krve **prostou difuzí** přes stěnu alveolu, plicní intersticiu a stěnu kapilár
- hnací silou dodávky O_2 (a recipročně CO_2) je postupný pokles jeho parciálního tlaku, tj. **konační gradient** mezi vdechovaným vzduchem, krví a tkáněmi:
 - parciální tlak = tlak, který by plyn měl pokud by byl ve směsi sám

Výměna plynů v plicích


- důvody poklesu PO_2 :
 - kompetice s CO_2 v alveolu (do výše atm. tlaku)
 - alveolární rovnice plynů $PAO_2 = PIO_2 - (PaCO_2/R)$
 - ne 100% difuze přes alveolokapilární membránu
 - fyzilogický pravo-levý zkrat
 - míchání okysličené a neokysličené krve (aa. bronchiales a vv. coronarie)
 - fyzilogicky malá část Hb jako Met-Hb a COHb
 - postupné spotřebování v průběhu acinu



41

Kvantitativně

- (1) vdechovaný **atmosférický** vzduch
 - 21% O_2 , 0.03% CO_2 , 78% N_2 , vodní páry 0.6% a zbytek tvoří další plyny (argon, helium, ..)
 - atmosférický tlak je 760 mmHg (101 kPa)
 - parc. tlak O_2 (PO_2): $0.21 \times 760 = 160$ mmHg
 - analogicky $PCO_2 = 0.3$ mmHg
- (2) **alveolární** vzduch (směs vdechovaného a vydechovaného vzduchu)
 - $PAO_2 = 100$ mmHg (13.3 kPa), $PACO_2 = 40$ mmHg (5.3 kPa)
 - parc. tlak O_2 v alveolu je o něco nižší než v atmosféře kvůli většímu zastoupení CO_2 v alveolu (vydechovaný vzduch)
- (3) **arteriální** krev
 - $PaO_2 = 90$ mmHg (12 kPa), $PaCO_2 = 45$ mmHg
 - difuze kyslíku není 100% a navíc existuje fyziologický zkrat
- (4) **venózní** krev
 - $PvO_2 = 30 - 50$ mmHg




Alveolar pressure = $P_{O_2} + P_{CO_2} + P_{H_2O} + P_{N_2}$

	vzduch (P)	alveolární (PA)	arteriální (Pa)	venózní (Pv)
O_2	21kPa/150mmHg	13.3 kPa/100mmHg	12kPa/90mmHg	5.3kPa/40mmHg
CO_2	0.03kPa/0.3mmHg	5.3kPa/40mmHg	5.3kPa/40mmHg	6.0kPa/45mmHg


42

Transport plynů krví




- krví (ve vazbě na Hb a fyzikálně rozpuštěný) je kyslík dodáván do všech částí těla, kde difunduje do tkání
 - při fyziologickém PaO_2 (90mmHg/12kPa) a fyziologickém hemoglobinu je téměř 100% saturace
 - a do poklesu PaO_2 na 12kPa saturace významně neklesá
 - saturace měřena pulzní oxymetrií
 - rozhodující je množství v mitochondriích
 - pro dostatečnou produkci ATP je nutné pO_2 v tkáních > 0.13 kPa (1mmHg) = kritická tenze kyslíku
 - organismus potřebuje kyslík:
 - cca 250ml/min → 350l/den v klidu
 - při zátěži mnohem více

Význam kyslíku v organismu





- v těle neexistují větší zásoby kyslíku
 - stačí cca na 5min
 - dýchání a dodávka kyslíku tkáním je proto nepřetržitý děj
 - jeho úplné přerušení znamená
 - ohrožení života (<5min)
 - reverzibilní ztráta zraku za cca 7s, bezvědomí za cca 10s
 - klinickou smrt (~5-7min), event. smrt mozku
 - smrt organismu (>10min)
- 85-90% využito v aerobním metabolismu při výrobě ATP na
 - udržení iontových gradientů
 - svalová kontrakce
 - syntézy
- pro zbytek procesů je pokles pO_2 méně kritický
 - hydroxylace steroidů
 - detoxikace (hydroxylace) cizorodých látek v játrech
 - syntéza oxidu dusnatého (→ vazodilatace)
 - degradace hemu hemoxygenázou

Sumárně: plíce jako součást „O₂ dráhy“




Ventilace a perfúze plic

- vztah mezi ventilací a perfúzí plic je variabilní
 - do jisté míry i u zdravých lidí
 - rozdíly mezi apexem a bazí plíce
 - apex: ventilace alveolů s redukovanou perfúzí (tzv. fyziologický mrtvý prostor, $V_A/Q = 3.3$)
 - báze: perfúze alveolů s redukovanou ventilací (fyziologický zkrat, $V_A/Q = 0.7$)
- ventilačně perfúzní (V_A/Q) nepoměr se významně zvyšuje u některých plicních nemocí a zodpovídá za jejich projevy
 - $\uparrow V_A/Q$ poměru (tj. \uparrow **mrtvého prostoru**)
 - např. plicní embolie
 - $\downarrow V_A/Q$ poměru (tj. \uparrow **plicního zkratu**)
 - obstrukční nemoci plic
 - kolaps plíce
- optimalizace $\downarrow V_A/Q$ - **vazokonstrikční reflex**
 - cévy okolo méně ventilované části plíce se kontrahují
 - ale!!! viz důsledky obstr. nemocí



46

Ventilation-perfusion inequality



more detail next time!!!


47

CONTROL OF RESPIRATION & ITS DISORDERS




48

Control of respiration




Copyright © 2001 Benjamin Cummings, an imprint of Addison Wesley Longman, Inc.

- central chemoreceptors in medulla oblongata





- peripheral chemoreceptors in aorta and glomus caroticum (via n. glossopharyngeus and vagus)
 - active when ↓PaO₂ below 10kPa
 - activation supported by hypercapnia
- pulmonary mechanoreceptors

Central chemoreceptors



- sensitive to ↑PaCO₂ (and subsequent formation of H⁺ in CF)
- H⁺ cannot go through hematoencephalic barrier therefore response to other than respiratory acidosis slower
 - increase in [H⁺] due to metabolic acidosis (e.g. diabetic ketoacidosis) will subsequently increase ventilation with a fall in PaCO₂ causing deep (Kussmaul) respiration
- very quick adaptation to acute or intermittent hypercapnia, however, gets adapted to chronic hypercapnia due to ↑HCO₃⁻ in cerebrospinal fluid
 - problem in COPD - in these patients hypoxaemia is the chief stimulus to respiratory drive
 - oxygen treatment may therefore reduce respiratory drive and lead to a further rise in PaCO₂


Peripheral chemoreceptors - oxygen sensors




- Glomus caroticus and aortic bodies - sensitive to change of PaO₂
 - decrease of O₂ in these cells closes K⁺ channels → depolarization → ↑ intracellular Ca²⁺ → excitation → activation of the respiratory centre
- When hypoxemia is not accompanied with hypercapnia, activation of this sensors is when PaO₂ < 7,3 kPa (55 mm Hg)

Respiratory stimuli

- Coordinated respiratory movements result from rhythmical discharges arising in interconnected neurones in the reticular substance of the brainstem (medulla oblongata), known as the **respiratory centre**
 - via the phrenic and intercostal nerves to the respiratory musculature (principal and auxiliary respiratory muscles)




- the pulmonary blood flow of 5 L/min carries 11 mmol/min (250 mL/min) of oxygen from the lungs to the tissues
- ventilation at about 6 L/min carries 9 mmol/min (200 mL/min) of carbon dioxide out of the body
- normal PaO₂ is between 11 and 13 kPa (83 - 98 mmHg)
- normal PaCO₂ is 4.8-6.0 kPa (36-45 mmHg)



53

Respiratory centres




- Respiratory centre is formed by several groups of neurons:
 - The basic automatic rhythm of respiration is due to activity of Dorsal Respiratory Group (DRG) – inspiration neurons – efferent impulses go to diaphragma and inspiration intercostal muscles
 - DRG also obtain afferent stimuli from the peripheral chemoreceptors and several pulmonary receptors
 - Ventral Respiratory Group (VRG) contains both inspiration and expiration neurons
 - inactive during to normal ventilation, increased ventilation leads to their activation

54

Higher respiratory centres


- Medulla
 - quiet inspiration
 - effort inspiration and forced expiration
- Pons - Pneumotaxic and apneustic centres can modulate depth of ventilation and its frequency
 - Apneustic centre:
 - supports inspiration by the activity of inspiration neurons
 - Pneumotaxic centre:
 - antagonises apneustic centre
 - inhibition of inspiration
- Ventilation can be modulate by cortex, limbic system and hypothalamus (emotions and diseases).



55

Dyspnea (breathlessness)


- on physical exertion is normal and not considered a symptom unless the level of exertion is very light, such as when walking slowly
- although breathlessness is a very common symptom, the sensory and neural mechanisms underlying it remain obscure
- the sensation of breathlessness is derived from at least three sources:
 - changes in lung volume
 - sensed by receptors in thoracic wall muscles signalling changes in their length
 - the tension developed by contracting muscles
 - this can be sensed by Golgi tendon organs
 - tension developed in normal muscle can be differentiated from that developed in muscles weakened by fatigue or disease
 - central perception of the breathing effort




common causes of dyspnea

56

Apnea



- suspension of external breathing
- causes
 - voluntarily achieved (free diving)
 - drug-induced (e.g. opiate toxicity)
 - during sleep
 - mechanically induced (e.g. OSA)
 - infants (sudden death)
 - central apnea syndromes
 - periodical breathing
 - Cheyne-Stokes breathing
 - patients with cardiac failure
 - consequence of neurological disease or trauma



Obstructive sleep apnea (OSA)

- Flow of air pauses or decreases during sleep because the airway has become narrowed, blocked, or floppy
 - breathing pauses can last from a few seconds to minutes
 - may occur 30 times or more an hour
 - typically, normal breathing then starts again, sometimes with a loud snort or choking sound
- During apnea deep sleep shifts to light sleep
 - as a result, the quality of sleep is poor, which makes one tired during the day (excessive daytime sleepiness)
- Commonly undiagnosed, typically overweight adults
- Risks – due to hypoxia



- changes in the neurons of the hippocampus and frontal cortex
- hypertension
- coronary artery disease
- type 2 diabetes
- depression
- sleepiness-related accidents





Figure 2-30. Normal, spontaneous breathing (eupnea). The I:E ratio typically is 1:2.




Figure 2-38. Cheyne-Stokes respiration: A gradual increase and decrease in the volume and rate of breathing, followed by 10 to 30 seconds of apnea.




Figure 2-40. Kussmaul's respiration: Increased rate and depth of breathing. This breathing pattern causes the P_{aCO_2} and P_{aO_2} to decrease and $P_{aH_2CO_3}$ and P_{aO_2} to increase.




Figure 2-35. Biot's respiration: Short episodes of rapid, uniformly deep inspirations, followed by 10 to 30 seconds of apnea.

