



Autonomic nervous system




AUTONOMIC NERVOUS SYSTEM

- **autonomic nervous system participates in innervation of the visceral part of human body, it controls autonomic functions, which take place independently of our will**
- **It is influenced by cerebral cortex, limbic system and hypothalamus**
- **It makes innervation of visceral organs and vessels, of smooth muscle, myocardium and glands**
- **There is inserted one another neuron into visceromotoric pathways, the neuron is located in autonomic ganglia**
- **preganglionic and postganglionic fibers**
- **enteric system –separate part**




Types of stimuli

- Nuclei within CNS → **visceromotor fibers** – through anterior roots of spinal cord → autonomic ganglia along the spine – to the organs of abdomen, thorax, pelvis
- Free nerve endings in the wall of organs → pressure, thrust, pain- **viscerosensory** → autonomic ggl.- to posterior roots of spinal cord - ggl. spinale or ggl. VII., IX., X. and into visceromotor nuclei




- **Autonomic tracts don't go from CNS directly – they switch over in ganglia outside CNS**
- They are formed at least by two neurons, which switch over in so-called **autonomic ganglion**
- **Preganglionic neuron** : myelinated axon that goes from CNS to autonomic ganglion
- **Postganglionic neuron**: unmyelinated axon that goes from autonomic ganglion as a proper autonomic nerve



- autonomic (visceromotor) nerve fibers are of two types
sympathetic *pars sympathica*
parasympathetic *pars parasympathica*
- Glands and smooth muscle of almost each visceral organ are innervated by both sympathetic and parasympathetic
- One system is usually activating and the other inhibiting
- Exception are **smooth muscle of the skin and skin glands**, which are innervated only by sympathetic

Main functions

- Contraction and relaxation of smooth muscle
- Function of all exocrine and some endocrine glands
- Heart rhythm
- Some metabolic processes



Division of autonomic nerve system

- sympathetic – *fight or flight*
- parasympathetic – *rest or digest*
- enteric system



Sympathetic
fight or flight



Parasympathetic
rest or digest



Pars sympathica: nuclei in CNS and in the spinal cord (C8 – L3)

Pars parasympathica: nuclei in CNS (which belong to the cranial nerves), spinal cord (S2 – S4)

craniosacral system (parasympathetic)

thoracolumbar system (sympathetic)

cranio-sacral system (parasympathetic)

Sympathetic and parasympathetic system differ in the arrangement of ganglia:

Sympathetic ganglia:


- are far from target organs (at spine) – paravertebral ganglia – truncus sympathicus dexter et sinister

Parasympathetic ganglia:

- closer to organs (ganglion ciliare, pterygopalatinum, oticum, submandibulare + scattered within organ walls)

Mediators of sympathetic and parasympathetic system:

- preganglionic – the same (from CNS) – acetylcholine
- postganglionic - sympathetic – noradrenalin
- postganglionic - parasympathetic - acetylcholine




SYMPATHETIC


„thoracolumbar system“

Functions


- It controls the **catabolic** functions, activates functions of the visceral organs
- **it accelerates** the heart activity and breathing
- It causes **contraction** of smooth muscle of vessels within the skin and visceral organs and thereby increases blood pressure
- **It increases** level of sugar in blood
- **It expands** pupils (mydriasis)
- It conversely slows digestion
- It induces a **state of wakefulness** and it is used in stress reactions

- Central part: nucl. hypothalamici medii, nucl. intermediolateralis C8-L3- known as **thoracolumbar system**
- it arises from spinal nerve as ***ramus communicans albus*** – it ends in sympathetic ganglion next to the backbone – **preganglionic** section- into **paravertebral ggl.**
- individual paravertebral ganglions form ***truncus sympathicus*** – peripheral section
- from ganglions arise proper sympathetic nerves- **postganglionic** section
- Sympathetic nerves enter (directly or via plexuses inside wall of big arteries) the innervated organs





— Sympathetic fibers
— Parasympathetic fibers




Sympathetic fibers
 Presynaptic ——— (solid red)
 Postsynaptic - - - - - (dashed red)

Parasympathetic fibers
 Presynaptic ——— (solid blue)
 Postsynaptic - - - - - (dashed blue)

Antidromic conduction ——— (solid green)

Truncus sympathicus



adix anterior (motoria)
anglion spinale

.....
orgány

↑
) | |

Cervical part


Ganglion cervicale superius



Ganglion cervicale medium

Ganglion cervicothoracicum /
stellatum

- It forms periarterial
plexuses around a. carotis
ext. et int. – intake of
sympathicus to neck and
head

- *nn. cardiaci* – innervation
of the heart





Ganglia thoracica (thoracic part)

- 10 pairs of ganglia
- nn. splanchnici – for smooth muscle of GIT and its vessels
- rr. communicantes grisei - to intercostal nerves
- Branches to heart, lungs, esophagus


Ganglia lumbalia (lumbar, abdominal part)



- 4-5 pairs of ganglia
- rr. communicantes grisei
- nn. splanchnici lumbales
- rr. vasculares

Ganglia sacralia (pelvic part)

- 4 pairs of ganglia
- rr. communicantes grisei – for pelvic organs
- Periarterial plexuses

13





PARASYMPATHETIC


Functions


- **pars cranialis:** most important is parasympathetic part of *nervus vagus* – it innervates the digestive tract till the border between colon transversum and colon descendens in the abdominal cavity
- **pars sacralis:** it innervates the digestive tract from the border between colon transversum and colon descendens till rectum and visceral organs located in the pelvis (urinary bladder, genital organs except gonads)

It controls **anabolic reactions** – preservation of energy, it induces **inhibition of organism:**

- **It slows** heart activity and breathing
- **It decreases** blood pressure
- **It narrows** pupils (miosis)
- **It accelerates digestion**, sweating and salivation
- It is used especially **at rest (sleep)** and during digestion

- **pars cranialis- III.,VII.,IX.,X.** (cranial parasympathetic)
- **pars sacralis S2-S4** (sacral parasympathetic) - **craniosacral system** - ganglia are located close to the innervated organs, preganglionic section is therefore long and postganglionic section is short
- mediator is **acetylcholin** in whole section - cholinergic system





Cranial parasympathetic

- **Nuclei of cranial nerves:**

ncl. oculomotorius accessorius to ganglion ciliare
(m. sphincter pupillae, m. ciliaris)

ncl. salivatorius superior (VII.) to ganglion pterygopalatinum and submandibulare (lacrimal gland, mucosa of nasal cavity, palate, tongue, gl. sublingualis and submandibularis)

ncl. salivatorius inferior (IX.) to ganglion oticum (glandula parotis and small salivatory glands of cheek)

ncl. dorsalis n. X (together with n. vagus to organs)

Sacral parasympathetic

Caudal elongation of ncl. intermediolateralis

S2-5 (pars sacralis, pelvica) – to pelvic organs
(terminal part of large intestine, rectum, urinary bladder, inner genital organs except sexualglands)

nn. splanchnici pelvici

Ganglion ciliare

- Here end preganglionic fibers of *n. oculomotorius*
- **parasympathetic** (*m. sphincter pupillae, m. ciliaris*)
sympathetic (*m. dilatator pupillae*)

Ganglion pterygopalatinum

- Here end preganglionic fibers of *n. facialis*
- *Mucosa of posterior part of nasal cavity, upper teeth, mucosa of hard palate, lacrimal gland*

Ganglion submandibulare

- Here end preganglionic fibers of *n. facialis*
- *gl. sublingualis, gl. submandibularis, salivary glands of tongue and bottom of oral cavity*


Ganglion oticum

- Here end preganglionic fibers of *n. glossopharyngeus*
- *skin, mucosa, teeth and gingiva of lower jaw, gl. parotidea*

Preganglionic fibers of ***n. vagus*** end in prevertebral ganglia of thoracic and abdominal cavity

Preganglionic fibers of ***sacral parasympathetic*** are switched over in pelvic plexuses

Ggl. ciliare



nn.ciliares breves

m.dilator pupillae

mydriasa

m.sphincter pupillae, m. ciliaris

Miosa, akomodace

Ggl. pterygopalatinum



rr.nasales posteriores sup. et inf.

n.palatinus major

nn.palatini minores


Žlázký dutiny nosní a patra

r.communicans cum n. lacrimali

Glandula lacrimalis

VII. -ncl. salivatorius sup

Ggl. submandibulare



rr. glandulares pro


**gl. submandibularis a
sublingualis**

Spojky do n. lingualis

Drobné slinné žl. jazyka

VII. Ncl. salivatorius sup.

Ggl. oticum



rr. comm. cum n. auriculotemporalis

Glandula parotis


r. comm. cum n. buccalis

Slinné žl. tváře


IX. Ncl salivatorius inf


Motorická vlákna pro m. tensor veli palatini a m. tensor tympani **VII**

m. pterygoideus medialis **V**




BRANCHES:





© www.kenhub.com






Mixed autonomic (prevertebral) plexuses

1) Mixed plexuses of thorax

- plexus cardiacus
- plexus aorticus thoracicus
- plexus pulmonalis
- plexus oesophageus


2) Mixed plexuses of abdomen and pelvis

- plexus aorticus abdominalis: plexus coeliacus, plexus renalis et suprarenalis, plexus testicularis (ovaricus), plexus uretericus, plexuses around vessels
- plexus hypogastricus superior: plexus rectales, plexus vesicales, plexus prostaticus, plexus deferentialis, plexus uterovaginalis




Enteric system

- In the wall of digestive tract
- **plexus submucosus**
- **plexus myentericus**
- Separate and independent of connection with sympathetic and parasympathetic
- It works also after interruption of connections with **ANS**
- It controls tension and mobility of digestive tract, it regulates secretion of all glands and blood flow
- innervation and regulation of function of gall bladder and pancreas



The neural tracts tractus nervosi



THE SENSORY TRACTS

- receptor→CNS
- A) specific:specific information
- B) nonspecific: through interneurons, general information, preparation of CNS for income of specific information
- 1th neuron: pseudounipolar cell of spinal ganglion (ganglion of cranial nerves)
→(cerebellum)→thalamus→cortex
- **SOMATOSENSORY TRACTS:** protopathic sensibility
epicritic sensibility
proprioception
- **VISCEROSENSORY TRACTS**

- **Protopathic sensibility**: tactile information (warmth, cold, pressure, pain, rough skin sensibility)

1) Limbs and trunk: *tractus spino-thalamo-corticalis*

1st Pseudounipolar neuron of spinal ganglion → 2nd nucleus proprius → 3rd thalamus → cortex (gyrus postcentralis, area 1, 2, 3)

2) Head area: *tractus trigemino-thalamo-corticalis*

1st Pseudounipolar neurons of sensory ganglia of CN (V., VII., IX., X.) → 2nd nucleus tractus spinalis (V.) → 3rd thalamus → cortex

- **Epicritic sensibility**: discriminatory sensation (tactile resolution of shape of object etc.)

1) Limbs and trunk: *tractus spino-bulbo-thalamo-corticalis*

1st Pseudounipolar neuron of spinal ganglion → fasciculus gracilis, fasciculus cuneatus → 2nd nucleus gracilis, cuneatus medialis → 3rd thalamus → cortex (gyrus postcentralis, area 1, 2, 3)

2) Head area: *tractus trigemino-thalamo-corticalis*

1st Pseudounipolar neurons of sensory ganglia of CN (V., VII., IX., X.) → 2nd nucleus principalis (V.) → 3rd thalamus → cortex

- **Proprioception**: from the locomotor system to the cerebellum

1) LL and trunk: 1st Pseudounipolar neuron of spinal ganglion → 2nd nucleus thoracicus → 3rd cerebellum → 4th thalamus → cortex

2) UL: 1st Pseudounipolar neuron of spinal ganglion → fasciculus cuneatus → 2nd nucleus cuneatus lateralis → 3rd cerebellum → 4th thalamus → cortex

3) Head area: *tractus trigemino-thalamo-corticalis*

1st Pseudounipolar neurons of nucleus mesencephalicus nervi V. → 2nd cerebellum → 3rd thalamus → cortex

- **VISCEROSENSORY TRACTS**


1st Pseudounipolar neuron of spinal ganglion → 2nd nucleus intermediomedialis

→ nucleus intermediolateralis

→ FR → thalamus → cortex


MOTOR TRACTS

- Set of all neural tracts, which are connected into the regulation of movement. To them belong **pyramidal** and **extrapyramidal tracts**.



PYRAMIDAL TRACTS (direct)

- projection **direct** motor tracts of **voluntary movement**
- They interconnect motor cortex of hemisphere with motoneurons of anterior spinal horns and with motoneurons of nuclei of cranial nerves
- It is only one-neuron way
- They start in primary motor cortex, to them belong **tractus cortico-spinalis (tract of voluntary movement of trunk and limbs)** and **tractus cortico-nuclearis (tract of voluntary movement of striated muscles of the head)**.




(a) Corticospinal pathway

EXTRAPYRAMIDAL TRACTS (indirect)

- Control of involuntary movement
- **Projection extrapyramidal tracts** (connect motor cortex of hemisphere with motoneurons of anterior spinal horns; they are switched over in motor nuclei of brainstem)
- **Connections of (motor) basal ganglia** (BG are interconnected with each other and with other motor structures of brain (e.g. motor cortex and motor parts of thalamus))
- **Tracts of cerebellum**
- They further interconnect **motor nuclei of thalamus, RF and e.g. nucleus ruber, substantia nigra etc.**

- Association tracts:
 - The same hemisphere: *fibrae arcuatae*, *fasciculus longitudinalis superior*, *et inferior*, *fasciculus uncinatus*, *fasciculus arcuatus*
- Commissural tracts:
 - Right and left side of CNS: *commissura anterior et posterior*, *commissura fornicis*, *corpus callosum*



Obrázky:

- **Atlas der Anatomie des Menschen/Sobotta. Putz,R., und Pabst,R. 20. Auflage. München:Urban & Schwarzenberg, 1993**
- **Netter: Interactive Atlas of Human Anatomy.**
- **Naňka, Elišková: Přehled anatomie. Galén, Praha 2009.**
- **Čihák: Anatomie I, II, III.**
- **Drake et al: Gray's Anatomy for Students. 2010**