

# Regulation of Blood Flow

Assoc. Prof. MUDr. Markéta Bébarová, Ph.D.

*Dept. of Physiology, Faculty of Medicine, Masaryk University*



**This presentation includes only the most important terms and facts. Its content by itself is not a sufficient source of information required to pass the Physiology exam.**

# Definition of Blood Flow

mathematical formulation – analogy with the electric current

Ohm's law

$$I = U / R$$



$$Q = \Delta P / R$$

Q

blood flow



difference of pressure at the beginning and at the end of a vessel

R

resistance of the vessel  
(peripheral resistance)

# Definition of Blood Flow

$$Q = \Delta P / R$$


$$R = 8\eta l / \pi r^4$$

$r$  radius of the vessel  
 $\eta$  viscosity of the blood  
 $l$  length of the vessel

This formula applies to the steady state flow. Viscosity of the blood is not constant, it varies with the velocity of blood flow. The blood flow is laminar in the middle (*plasma skimming*). The elasticity of vessels.

Hagen - Poiseuille formula

$$Q = \Delta P \cdot \pi r^4 / 8\eta l$$




# Methods for Measuring Blood Flow


- A. with a cannula inserted into a vessel
- B. without direct contact with the blood flow
  1. Electrical Induction Principle
  2. Doppler Effect
  3. Plethysmography
  4. Fick Principle

# Methods for Measuring Blood Flow

## 1. Electrical Induction Principle (Faraday, 1791-1867)



- ❖ the electromagnetic flowmeter
- ❖ an electromotive force is generated in the blood (as a conductor) when it moves through a magnetic field




- ❖ this electromotive force (measured with an electrode placed on the vessel surface) is proportional to the velocity of blood flow
- ❖ can detect changes in the velocity  $< 0.01$  s recording of both steady blood flow and its pulsatile changes

# Methods for Measuring Blood Flow

## 2. Doppler Effect (Christian Doppler, Prague 1842)




- ❖ the ultrasonic Doppler flowmeter; most common
  - ultrasonic waves of a known wave length (frequency) are sent into a vessel diagonally along the blood stream from a subtle piezoelectric crystal
  - waves reflect from the red and white blood cells ████ change ( $\uparrow$ ) of the wave length ( $\downarrow$  frequency)
  - reflected waves are picked up by a sensor
- ❖ change of the wave length (frequency) is proportional to the velocity of blood flow
- ❖ both steady blood flow and its pulsatile changes can be measured



# Methods for Measuring Blood Flow

## 3. Plethysmography

- ❖ usually as the venous occlusion plethysmography
  - ❖ can be used on limbs
- 
- venous drainage of the limb is stopped (e.g. with an arm cuff)
  - increasing volume of the limb (expelling water from closed chamber, measured as a change of its volume) is linearly proportional to the arterial inflow of blood






# Methods for Measuring Blood Flow

## 3. Plethysmography

- ❖ usually as the venous occlusion plethysmography
- ❖ can be used on limbs

- venous drainage of the limb is stopped (e.g. with an a cuff)
- increasing volume of the limb (expelling water from closed chamber, measured as a change of its volume) is lineary proportional to the arterial inflow of blood



[http://schueler.ws/?page\\_id=21](http://schueler.ws/?page_id=21)

# Methods for Measuring Blood Flow

## 4. Fick Principle - Direct Fick Method

$$Q = \frac{A / \text{time}}{AV \text{ diff}}$$

- **blood flowing from the right heart to the lungs** – about 150 ml O<sub>2</sub> / 1 l (a sample of the mixed venous blood bled from the pulmonary artery with a catheter inserted to the brachial vene)
- **blood flowing from the lungs to the left heart** – about 200 ml O<sub>2</sub> / 1 l (a sample of the arterial blood from any artery, arterial O<sub>2</sub> content is uniform)




The blood catches 50 ml O<sub>2</sub> / 1 l during passage through the lungs.

- **The total O<sub>2</sub> consumption is 250 ml / 1 min.**  
(O<sub>2</sub> decay in the expired air compared to the inspired air, oximeter)



$$CO = \frac{250 \text{ ml O}_2 / \text{min}}{50 \text{ ml O}_2 / \text{l}} = 5 \text{ l} / \text{min}$$



# Methods for Measuring Blood Flow


$\dot{Q}$  = Cardiac output

$$\dot{Q} = \frac{\text{O}_2 \text{ consumption}}{A-V}$$

$$\dot{Q} = \frac{250 \text{ mL O}_2/\text{min}}{0.05 \text{ mL O}_2/\text{mL}}$$

$$\dot{Q} = 5,000 \text{ mL/min}$$

O<sub>2</sub> consumption =  
250 mL/min



# Methods for Measuring Blood Flow

## 4. Fick Principle – Method of Indicatory Gas

- ❖ to determine the instantaneous blood flow through a specific tissue
- ❖ for example the cerebral or coronary blood flow using inhaled nitrous oxide  $N_2O$  – **Kety method**


$$\text{cerebral blood flow} = \frac{\text{N}_2\text{O concentration in the venous blood} \downarrow \text{N}_2\text{O removed from blood by brain / time}}{\text{averaged arteriovenous difference of N}_2\text{O}}$$

# Methods for Measuring Blood Flow

## 4. Fick Principle - Indicator Dilution Technique

- known amount of an indicator (dye or radioactive isotope) is injected into a peripheral (an arm) vein ( $A$ , [mg])
- concentration of the indicator in serial samples of the arterial blood is determined
- estimation of the averaged concentration of the indicator in the arterial blood after a single circulation ( $C$ , [mg/ml])


$$CO = \frac{A}{C (t_2 - t_1)} \quad \begin{matrix} \text{[mg]} \\ \text{[mg.ml}^{-1}\text{.s]} \end{matrix}$$



# Methods for Measuring Blood Flow

## 4. Fick Principle - Indicator Dilution Technique

- known amount of an indicator (dye or radioactive isotope) is injected into the proximal (PA) infusion line
- concentration of indicator in serial samples of blood is determined
- estimation of blood flow based on concentration of indicator in arterial blood and venous blood circulation



### ❖ thermodilution

a cold saline (indicator) is injected into the right atrium through a double lumen catheter; the change of blood temperature (inversely proportional to the blood flow) is recorded in the pulmonary artery using a thermistor in the other side of the catheter

# Regulation of Blood Flow

$$Q = \frac{\Delta P \cdot \pi r^4}{8 \eta l}$$

**Resting Tone** (intermediary vascular muscle tone at rest)

- ❖ due to tonic activity of vasoconstrictive sympathetic fibres
- ❖ a role might play also: myogenic response of vessels to the blood pressure (later), high concentration of O<sub>2</sub> in the arterial blood, Ca<sup>2+</sup>

**Basal Tone**

- ❖ in response to denervation; due to spontaneous depolarizations of the vascular smooth muscles


Regulation

Local

Systemic



## B. Vasoconstriction and vasodilatation





# Regulation of Blood Flow - Local

## A. Acute

*seconds to minutes, but incomplete (about  $\frac{3}{4}$  of the desired effect)*

1. **Metabolic Autoregulation**
2. Myogenic Autoregulation
3. Regulation Mediated by Local Substances

## B. Chronic

*hours, days to weeks , even months*

# Regulation of Blood Flow - Local

## Metabolic Autoregulation

insufficient blood flow  $\begin{cases} \nearrow \uparrow \text{ metabolic demands of a tissue} \\ \searrow \downarrow \text{ or stopped blood supply} \end{cases}$

→  $\uparrow$  concentration of metabolites ( $\text{CO}_2$ , lactic acid, adenosine,  $\text{K}^+$ , phosphate),  $\downarrow$  pH,  $\uparrow$  osmolarity in the interstitium,  $\uparrow$  tissue temperature (the metabolic heat);  $\downarrow$   $\text{pO}_2$  (the second theory based on the lack of  $\text{O}_2$  and nutrients)

→ **vasodilatation**

**Preferred to the systemic regulation in case of hypoxia (to preserve the adequate tissue perfusion).**

*It plays the key role in e.g. brain, heart and skeletal muscles.*

# Regulation of Blood Flow - Local

## Metabolic Autoregulation


### active hyperemia

(increase of the blood flow induced by an increased metabolic activity of the tissue)

### reactive hyperemia

(transient increase of the blood flow exceeding its common level after release of an occlusion; it gradually returns to the control level)

(15-, 30- and 60-s occlusions of the femoral artery in a dog)



# Regulation of Blood Flow - Local

## A. Acute

*seconds to minutes, but incomplete (about  $\frac{3}{4}$  of the desired effect)*

1. Metabolic Autoregulation
2. Myogenic Autoregulation
3. Regulation Mediated by Local Substances

## B. Chronic

*hours, days to weeks , even months*

# Regulation of Blood Flow - Local

## Myogenic Autoregulation (Bayliss effect)

↑ blood pressure

→ ↑ blood flow and ↑ tension in the vascular wall

$$Q = \Delta P / R$$

Law of Laplace

$$T = P \cdot r$$


→ mechanical stimulation, depolarization and subsequent contraction of the smooth muscle cells in the vascular wall → vasoconstriction

→ return of the blood flow back on the original level

*It plays an important role in the brain and kidneys.*

# Regulation of Blood Flow - Local

## Myogenic Autoregulation



# Regulation of Blood Flow - Local

## A. Acute

*seconds to minutes, but incomplete (about  $\frac{3}{4}$  of the desired effect)*

1. Metabolic Autoregulation
2. Myogenic Autoregulation
3. Regulation Mediated by Local Substances

## B. Chronic

*hours, days to weeks , even months*



# Regulation of Blood Flow - Local


## Regulation Mediated by Local Substances

*endothelial-derived relaxing factor (EDRF) – NO*  
(half-life in the blood only 6 s) **vasodilatation**

- ❖ important in the **intermediate and larger arteries** back upstream where the metabolic tissue changes causing dilatation of the microvessels cannot directly reach
- ❖ **synthesized in the endothelial cells** of arteriols and small arteries **due to the shear stress** induced by the flowing blood (deforms the endothelial cells in the direction of flow)
- ❖ its **synthesis stimulated by the products of thrombocyte aggregation** (to keep vessels with intact endothelium permeable) and also by many **primary vasoconstrictive substances**




## B. Vasoconstriction and vasodilatation



# Regulation of Blood Flow - Local

## Regulation Mediated by Local Substances

*endothelial-derived relaxing factor (EDRF) – NO*



# Regulation of Blood Flow - Local

## Regulation Mediated by Local Substances

### prostacyclin

- ❖ synthesized in the endothelial cells from the arachidonic acid
- ❖ inhibition of thrombocyte aggregation and **vasodilation**

### thromboxane $A_2$

- ❖ synthesized from the arachidonic acid by thrombocytes
- ❖ support of thrombocyte aggregation and **vasoconstriction**

*A balance between them is crucial for formation of the localized clot and preservation of the blood flow. (aspirin)*

# Regulation of Blood Flow - Local

## Regulation Mediated by Local Substances

### endothelins

- ❖ several similar polypeptides synthesized by the endothelial cells (ET-1, ET-2, ET-3 )
- ❖ 2 endothelin **receptors**:
  - ET<sub>A</sub> – specific for ET-1, in many tissue vessels, → vasoconstriction
  - ET<sub>B</sub> – ET-1 to ET-3, unknown function (maybe vasodilatation – *through increased synthesis of NO* - and developmental effects)
- ❖ **ET-1 – one of the most potent vasoconstrictive substances**
- ❖ the exact physiological role not known
- ❖ released from the endothelial cells in the damaged tissue  
vasoconstriction restricts bleeding
- ❖ play a role in closing *ductus arteriosus* at birth

# Regulation of Blood Flow - Local

## Serotonin (5-OH tryptamine)

### ❖ vasoconstrictive effect

- in a damaged tissue
- direct local effect
- released from thrombocytes

### ❖ vasodilatory effect

- in an undamaged tissue
- through increased activity of NO synthase

# Regulation of Blood Flow - Local

## Other, specific mechanisms

- ❖ local vasoconstriction of damaged arteries and arteriols  
(due to release of serotonin and thromboxane A<sub>2</sub> from thrombocytes and endothelin-1 from the endothelial cells)
- ❖ vasoconstriction (vasodilatation) induced by a decrease (increase) of the tissue temperature
- ❖ specialized tissues (kidneys, brain, etc.)

# Regulation of Blood Flow - Local

## A. Acute

*seconds to minutes, but incomplete (about  $\frac{3}{4}$  of the desired effect)*


1. Metabolic Autoregulation
2. Myogenic Autoregulation
3. Regulation Mediated by Local Substances

## B. Chronic

*hours, days to weeks , even months*

# Regulation of Blood Flow - Local

## Chronic regulation



*It is especially important in case of the long-term change of metabolic demands of a tissue - to provide sufficient blood flow without circulation overload.*



# Regulation of Blood Flow - Local


## Chronic regulation

- ❖ mediated by changes of the tissue vascularity
- ❖ the key role – lack of O<sub>2</sub> (higher altitude, retrolental fibroplasia in premature newborns after the curative stay in the oxygen tent) and also nutrients
- ❖ identified number of factors increasing grow of new vessels - angiogenic or vascular growth factors - small peptides, best characterized: vascular endothelial growth factor (VEGF), fibroblast growth factor, and angiogenin
- ❖ proceeds fast (within days) in the young individuals and in newly formed tissue (new scar, tumor tissue) vs. within even months in the elderly and differentiated tissues


# Regulation of Blood Flow - Local

## Chronic regulation

unstimulated muscle



regularly stimulated muscle



# Regulation of Blood Flow

Local

**Systemic**

A. Neural

**B. Humoral**

# Regulation of Blood Flow - Systemic

## Humoral regulation

### Vasoconstrictive substances

#### ❖ norepinephrine

Generalized vasoconstriction ( $\alpha_1$ -rec.)  
( $\uparrow$  BP  $\rightarrow$  reflex bradycardia,  $\downarrow$  CO)

#### ❖ epinephrine (high levels)


Vasodilatation in the skeletal muscles, liver and coronary arteries ( $\beta_2$ -rec.)  
Vasoconstriction in other tissues

#### ❖ angiotensin II

$\downarrow$  BP  $\rightarrow$  secretion of renin  $\rightarrow$  formation of angiotensin II  
Generalized vasoconstriction (+  $\uparrow$  water intake and  $\uparrow$  aldosterone)

#### ❖ vasopressin (antidiuretic hormone)

Generalized vasoconstriction (+  $\uparrow$  reabsorption of water in the kidneys)



# Regulation of Blood Flow - Systemic

## Humoral regulation

### Vasodilatory substances

#### ❖ atrial natriuretic peptide (ANP)

■ reactivity of the vascular smooth muscles on vasoconstrictive stimulation (+ ↑ natriuresis – relaxation of the mesangial cells and, thus, ↑ glomerular filtration rate, + inhibition of vasopressin secretion, + ↓ aldosterone)

#### ❖ VIP (vasoactive intestinal peptide)

■ vasodilatation (+ many other effects in GIT, namely relaxation of the intestinal smooth muscles including sphincters)

#### ❖ histamine

- released in tissues (from the mast cells), or from basophiles in the blood, during tissue damage or inflammation (also allergic)

→ vasodilatation of arteriols + ↑ permeability of capillaries  
(edemas; anaphylactic shock)

← through EDRF (vasoconstrictor by itself)

# Regulation of Blood Flow - Systemic

## Humoral regulation

### Vasodilatory substances

#### ❖ kinins - bradykinin and lysylbradykinin (kallidin)


- small polypeptides, half-life - several minutes

→ vasodilatation of arteriols + ↑ permeability of capillaries  
(similar to histamine)

regulation of the blood flow and leak of fluids from capillaries in the inflamed tissue

+

regulation of the blood flow in the skin, salivary and GIT glands in common conditions



# Regulation of Blood Flow - Systemic

## Humoral regulation

### Other factors

#### ❖ ions

vasoconstriction:  $\uparrow \text{Ca}^{2+}$ , slightly  $\downarrow \text{H}^+$

vasodilatation:  $\uparrow \text{K}^+$ ,  $\uparrow \text{Mg}^{2+}$ ;  $\uparrow \text{H}^+$ , notably  $\downarrow \text{H}^+$   
acetate, citrate (anions) – only mild effect