

SYMPATHOTROPIC DRUGS

SYMPATHOLYTICS

Copyright notice

The presentation is copyrighted work created by employees of Masaryk university.

Students are allowed to make copies for learning purposes only.

Any unauthorised reproduction or distribution of the presentation or individual slides is against the law.


Sympatholytics (direct and indirect)

Indications:

- hypertension (mild and moderate)
- antimigraine drugs
- disorders of peripheral vascularity
- benign prostatic hyperplasia
- urinary obstruction – postoperative atonia
- pheochromocytoma

Direct sympatholytics

α



Direct sympatholytics α non-selective

- **ergot alkaloids** (ergotamine, ergometrine, ergotoxine, methylergometrine, dihydroergotamine, dihydroergotoxine, dihydroergocristine)

Direct sympatholytics α non-selective

Ergot alkaloids and their derivatives → reversible α -lytics

- in *Secale cornutum*, product of *Claviceps purpurea*, fungus that infects cereal crops
- derivatives of lysergic acid
- effects:
 - CNS (hallucinations, ↓ prolactin secretion)
 - smooth muscle of blood vessel (effects mimetic or **lytic**)
 - uterine muscle → contractions

Direct sympatholytics

α non-selective

Ergot alkaloids

- **ergotamine, ergometrine**
 - partial α -agonistic effects
 - uterotonic effect, amplified by methylation of derivatives (methylergometrine)
- **ergotoxine**
 - mixture of alkaloids, mainly **ergocristine, ergocriptine** and **ergocornine** - especially α -lytic effects
- **α -lytic effects are increased in dihydro-derivatives**
(*dihydroergotamine, dihydroergotoxine, dihydroergocristine*)

!

Direct sympatholytics

α non-selective

Ergot alkaloids

- **methylergometrine**

uterotonic effect

- therapy and prevention of uterine bleeding after childbirth (in hypotony and atony of myometrium)

Direct sympatholytics

α_1 selective

α_1 sympatholytics

Overview of drugs, use

- terazosin, doxazosin, alfuzosin, tamsulosin...
- prazosin (in Czech Rep. non registered)
- Use:
 - hypertension (relaxation of arterial and venous smooth muscle)
 - benign prostatic hyperplasia
 - urinary obstruction

Direct sympatholytics with combined effect

- **urapidil**
- combined central and peripheral action, blocks α_1 receptors, in CNS blocks H_1 receptors, activates $5-HT_{1A}$ receptors
- **Use:**
 - hypertension (hypertension crisis, severe, respectively, very severe forms of hypertension and hypertension resistant to standard therapy)

Direct sympatholytics

α_2 selective

α_2 sympatholytics

- **yohimbine** (in Czech Rep. non registered)
- vasodilation in the pelvic area, afrodisiac effect
- it is contained in some dietary supplements

Direct sympatholytics β -blockers, β -sympatholytics

- **competitive antagonists** (intrinsic activity = 0) or **partial agonists**
(ISA - *intrinsic sympathomimetic activity*) = dualists
- **nonselective** or **cardioselective** (selectively block β_1 receptors)
- sufficient solubility in fats \rightarrow penetration across HEB

Direct sympatholytics β -blockers, β -sympatholytics

Organ effects

cardiovascular system: negatively chronotropic and inotropic effect →

↓ **BP** and **HR**

- inhibition of vasodilation by β_2 -receptor blockade → **peripheral vascular resistance increase**

- renine secretion reduction

bronchi: bronchoconstriction

eye: intraocular pressure decrease

metabolic effects: glycogenolysis reduction, lipolysis inhibition

Direct sympatholytics β -blockers, β -sympatholytics

NONSELECTIVE ($\beta_1 + \beta_2$) propranolol, metipranolol

(CARDIO)SELECTIVE (β_1) atenolol, metoprolol

NONSELECTIVE ($\beta_1 + \beta_2$) WITH ISA pindolol,
bopindolol (in
Czech Rep. non
registered)

(CARDIO)SELECTIVE (β_1) WITH ISA acebutolol

WITH COMBINED EFFECTS $\alpha + \beta$ labetalol
carvedilol

Direct sympatholytics β -blockers, β -sympatholytics

Use, indications:

- hypertension
- Ischemic heart disease, non-stabil *angina pectoris*, status after acute myocardial infarction
- arrhythmia
- glaucoma
- hyperthyreosis
- anxiety (moderate effect)

Direct sympatholytics β -blockers, β -sympatholytics

Side effects:

- *asthma bronchiale*, dyspnoea
- heart insufficiency
- bradycardia, blockade of heart impuls conduction
- masking of hypoglycemia symptoms
- disorders of peripheral blood circulation
- sleep disorders, depression (lipophilic drugs)
- rash, fever and other allergic symptoms (rarely)
- abrupt discontinuation of therapy – „**rebound phenomena**“

Nondirect sympatholytics

decreases catecholamine concentration in the synaptic cleft by:

- inhibition of NT synthesis
- inhibition of NT storage
- inhibic  of NT release
- false precursors