(XI.) Digital model of aortic function (XVI.) Blood flow in veins


Definitions of key words and symbols

Stroke volume (SV) – volume of blood ejected from the left ventricle to the aorta during one contraction


Heart rate (HR) – number of heart contractions per1 minute

Peripheral vascular resistance (R) – resistance of small arteries (mainly arterioles and capillaries)

Compliance of aorta (C_a) – ability of aorta to change its volume according to changes of blood pressure




Arterial blood pressure in case of changing circulatory parameters and cardiac output



Model of aortic function






Modelled situations

- **SV increase**: hyperhydration rapid administration of i.v. infusion, intake of large amount of water in short time; **decrease**: dehydration, loss of blood (haemorrhage)
- **HR increase:** activation of sympathetic nervous system stress, physical activity; **decrease:** increase of vague tonus, adaptation of heart in sportsmen (athletic heart)
- **R increase:** predominance of vasoconstriction e.g. in cold environment; **decrease:** predominance of vasodilation sauna, distributive shock (anaphylaxis, adrenal crisis)
- **C higher values:** in children, young people; **lower values:** in elderly people, atherosclerosis, elastic fibers degeneration isolated systolic hypertension (systolic blood pressure is higher than normal, diastolic blood pressure is predominantly at normal level)

Mechanisms of venous return



- 1. Pressure gradient between venous system and right atrium ("a force acting from behind" vis a tergo)
- 2. Suction effect of systole ("a force acting from in front" vis a fronte)
- 3. Skeletal muscle contractions muscle pump
- 4. Suction effect of inspirium increased intraabdominal pressure and decreased intrathoracic pressure
- 5. Venous valves

Picture reference:

Slide 7 – Atlas Of Physiology, Silbernagl & Despopoulos, Georg Thieme Verlag 2003