


PEDIATRIC'S PHYSIOLOGY

Physiological peculiarities in children



GROWTH PERIODS

in relation to age

- **CHILDREN**

- ✓ Newborn: 0 –28 days after born (1 month)

- ✓ Suckling: 2 – 12 month

- ✓ Common term: **infancy**

- ✓ 1 – 4 years old

- ✓ (Toddler 1 –3 years old)

- ✓ Common term: **early childhood**

GROWTH PERIODS

- ✓ 5 – 12 years old – **late childhood**
- ✓ Other special terms:
 - ✓ Pre-school period 5 – 7 years
 - ✓ School period – younger, older

GROWTH PERIODS

- **ADOLESCENCE**

- ✓ 13 – 20 years old
- ✓ The other special terms:
 - ✓ Teenager -19 years
 - ✓ Pubertas 11-15 years

Anthropometric characteristics

body WEIGHT

- Birthweight: 2 500 - 3 800 g
 - Double birthweight: 4-5 mo
 - Triple birthweight: 1 yr
 - Quadruple birthweight: 2 yr
-
- **Pecularity in newborn: Weight loss** in first few days : 5-10% of birthweight
 - Return to birthweight: 4 -7 days of age

- AVERAGE weights:
 - at birth: 3 500 g
 - at 1 yr: 10 kg
 - at 5 yrs: 20 kg
- DAILY body weight gain – important for evaluation of nutritional state:
 - 20-30g/day for the first 3-4 months of life
 - 15-20g/day - until the end of the 1st year





Figure 1-9. Weight by age percentiles for boys, ages birth to 36 mo, including highest and lowest values at each age. (From Pomerance HH: Growth Standards in Children. New York, Harper and Row, 1979, p 25.)

Percentiles determines how many percent of population has lower values than the measured values in examined person

body HEIGHT

- AVERAGE length: 20 in (50 cm) at birth
30 in (75 cm) at 1 yr
at age 3 yr - the average child is 3 ft tall
at 4 yr - the average child is 40 in (100 cm)
tall (double birth length)
- Average ANNUAL length increase: 2-3 in
(5-7 cm) between age 4 yr and puberty

Figure 1-8. Length by age percentiles for boys, ages birth to 36 mo, including highest and lowest values at each age. (From Pomerance HH: Growth Standards in Children. New York, Harper and Row, 1979, p 29.)




HEAD CIRCUMFERENCE (HC)

- Average HC: 35 cm at birth (13.5 in)
- HC increases:
2 cm/month - for first 3 months, then slower
1 cm/month for first year

Newborn according the weeks of gestation and birth weight


- Preterm infant (premature – earlier than 38 weeks of gestation)
 - ✓ Low birthweight infants (LBW):
less than 2 500 g
 - ✓ Very low birthweight (VLBW):
less than 1 500 g
- Full-term infant (38 – 40 weeks of gestation)
birthweight 3 000 - 3 500 g, 48-52 cm length, head circumference 35cm
- Ower-term infant (41 – 42 weeks of gestation)
4000 - 6 000g, 53 - 56 cm



A

36. týden


36th week



B

38. týden

38th week



C

40. týden

40th week

G e s t a č n í v ě k
week of gestation

(podle Ushera)

The skin is covered by white muzzle – *vernix caseosa*

The skin is covered by white muzzle – *vernix caseosa*





19.04.2017 09:10

Examination of newborn at the delivery room

• Apgar score

Signs	Points		
	0	1	2
✓ Heart rate:	0	<100 /min	>100/min
✓ Respiration:	none	weak cry	vigorous cry
✓ Muscle tone	none	hypotonic-hypertonic	limb flexion
✓ reflex irritability:	none	some motion	cry, withdrawal
✓ Color of body:	blue	pink body, blue extremities	pink all over

- more than 50 years - determined the newborn hope of survival
- now - as a recommendation for nursing practice

Virginia Apgar



Physician

1909 - 1974

USA
20

TRANSITION FROM FETAL TO NEONATAL PHYSIOLOGY

- **Specialities of fetal circulation:**
 - ✓ **Placenta**, where deoxygenated blood becomes oxygenated
 - ✓ **1 Umbilical vein** – well-oxygenated blood
 - ✓ **2 Umbilical arteries** – deoxygenated blood
 - ✓ **Foramen ovale**
 - ✓ **Ductus arteriosus Botalli**
 - ✓ **Ductus venosus**




Figure 83-4. Organization of the fetal circulation. (Modified from Arey: *Developmental Anatomy*. 7th ed. Philadelphia, W. B. Saunders Company, 1974.)

RESPIRATORY SYSTEM

- Respiratory movements – started about 20 weeks of gestation

- Surfactant secretion:

- ✓ A substance normally secreted into the alveoli that **decreases the surface tension** of the alveolar fluid, therefore allowing the alveoli to open easily during inspiration
 - ✓ The surfactant secreting cells (the type II alveolar epithelial cells) started secretion about 20 weeks of gestation
- Estimation of pulmonary maturity:

ratio Lecithin/sphingomyelin production 2:1

Decrease of surfactant: Respiratory distress syndrome

NEONATAL JAUNDICE


- Bilirubin formed **in the fetus** can cross the placenta into the mother and be excreted through **the liver of the mother**

But


- Immediately after birth the only means for ridding the neonate of bilirubin is through the neonate's own liver, which for the 1st weeks has poorly functions (without any reserves), and decrease capacity for conjugating system of bilirubin and its excretion into the bile

- The plasma bilirubin concentration rises during the first 3 days of life and then gradually falls back to normal as the liver becomes functional
- This condition called **physiologic hyperbilirubineamia** and it is associated with a mild jaundice of the infant's skin and especially of the sclerae of its eyes


Normal bilirubin metabolism



Bilirubin metabolism during phototherapy




Increasing skin transmittance →



Spectrum of light

Blue most effective
(Especially around
460–490 nm)


Distance
Maximize irradiance
by minimizing
patient-to-light-source
distance



Irradiance
Standard PT:
about $10 \mu\text{W}/\text{cm}^2/\text{nm}$

Intensive PT:
 $\geq 30 \mu\text{W}/\text{cm}^2/\text{nm}$
(430–490 nm)

Skin area exposed
Maximize for intensive phototherapy
with additional light source below infant



TEMPERATURE

- **In utero** thermoregulation of the fetus is performed by the placenta, which is as an efficient heat exchanger
- **Fetal temperature** is higher than the mother's temperature: about 38.5 °C
- **After birth**, the newborn infant begins life covered by amniotic fluid and situated in a cold environment: 20-25 °C
- An infant's skin temperature may fall 0.3 °C/min and the core temperature may decline 0.1 °C/min in the delivery room

- Because the body surface area is large in relation to body mass, **heat is readily lost from the body**
- The ideal environmental temperature is called as **the neutral thermal environment**: the ambient temperature resulting in the lowest rate of heat production and the lowest consumption of oxygen by the infants while maintaining normal body temperature
- 1 hour after birth: 33-34 °C
- 1 day after birth: 31-33 °C
- 1 weeks after birth: 27-33 °C







Figure 83-7. Fall in body temperature of the neonate immediately after birth, and instability of body temperature during the first few days of life.





Empress Catherine the Great, Russia, 18th century
survival against the russian winter



Immune system

- a main prenatal immunoglobulin = **IgG**:
 - passes through the placenta
 - At the birth – the same level as in mother
 - at birth is the same concentration as in the mother's body
 - gradually decreasing its concentration
 - in 3rd to 10th week reached the lowest values then again levels increase

- **IgM** forms newborns aged 1-2 weeks
- **IgA** occurs at the age of one month, then the concentration slowly increases
(IgA is rich colostrum and breast milk)

BLOOD - composition

- **After birth:**

- ✓ **Erythrocytes = $5-6 \times 10^{12}/l$**

- ✓ **Leukocytes = $20-22 \times 10^9 /l$**

- ✓ **Hemoglobin = 190 g/l**

- **At 3 month of live:**

- ✓ **Erythrocytes = $4 \times 10^{12}/l$**

- ✓ **Leukocytes = $10.5 \times 10^9/l$**

- ✓ **Hemoglobin = 110 g/l**




Figure 5-2. Hemoglobin-oxygen dissociation curves. The position of the adult curve depends on the binding of adult hemoglobin to 2,3-diphosphoglycerate (DPG), temperature, carbon dioxide tension ($p\text{CO}_2$), and hydrogen ion concentration (pH).

CARDIOVASCULAR SYSTEM

Heart rate according age

- Newborn 135-140 beat per minute
- 6 month 130-135
- 1 year 120-125
- 2 years 110-115
- 5 years 98-100
- 8 years 80-85
- 15 years 70-76

Elektrocardiography

- More difficult evaluation than in adult
- The ECG curve is changing with respect to:
 - Ratio between right ventricle muscle/left ventricle muscle
 - Spred of activation from atrium to ventricle myocardium
 - repolarization

Evaluation of ECG curve in children is nescesary made with respect to anamnesis, clinical state and laboratory view

- **Generally accepted:**

- In newborn – predominance of the right ventricle
- To 3 month after birth – increase of left forces
- At 2 years – right and left ventricle in equilibrium
- 3 years to adult – prevalence (superiority) of left ventricle

Blood pressure

- Immediately after birth – high blood pressure:
 - Stress after delivery, increase concentration of catecholamine and cortizol
- After 1st day 70/50 mmHg:
 - Open of pulmonary and intestine circulation
- During pubertas:
 - Development of regulatory mechanism
 - Stimulation of external world

- Newborn **80/46 mmHg** 10,6/6,1 kPa
- 3 years **100/67** 13,3/8,9
- 10-11 years **111/58** 14,8/7,7
- 13-14 years **118/60** 15,7/8,0

The size of cuffs

<u>Body weight</u>	<u>age</u>	<u>size of cuff</u>
1 500 g	*	2,5 cm
5 kg	3 měsíc	4,5 cm
10 kg	15 měsíců	6 cm
30 kg	9 let	7,5 cm
30 a více kg	10 a více let	12 cm

THEORY of AGE



Elderly period

- **Earlier senior: 65 - 75 years old**
 - **Middle senior: 75 - 85 years old**
 - **Late senior: above 85 years old**
-
- **The „AGING“ is programming biological process**

Theory of „aging“

- „Free radicals“

– primary reason for aging is: damages of macromolecules and structures of cells by biochemistry reactions of free oxygen radicals

(oxygen free radicals damaging our bodies are „taxes“ that people breathe oxygen on the Earth)

- **Neuroendocrine theory**

- This theory is based on the fact, that the secretion of hormone melatonin is reduce with age (as „youth hormone“; pineal gland – coordinates of circadian rhythms)

- **Gene theory**
- Increase a lot of mutations in the cells during all of the lifetime, the mutations are a primary cause of the aging
- Theory of programming of aging is based on the idea that the function of genes is reduced in time (e.g. Apoptosis – programming death of the cells)
- Theory based on the hypothesis that exist „any genetic programme“ (Hayflick 1985 – observing the families with longevity)

The symptoms of aging

- **Reduction of function of all organ systems:**
- | loco-motor function – as general and final, decreases of forces of skeletal muscles
- reduction of capacity of the lungs, cardiac output, cardiac reserve, function of excretory system, liver, metabolism
- reduction of number of neurons in the brain (central nervous system)

- **The other symptoms:**
- Changes in places of fat deposits
- Changes of the skin hair
- Changes in the memory – main in the short-term memory
- Changes of the behavior – non-tolerance, depression

„Everyone is old, depending on how he/she feels to be old“

Thank you for your attention

