

Tissue concept and classification

Petr Vaňhara, PhD

Department of Histology and Embryology,
Faculty of Medicine MU

pvanhara@med.muni.cz

3. PŮVOD TKÁNÍ A JEJICH ROZDĚLENÍ

Tkáň lze definovat jako soubor morfologicky shodných nebo velmi podobných funkcími složkami orgánů lidského těla.

Tkáň se vyvíjejí ze zárodečných listů v průběhu embryonálního vývoje.

Mezenchym je embryonální tkáň, která vzniká z mezodermu (ektodermu) a vyplňuje tělo organismu sifovitou texturou.

Tkáň se dělí na pojivovou, svalovou, nervovou a epitelenou.

Tkáň pojivová je mezenchym, skládá se z buněk a mezibuněčné hmoty.

Tkáň svalová je mezenchym, skládá se z buněk a mezibuněčné hmoty.

Tkáň nervová je mezenchym, skládá se z buněk a mezibuněčné hmoty.

Tkáň epitelová je ektoderm, skládá se z buněk a mezibuněčné hmoty.

Tkáň pojivová
Pojivová tkáň je mezenchym
Skládá se z buněk a mezibuněčné hm.

Tkáň svalová
Svalová tkáň je mezenchym
Skládá se z buněk a mezibuněčné hm.

Tkáň nervová
Nervová tkáň je mezenchym
Skládá se z buněk a mezibuněčné hm.

Tkáň epitelová
Epitelová tkáň je ektoderm
Skládá se z buněk a mezibuněčné hm.

Část II. Čtyři základní typy tkání

Epitelová tkáň

CÍLE STUDIA
Tato kapitola by měla studentovi pomoci
- poznat čtyři základní typy tkání
- poznat strukturální a funkční charakteristiky, které odlišují epitelovou tkáň od dalších tří základních typů tkání
- poznat typy epitelové tkáně a uvést příklady míst, kde se jednotlivé typy mohou nacházet
- popsat speciální funkce jednotlivých typů epitelových buněk a uvést příklady míst, kde se jednotlivé typy mohou nacházet
- na mikrofotografiích poznat epitelu a určit jejich funkci podle struktury a lokalizace
- znát kritéria, která se užívají při klasifikaci žláz
- znát druhy žláz u člověka a uvést příklady míst, kde se mohou nacházet
- na mikrofotografiích a schématech poznat žlázy a určit jejich typ

OTÁZKY KE STUDIU
1. Vyměňte hlavní funkce epitelových tkání (II.A.1).
2. Ze kterého(ých) embryonálního(ých) zárodečného(ých) listu(ů) se epitelová tkáň vyvíjí? Uveďte příklady epitelů odvozených od jednotlivých zárodečných listů (II.H.; tabulka 4-1).
3. Vyměňte strukturální a funkční charakteristiky epitelových tkání, které je odlišují od ostatních typů tkání. Vezměte v úvahu polární buněk (IV.), specializace apikálních (IV.A.), laterálních (IV.B.) a bazálních (IV.C.) povrchů, způsob výživy (II.F.) a intenzitu mitotického dělení (II.E.).
4. Popište bazální lamínu s ohledem na její lokalizaci, složení a barvicí vlastnosti (IV.C.1.).
5. Které struktury a molekuly pomáhají připevnit epitelové buňky k základní membráně (IV.B.2.)?
6. Porovnejte bazální lamínu s mezibuněčnou hmotou (IV.B.2.).

Tissues: Concept and Classification



Úvod

Tkáň je tvořena skupinou buněk, které jsou uspořádány do souborů. Soubor stejného typu buněk spojených mezibuněčnými kontakty a mezibuněčnou hmotou tvoří tkáň. Rozlišujeme čtyři základní typy tkání: epitelu, pojivu, svalovinu a tkáň nervovou.

Epitelum je tvořeno skupinou buněk, které jsou uspořádány do souborů. Soubor stejného typu buněk spojených mezibuněčnými kontakty a mezibuněčnou hmotou tvoří tkáň. Rozlišujeme čtyři základní typy tkání: epitelu, pojivu, svalovinu a tkáň nervovou.

Pojivová tkáň. Soubor buněk s četnými vzájemnými kontakty a minimem mezibuněčné hmoty. Základní dělení: krycí epitelu, žlázové epitelu, minerální. Její uspořádání je rozhodující pro specifické biomechanické vlastnosti tkáně, jako jsou pružnost, pevnost a pružnost.

Nervová tkáň. Soubor nervových buněk včetně jejich výběžků a gliových buněk; specializována na přenos a zpracování informací, které jsou založeny na elektrochemických mechanismech.

Tkáň svalová. Je to soubor buněk schopných koordinovaných, makroskopicky patrných kontrakcí. Rozdělení: příčně pruhované svalstvo (kosterní a srdeční), hladké svalovina.

Organ je vždy tvořen z většího počtu tkání. Tkáň specifická pro organ - většinou epitelu - se označuje jako **parenchym**, na rozdíl od vazivového **stromatu**, které poskytuje organu mechanickou soudržnost a ve kterém jsou uloženy cévy (krevní a lymfatické) a nervy. Původ různých typů tkání a organů ze tří zárodečných listů (ektoderm, mesoderm, entoderm) mladého embrya je rekapitulován na str. 447.

OVERVIEW OF TISSUES

Tissues are aggregates or groups of cells organized to perform one or more specific functions.

At the light microscope level, the cells and extracellular components of the various organs of the body exhibit a recognizable and often distinctive pattern of organization. This organized arrangement reflects the cooperative effort of cells performing a particular function. Therefore, an organized aggregation of cells that function in a collective manner is called a **tissue** [Fr. *tissu*, *woven*; L. *texo*, *to weave*].

Although it is frequently said that the cell is the basic functional unit of the body, it is really the tissues, through the collaborative efforts of their individual cells, that are responsible for the body's overall function.

Histologie (z řeckého *histos* = tkáň, *logia* = studium) je nauka o stavbě tkání. Tkáň lze definovat jako komplex morfologicky podobných buněk, specializovaných k výkonu určité funkce. Jsou materiálem pro stavbu orgánů těl mnohobuněčných organismů, metazoí. Za embryonálního vývoje jedince (ontogeneze) se tkáň diferenciuje ze 3 zárodečných listů, ektodermu, entodermu a mezodermu, procesem zvaným histogeneze. Na jejím podkladě vznikají čtyři základní typy tkání: 1. **Tkáň epitelová** – vzniká ze všech tří zárodečných listů. Tvoří ji buňky těsně k sobě přiléhající s malým množstvím mezibuněčné hmoty. Uspořádána je buď v listy, kryjící povrchy, nebo v epitelové masě.

2. **Tkáň pojivová, podpurná** – pochází z mesenchymu (derivát mezodermu). Vyznačuje se hojnou účastí mezibuněčné základní hmoty, ve které jsou uloženy rozličné typy buněk, plnicí řadu funkcí.

3. **Tkáň svalová** – je původu převážně mesodermového. Tvoří ji buňky nebo syncytium. Její elementy jsou protáhlého tvaru. Jejich cytoplazma je opatřena prvky, které umožňují její kontrakci, a tím i pohyb orientovaný v příslušném směru.

4. **Tkáň nervová** – pochází z ektodermu. Její nejvýznamnější komponentou jsou nervové buňky – neurony, schopné vytvářet nervový vzruch a předávat jej z buňky na buňku.

Epitelová (epitely)

Epitelová tkáň je tvořena skupinou buněk, které jsou uspořádány do souborů. Soubor stejného typu buněk spojených mezibuněčnými kontakty a mezibuněčnou hmotou tvoří tkáň. Rozlišujeme čtyři základní typy tkání: epitelu, pojivu, svalovinu a tkáň nervovou.

Mezibuněčné spoje, kontakty epitelových buněk
Kontakty epitelových buněk je podmíněna specializací buněk v souvislosti s jejich funkcí, která je zabezpečena jejich kohezí. Koheze je způsobena volným povrchem buněk je intercelulární a utváří tzv. tmele. Na řezu vede k obrazu šestiúhelníku. Na řezu kolmém k tmele jsou tmelevisky viditelné jako tmavé body (obr. 67).

Epitelová tkáň je tvořena skupinou buněk, které jsou uspořádány do souborů. Soubor stejného typu buněk spojených mezibuněčnými kontakty a mezibuněčnou hmotou tvoří tkáň. Rozlišujeme čtyři základní typy tkání: epitelu, pojivu, svalovinu a tkáň nervovou.

Despite the variations in general appearance and structural organization, and physiologic properties of the various body organs, the tissues that compose them are classified into four basic types.

- **Epithelium (epithelial tissue)** covers body surfaces, lines body cavities, and forms glands.
- **Connective tissue** underlies or supports the other three basic tissues, both structurally and functionally.
- **Muscle tissue** is made up of contractile cells and is responsible for movement.
- **Nerve tissue** receives, transmits, and integrates information from outside and inside the body to control the activities of the body.

FOUNDING FATHERS OF HISTOLOGY – DISCOVERY OF CELLS

Robert Hooke





1665

Wow! I see
cells!!

Sche. xi

Fig. 1.



MICROGRAPHIA:

OR SOME

Physiological Descriptions

OF

MINUTE BODIES

MADE BY

MAGNIFYING GLASSES.

WITH

OBSERVATIONS and INQUIRIES thereupon.


By R. HOOKE, Fellow of the ROYAL SOCIETY.

*Non possis oculo quantum contendere Linceus,
Non tamen idcirco contemnas Lippus innugi. Horat. Ep. lib. 1.*



LONDON, Printed by *Jo. Martyn*, and *Ja. Allestry*, Printers to the
ROYAL SOCIETY, and are to be sold at their Shop at the Bell in
S. Paul's Church-yard. M DC LX V.

Xavier Bichat



1801

So different
tissues!

„I see different structures in human body. I do not need a microscope to distinguish 21 types! I will call them tissues.

In a diseased body the tissues have altered, abnormal structure!

FOUNDING FATHERS OF HISTOLOGY – MODERN CELL THEORY

Matthias Jacob Schleiden




Cells are the **basic units** of any organism

- New cells **origin** only from **other** cells
- Cells **exchange energy** (open thermodynamic system)
- Genetic **information** is **inherited** in new generations
- Chemical and structural composition of cells is generally **identical**

Theodor Schwann



J.E.P.



Robert Remak










*Omnis cellula
e cellula!*

1800-1850

Rudolf Virchow







CELL AND TISSUE VARIABILITY IN A MULTICELLULAR BODY




TISSUES AND ORGANS

- 6×10^{13} **CELLS** of **200** different types
- cells form **functional, three-dimensional**, organized **aggregations** of morphologically similar **cells** and their **products** and derivatives - **TISSUES**
- tissues constitute **ORGANS** and organ systems



Parenchyma: functional component of a tissue
(liver, lung, pancreatic, kidney parenchyma)

Stroma: surrounding, essential supportive tissue



**Example:
LIVER**

Parenchyma:

- Hepatocytes
- Sinusoids and adjacent structures


Stroma:

- Connective tissue and adjacent structures
- Vessels
- Nerves
- Bile ducts

CONTEMPORARY TISSUE CLASSIFICATION

Based on **morphology and function:**


Epithelium



Continuous, avascular layers of polarized cells with different functions, oriented to open space, with specific junctions and minimum of ECM and intercellular space.

Derivates of all three germ layers

Muscle




Myofibrils → contraction

Mesoderm – skeletal muscle, myocard, mesenchyme

– smooth muscles

Rarely ectoderm (eg. m. sphincter a m. dilatator pupillae)

Nerve




Neurons and neuroglia

Reception and transmission of electric signals

Ectoderm, rarely mesoderm (microglia)

Connective




Dominant extracellular matrix

Connective tissue, cartilage, bone...



Mesenchyme

What is a tissue?


Functional, three-dimensional, organized aggregation of **morphologically similar cells, their products and derivatives**



- classical histological definition is based on microscopic visualization




How to build a tissue?



Stem cells are essential

Stem cells are capable of **differentiation** and **self-renewal**



STEM CELLS

Totipotent

- Constitute all cells of the body incl. extraembryonic tissues
- Zygote and early stages




Pluripotent

- All cells in the body except for trophoblast
- Blastocyst – Inner cell mass - ICM (embryoblast)
- Embryonic stem cells



Multipotent



- Give rise to various cell types of a particular tissue
- Mesenchymal SC, hematopoietic SC



Oligo- a unipotent

- One or several cell types – hematopoietic, tissue precursors for renewal of intestinal epithelia, etc.



STEM CELLS



STEM CELLS IN BIOMEDICAL RESEARCH AND HUMAN BIOLOGY



Embryonic stem cells (ESCs)

- embryoblast of blastocyst
- pluripotent
- modelling of early embryogenesis, regenerative medicine




Tissue (adult) stem cells

- regeneration and renewal of tissues
- GIT, CNS, mesenchyme
- regenerative medicine, cancer biology



Induced pluripotent stem cells (iPSc)

- adult differentiated cell (fibroblast) is reprogrammed into pluripotent state
- differentiation into desired cell type
- regenerative medicine, cell and gene therapy




Nobel prize 2012

Cell

Induction of Pluripotent Stem Cells from Mouse Embryonic and Adult Fibroblast Cultures by Defined Factors



Kazutoshi Takahashi¹ and Shinya Yamanaka^{1,2,3,*}
¹Department of Stem Cell Biology, Institute for Frontier Medical Sciences, Kyoto University, Kyoto 606-8507, Japan
²CREST, Japan Science and Technology Agency, Kawaguchi 332-0012, Japan
³Contact: yamanaka@frontier.kyoto-u.ac.jp
DOI 10.1016/j.cell.2006.07.024



Disease modelling
Drug testing
Tissue replacement

...

iPSCs SHARE FUNDAMENTAL PROPERTIES WITH hESCs



STEM CELLS AS THERAPY


Age-related macular degeneration




neovascularisation



hiPSCs



Retinal pigment epithelium




Clinical trial




Cancer stem cells

- solid tumor is always heterogeneous
- small population of cells with stem cell character can repopulate tumor tissue after cytotoxic therapy




Tissue stem cells




Cancer stem cells

WHY ARE TISSUES SO DIFFERENT?



Essential terminology




- Induction of differentiation
- Determination and commitment
- Terminal differentiation




-blast




eg. myeloblast






-cyte




e.g. granulocyte






Essential mechanisms 1






Nature Reviews | Genetics



TISSUE DIFFER IN THEIR GENETIC AND EPIGENETIC PROFILES



Essential mechanisms 2





Huge number of **biological** and **physically-chemical** parameters

Stem cell niche

- Embryonic development
- Intercellular interaction
- Space organization (dimensionality)
- Gradient of morphogenes
- Epigenetic profile
- Gene expression dynamics
- Partial pressure of gases
- ECM composition
- Mechanical stimulation
- Perfusion and interstitial flows
- Local immunity response
- Metabolites

STEM CELL NICHE




- Osteoclasts
- Osteocytes
- Adipocytes
- Fibroblasts
- Stromal cells
- Vascular endothelial cells
- Immune cells
- Hematopoietic stem cells(HSC) and their differentiated progenies
- Mesenchymal stem cells (MSC)


- ECM components**
- Fibronectin
 - Laminin
 - Collagens
 - Apatite crystals (calcium 38%, phosphorus 18%)
 - Bone promoting proteins
 - Bone sialoproteins
 - Osteonectin
 - Osteoprotegerin
 - Osteocalcin
 - Integrins
 - Alcaline Phosphatase
 - Proteoglycans, Glycosaminoglycans
 - Osteopontin
 - MMPs & TIMPs
 - Receptors
 - Adhesion molecules

- Physico-chemical Effectors**
- Cytokines
 - Chemokines
 - Growth factors
 - Hormones
 - Physico-mechanical forces
 - Biochemical regulators (pH, oxygen concentration, nutrients...)

HEMATOPOIETIC NICHE



MICROENVIRONMENT IS NECESSARY FOR TISSUE HOMEOSTASIS





Apoptosis

Regeneration

Senescence



Transformation

MICROENVIRONMENT MIGHT BE CLINICALLY IMPORTANT



- Angiogenesis
- Inflammation
- Invasion and metastasis
- Self-sufficiency in growth signals
- Insensitivity to anti-growth signals

Essential mechanisms 3





Wild type



Mutant



LEWIS WOLPERT AND FRENCH FLAG MODEL








Cellular phenotype: A A B B C C



WHY DO TIGERS HAVE STRIPES?

Reaction-diffusion system



THE CHEMICAL BASIS OF MORPHOGENESIS

By A. M. TURING, F.R.S. *University of Manchester*

(Received 9 November 1951—Revised 15 March 1952)


It is suggested that a system of chemical substances, called morphogens, reacting together and diffusing through a tissue, is adequate to account for the main phenomena of morphogenesis. Such a system, although it may originally be quite homogeneous, may later develop a pattern or structure due to an instability of the homogeneous equilibrium, which is triggered off by random disturbances. Such reaction-diffusion systems are considered in some detail in the case of an isolated ring of cells, a mathematically convenient, though biologically unusual system. The investigation is chiefly concerned with the onset of instability. It is found that there are six essentially different forms which this may take. In the most interesting form stationary waves appear on the ring. It is suggested that this might account, for instance, for the tentacle patterns on *Hydra* and for whorled leaves. A system of reactions and diffusion on a sphere is also considered. Such a system appears to account for gastrulation. Another reaction system in two dimensions gives rise to patterns reminiscent of dappling. It is also suggested that stationary waves in two dimensions could account for the phenomena of phyllotaxis.

The purpose of this paper is to discuss a possible mechanism by which the genes of a zygote may determine the anatomical structure of the resulting organism. The theory does not make any new hypotheses; it merely suggests that certain well-known physical laws are sufficient to account for many of the facts. The full understanding of the paper requires a good knowledge of mathematics, some biology, and some elementary chemistry. Since readers cannot be expected to be experts in all of these subjects, a number of elementary facts are explained, which can be found in text-books, but whose omission would make the paper difficult reading.




1. A MODEL OF THE EMBRYO. MORPHOGENS

In this section a mathematical model of the growing embryo will be described. This model will be a simplification and an idealization, and consequently a falsification. It is to be hoped that the features retained for discussion are those of greatest importance in the present state of knowledge.

The model takes two slightly different forms. In one of them the cell theory is recognized but the cells are idealized into geometrical points. In the other the matter of the organism is imagined as continuously distributed. The cells are not, however, completely ignored, for various physical and physico-chemical characteristics of the matter as a whole are assumed to have values appropriate to the cellular matter.




TISSUE PATTERNS ARE DRIVEN BY GRADIENTS OF MORPHOGENES



Belousov-Zabotinsky

HOX COMPLEX



Hox genes


Highly conserved family of transcription regulators that determine body polarity, orientation and axis

Tissue differentiation along antero-posterior axis







Human (39 genes)

Cluster	Chromosome	# Hox genes
HoxA	7	11
HoxB	17	10
HoxC	12	9
HoxD	2	9


doi:10.1038/sj.hdy.6800872




TEMPORO-SPATIAL EXPRESSION OF MORPHOGENES DRIVES FINAL LOCALIZATION, ORIENTATION AND MORPHOLOGY OF TISSUES AND ORGANS



MANIPULATING AER ALTERS INSTRUCTIONS FOR LIMB DEVELOPMENT





MORPHOGENES FROM AER AND ZPA DEFINES LIMB FORMATION





HOX PATTERN DRIVES TRANSCRIPTIONAL RESPONSE



(A) Phase I
Stylopod








(B) Phase II
Zeugopod





(C) Phase III
Autopod



Hox paralogue groups





STORY OF THALIDOMID



Proliferation

Vascularisation



Thalidomid



Thalidomid embryopathy

- phocomelia
- amelia
- anophthalmia/microphtalmia
- abnormal kidneys, heart, GIT, genitalia

Frances Oldham Kelsey,
FDA USA




HISTOGENESIS AND ORGANOGENESIS

Ectoderm

Endoderm

Mesoderm



EMBRYONIC DEVELOPMENT

Ectoderm

Surface ectoderm

- Epidermis, hair nails, cutaneous and mammary glands
- Corneal epithelium and lens of eye
- Enamel of teeth
- Internal ear
- Anterior pituitary gland
- Epithelium of oral cavity and part of anal canal

Neuroectoderm

- **Neural tube** and derivatives
 - CNS
 - Retina
 - Posterior pituitary gland
 - Pineal body
- **Neural crest** and derivatives:
 - Cranial and sensory ganglia and nerves
 - Schwann cells
 - adrenal medulla
 - Enteroendocrine cells
 - Melanocytes
 - Head mesenchyme and connective tissue
 - Odontoblasts

Mesoderm

head

- Connective tissue of head
- Cranium, dentin

Paraxial

- Skeletal muscle of trunk and limbs except cranium
- Dermis of skin
- Muscles of head

Intermediate

- Urogenital system + ducts, glands and gonads


Lateral


- Visceral muscle and connective tissue
- Serous membranes of pleura, peritoneum and pericardium
- Blood cells, leukocytes
- Cardiovascular and lymphatic system
- Spleen
- Adrenal cortex

Endoderm

- GIT epithelium except oral cavity and part of anal canal
- Extramural glands of GIT
- Epithelium of bladder
- Epithelium of respiratory system
- Thyroid gland, parathyroid glands, thymus
- Tonsils
- Epithelium of cavum tympani and Eustachian tube



COFFEEBREAK






CONNECTIVE TISSUE

Not just a tissue glue...




Mechanical and biological properties

→ surrounds other tissues, allows compartmentalization, provides support, defines physico-chemical environment, brings immunological support, provides storage of energy, ...



Cells and extracellular matrix (ECM)

- **Cells**
 - **Connective tissue** – permanent and transient cell populations (e.g. fibroblasts/myofibroblasts, immune cells, adipocytes, adult stem cells)
 - **Cartilage** – chondroblasts/chondrocytes
 - **Bone** – osteoblasts/osteocytes/osteoclasts
- **Extracellular matrix** – fibrous and amorphous
 - **Fibrous component**
 - collagen fibers (prototypically col. I, II)
 - reticular
 - elastic
 - **Amorphous component** (amorphous ground substance)
Complex matrix consisting of
 - glycosaminoglycans
 - glycoproteins
 - proteoglycans



Specific composition depends on a tissue type (connective × ligament × cartilage × bone)

CLASSIFICATION OF CONNECTIVE TISSUE

Collagen	Structure	Function and distribution
Loose collagen CT	Abundant ground substance, few collagen fibers with random arrangement	Microvascularisation Innervation
<u>Irregular dense collagen CT</u>	Few ground substance, few cells, many collagen fibers, random arrangement	Mechanically resistant organ capsules
<u>Regular dense collagen CT</u>	Tightly arranged collagen fibers with fibroblasts intercalated between them	Part of musculoskeletal system. Tendons, ligaments
Embryonic		
Mesenchyme	Undifferentiated cells uniformly dispersed in the ground substance, few collagen fibers	Undifferentiated progenitors
Wharton's jelly	Viscous amorphous matrix with collagen fibers. Fibroblasts.	Matrix of umbilical cord
Special		
Reticular CT	Network of collagen III fibers and reticular cells	Support of hematopoietic and lymphatic cells
Elastic	Rich in elastic fibers	Flexible support to the elastic arteries and aorta
Adipose	Adipocytes	Energy storage (white fat), heat production (brown fat)
Cartilage	Chondroblasts, chondrocytes	Mechanical support
Bone	Osteoblasts, osteocytes, osteoclasts	Mechanical support, calcium and phosphate metabolism
Blood	See lecture on blood & hematopoiesis this semester	








LOOSE COLLAGEN CONNECTIVE TISSUE

Cells

- Fibroblasts/fibrocytes/myofibroblasts
- Heparinocytes
- Macrophages of CT = histiocytes
- Plasma cells
- Lymphocytes
- Adipocytes
- Adult stem cells



Extracellular matrix

- Fibrous compound
- Amorphous ground substance





CELLS OF LOOSE COLLAGEN CONNECTIVE TISSUE

Mesenchymal (adult) stem cells




Collagen fibers

- family of fibrous proteins encoded by >35 genes (2013)
- polymer – subunit = tropocollagen; triple helix
- different structural and mechanical properties (strength, elasticity, pliability...)
- most abundant protein in human body (30% dry weight)



COLLAGEN



COLLAGEN



Type	Localization	Structure	Main function
I	Bone, tendons, meniscus, dentin, dermis, capsules of organs, loose CT 90% of type I	Fibrils (75nm) – fibers (1-20µm)	Resilience in pull
II	Hyaline and elastic cartilage	Fibrils (20nm)	Resilience in pressure
III	Skin, veins, smooth muscles, uterus, liver, spleen, kidney, lung	Like I, high content of proteoglycans and glycoproteins, reticular network	Shape formation
IV	Basal lamina of epithelium and endothelium, basal membranes	No fibrils or fibers	Mechanical support
V	Lamina of muscle cells and adipocytes, fetal membranes	Like IV	
VI	Interstitial tissue, chondrocytes – adhesion		Connecting dermis and epidermis
VII	Basal membrane of epithelium		
VIII	Some endothelia (Cornea)		
IX, X	Growth plate, hypertrophic and mineralized cartilage		Growth of bones, mineralization

COLLAGEN IN LIGHT MICROSCOPE

HE

HES


AZAN



Julian Voss-Andreae
"Unraveling Collagen"



2005

Orange Memorial Park
Sculpture Garden, City of
South San Francisco, CA




ELASTIC FIBERS

- less abundant than collagen
- polymer – tropoelastin
- minimal tensile resistance, loss of elasticity if overstretched
- reduction of hysteresis = allow return back to original state after mechanic change



RETICULAR FIBERS

- collagen 3D meshwork
- bone marrow, spleen, lymphatic nodules
- microenvironment for e.g. hematopoietic stem cells and progenitors





RETICULAR CONNECTIVE TISSUE



EXTRACELLULAR MATRIX – GROUND SUBSTANCE

Amorphous extracellular matrix

Colorless, transparent, homogenous substance consisting of glycosaminglycans, proteoglycans and structural glycoproteins




GLYCOSAMINOGLYCANS

linear polysaccharides composed of two disaccharide subunits
– **uronic acid and hexosamine**

polysaccharides rich in hexosamines = acid mukopolysaccharides

glucuronic or iduronic acid



glucosamin or galactosamin

GLYCOSAMINOGLYCANS

They bind to protein structures (except for hyaluronic acid)

Glyc


Hyalu

Chon

Derm

Hepa

Kerat




reum,

s


PROTEOGLYCANS

- protein + dominant linear saccharide component
- proteoglycan aggregates
- water-binding, volume dependent of hydration
- aggrecan (cartilage)
- syndecan
- fibroglycan




STRUCTURAL GLYCOPROTEINS



- dominant protein + branched saccharide component
- interaction between cells and ECM



- **fibronectin** – connects collagen fibers and glykosaminoglycans, cell adhesion and migration
- **laminin** – basal lamina – epithelial integrity
- **chondronectin** – cartilage – adhesion of chondrocytes to collagen



COMPOSITION OF ECM




CLASSIFICATION OF SPECIALIZED CONNECTIVE TISSUE

**Dense
Connective Tissue**



**Adipose Tissue
(Connective Tissue)**




**Areolar Tissue
(Connective Tissue)**



**Compact Bone
(Connective Tissue)**




**Blood
(Connective Tissue)**




ADIPOSE TISSUE

- Adipocytes, fibroblasts, reticular, collagen and elastic fibers, capillaries
- White and brown adipose tissue




WHITE ADIPOSE TISSUE

- adipocytes are actively formed until 2nd year of life
- no innervations, but rich vascularisation
- adipocytes with only one lipid droplet
- leptin (adipokinins)



BROWN ADIPOSE TISSUE




- fetus and children up to 1st year of life
- fast source of energy
- typical localization – between shoulder blades, axilla, mediastinum, around kidneys, pancreas, small intestine
- small cells with numerous fat droplets







EMBRYONIC ORIGIN OF CONNECTIVE TISSUE

- Mesenchyme = loose tissue between germ layers
- Complex network of star- or spindle-shaped cells
- Jelly-like amorphous ground substance


DAY 12 of embryonic development




DERIVATIVES OF MESENCHYME




DERIVATIVES OF MESENCHYME







MESENCHYMAL STEM CELLS



APPLICATIONS OF MESENCHYMAL STEM CELLS




FURTHER STUDY



Guide to General Histology and Microscopic Anatomy

Petr Vaňhara, Miroslava Sedláčková,
Irena Lauschová, Svatopluk Čech, Aleš Hampel



Masaryk University, Brno 2017

<http://www.histology.med.muni.cz>

pvanhara@med.muni.cz

Thank you for attention