

**M U N I
M E D**


Physiology of the respiratory system.

Questions for the oral exam

- A22: Hypoxia and ischemia
- A25: Lung ventilation, volumes, measurement
- A26: Dead space, measurement
- A27: Resistance of airways, measurement
- A28: Maximal respiratory flow - volume curve (spirogram)
- A45: Alveolar surface tension. Surfactant
- A46: Compliance of lungs. Respiratory work. Pneumothorax
- A47: Composition of atmospheric and alveolar air. Gas exchange in lungs and tissues
- A48: Transport of O₂. Oxygen - haemoglobin dissociation curve. Transport of CO₂
- A49: Regulation of ventilation
- A50: Respiratory responses to irritants

A22: Hypoxia and ischemia

- Hypoxia is a general name for a lack of oxygen in the body or individual tissues
- Ischemia, meaning insufficient blood flow to a tissue, can also result in hypoxia
- The most common types of hypoxia:
 - Hypoxic
 - Transport (anemic)
 - Ischemic (stagnation)
 - Histotoxic



ERY: ♀ $3.4 - 4.4 \cdot 10^{12}/l$


♂ $4.5 - 5.5 \cdot 10^{12}/l$

$pO_2: 21kPa$

A22: Hypoxia and ischemia


– Hypoxic:

- physiological: stay at higher altitudes
- $\downarrow pO_2$; \uparrow Ery
- pathological: hypoventilation during lung or neuromuscular diseases
- \downarrow ventilation; $\downarrow pO_2$; \uparrow Ery




– Transport (anemic):

- reduced transport capacity of blood for oxygen (anemia, blood loss)
- $\downarrow pO_2$; \downarrow Ery/Hb




– Ischemic (stagnation):

- restricted blood flow to tissue (heart failure, obstruction of an artery)
- $\downarrow pO_2$; \uparrow Ery




– Histotoxic

- cells are unable to utilize oxygen (cyanide poisoning)
- $\downarrow pO_2$; \uparrow Ery




A25: Lung ventilation, volumes, measurement

- *Ventilation*, or breathing, is the movement of air through the conducting passages between the atmosphere and the lungs
- *Principle*: determination the air flow velocity from the measured pressure differences between the inner and outer spirometer membranes, the volumes being calculated (PowerLab spirometry)




A25: Lung ventilation, volumes, measurement



- **Tidal volume (TV)** – the volume of air that enters the lungs during each inspiration (or the volume that is exhaled during every expiration).
- **Inspiratory reserve volume (IRV)** – the maximal amount of additional air that can be drawn into the lungs by determined effort after a normal inspiration at rest.
- **Expiratory reserve volume (ERV)** – the additional amount of air that can be exhaled from the lungs by determined effort after a normal expiration.
- **Residual volume (RV)** – the volume of air still remaining in the lungs after the most forcible expiration possible.

A25: Lung ventilation, volumes, measurement



Lung capacity:


- $VC = VT + IRV + ERV$
- $TLC = VC + VC$
- $FRC = ERV + RV$
- $IC = IRV + VT$
- $EC = ERV + VT$

Dynamic lung volumes:

- VE
- MMV


A25: Lung ventilation, volumes, measurement

Dynamic lung volumes



- **FVC** – the maximum volume of air that can be exhaled after maximum inhale
- **FEV_1** – the volume of air exhaled with the greatest effort in 1 second after maximum inhale
- **FEV_1/FVC (%)** – Tiffeneau index – around 0,8 (80 %)


A25: Lung ventilation, volumes, measurement



Obstructive lung disease

($FVC=N$; $FEV_1=\downarrow$)

- tracheal stenosis
- astma bronchiale
- CHOPN
- tumor



Restrictive lung disease

($FVC=\downarrow$; $FEV_1=N$)


- ### Pulmonary etiology
- pulmonary fibrosis
 - lung resection
 - pulmonary edema
 - pneumonia

Extrapulmonary etiology

- ascites
- kyphoscoliosis
- burns
- high diaphragm condition

A25: Lung ventilation, volumes, measurement


Helium dilution method – residual volume



$$C = \frac{n}{V}$$

$$V_1 \times C_1 = (RV + V_1) \times C_2$$

$$RV = \frac{V_1 \times C_1}{C_2} - V_1$$




A28: Maximal respiratory flow - volume curve (spirogram)


Principle: the measurement of the air flow velocity according to the speed of the turbine and the volumes are calculated (Cosmed).




- **PEF** – peak expiratory flow; the highest speed of air flow at peak of exhale
- **MEF** – maximum expiratory flow rates at different FVC levels, which is still to be exhaled (75 %, 50 % and 25 % of FVC)




A26: Dead space, measurement




[N_2] PROFILE OF EXPIRED AIR WITH NO MIXING



MEASURED [N_2] PROFILE



A26: Dead space, measurement



Bohr equation

$$\text{Dead space } V_D = \frac{V_T (F_{A_{CO_2}} - F_{E_{CO_2}})}{F_{A_{CO_2}}}$$

Using normal values:


$$V_D = \frac{0.5 (0.056 - 0.040)}{0.056}$$

$$\text{Dead space } V_D = 143 \text{ mL}$$

A27: Resistance of airways, measurement

Pneumotachograph:

- tubes of the same diameter, parallel arranged
- measures the differences in air pressure at the beginning and end of the pneumotachograph in proportion to the velocity of the inhaled or exhaled air




$$\dot{V} = \frac{\Delta P}{R}$$

$$\Delta P_p = P_p - P_{atm}$$



$$\Delta P_{alv} = P_{alv} - P_{atm}$$

$$\frac{P_p - P_{atm}}{R_p} = \dot{V} = \frac{P_{alv} - P_p}{R_d}$$

$$R_d = R_p \cdot \left(\frac{\Delta P_{alv}}{\Delta P_p} - 1 \right)$$



A46: Compliance of lungs. Respiratory work. Pneumothorax



A46: Compliance of lungs. Respiratory work. Pneumothorax


– According to etiology:

- **traumatic** pneumothorax (due to an injury) occurs if the chest wall is perforated or during an injury of the esophagus, bronchi, and during rib fractures.
- **spontaneous** pneumothorax
- **primary** idiopathic pneumothorax (without any known cause) may occur in tall healthy young men with an incidence of pneumothoraxes in the family,
- **secondary** pneumothorax arises as a consequence of lung diseases (such as COPD or cystic fibrosis),
- **iatrogenic** pneumothorax (due to medical procedures) occurs during invasive medical examinations such as transparietal aspiration biopsy, subclavian vein catheterization, or mechanical ventilation with positive pressure.
- **artificially induced** (deliberate) pneumothorax is used during thoracoscopy, an endoscopic examination the thoracic cavity.


– According to the communication of the pleural space with its surroundings

- **open pneumothorax** (when the hole in the pleural space remains open, the air in the pleural cavity moves back and forth with each breath of the patient)
- **closed pneumothorax** (when a small opening through which air enters the pleural cavity closes)
- **valvular pneumothorax** (the tissue of the lungs or the chest wall covers the hole in such a way that a valve emerges, this valve allows air to flow inside during inspiration, but it prevents the air from leaving the pleural cavity during exhalation).

A45: Alveolar surface tension. Surfactant



A



B

- pneumocytes typ II
- reduces the surface tension depending on the size of the alveolus
- increases lung compliance, reduces breathing work

The Laplace law (in constant tension):

the alveolus with bigger radius has lower pressure

→ the air would move from a smaller alveolus to a bigger one


→ collapse of smaller alveoli

$$P = \frac{2T}{r}$$

A46: Compliance of lungs. Respiratory work. Pneumothorax

Respiratory system resistance



- Elastic resistance:
 - elastic fibers
 - alveolar surface tension
- Nonelastic resistance:
 - viskose resistance
 - airway resistance




Respiratory work:

- Elastic
- Viskose
- Work of airway resistance


A46: Compliance of lungs. Respiratory work. Pneumothorax



Pt: P_{atm} and P_{pl}





Pt: P_{alv} and P_{pl}



Pt: P_{atm} and P_{alv}

A46: Compliance of lungs. Respiratory work. Pneumothorax



Respiratory work:
1 – elastic
2 – viscos
3 – airway resistance

Case report

Herman Neiswander is a 65-year-old retired landscape architect. One cold January morning, he decided to warm his car in the garage. Forty minutes later, Mr. Neiswander's wife found him slumped in the front seat of the car, confused and breathing rapidly. He was taken to a nearby emergency department, where he was diagnosed with acute carbon monoxide (CO) poisoning and given 100% O₂ to breathe. An arterial blood sample had an unusual cherry-red color. The values obtained in the blood sample are given in table below.

pO ₂	660 mm Hg (normal, 100 mm Hg)
pCO ₂	36 mm Hg (normal, 40 mm Hg)
% O ₂	50 (normal, 95–100)
Pulse oximetry	100%

Case report

1. Why was Mr. Neiswander's O₂ saturation reduced to 50%?
2. If Mr. Neiswander's % O₂ saturation was 50%, why was his pulse oximetry value 100%?
3. What percentage of the heme groups on his hemoglobin were bound to CO?
4. O₂-hemoglobin dissociation curve. What effect did CO poisoning have on the affinity of hemoglobin for O₂?
5. How did CO poisoning alter O₂ delivery to Mr. Neiswander's tissues?
6. What was the rationale for giving Mr. Neiswander 100% O₂ to breathe?

A47: Composition of atmospheric and alveolar air. Gas exchange in lungs and tissues.

COMPOSITION OF DRY ATMOSPHERIC AIR

O ₂	20.95 %	F _{O₂}	≅ 0,21
N ₂	78.09 %	F _{N₂}	≅ 0,78
CO ₂	0.03 %	F _{CO₂}	≅ 0,0004

BAROMETRIC PRESSURE IN SEA LEVEL


1 atmosphere = 760 mm Hg

PARTIAL PRESSURE OF DRY AIR IN SEA LEVEL


$$\begin{aligned} P_{O_2} &= 760 \times 0,21 &= \sim 160 \text{ mm Hg} \\ P_{N_2} &= 760 \times 0,78 &= \sim 593 \text{ mm Hg} \\ P_{CO_2} &= 760 \times 0,0004 &= \sim 0,3 \text{ mm Hg} \end{aligned}$$

23 1 kPa = 7,5 mm Hg (torr)

A47: Composition of atmospheric and alveolar air. Gas exchange in lungs and tissues.



A48: Transport of O₂. Oxygen - haemoglobin dissociation curve. Transport of CO₂




O₂ is transported in two forms :

- physically dissolved(1%)
- in chemical bond with Hb (99%)

- Fetal hemoglobin(2 α , 2 γ)
- Methemoglobin (Fe³⁺)
- Carboxyhemoglobin (CO)
- Carbaminohemoglobin (CO₂)
- Oxyhemoglobin (O₂)
- Deoxyhemoglobin (without any gases)


A48: Transport of O₂. Oxygen - haemoglobin dissociation curve. Transport of CO₂



Dissociation curve of Hb is influenced by:



- pH of blood
- pCO₂ of blood
- Temperature
- Concentration of 2,3 - BPG

A48: Transport of O₂. Oxygen - haemoglobin dissociation curve. Transport of CO₂




CO₂ is transported in next forms :


- physically dissolved (5 %)
- in the form of bicarbonate anions (85%)
- in chemical bond with Hb (10%)




A49: Regulation of ventilation




A49: Regulation of ventilation



A49: Regulation of ventilation



A49: Regulation of ventilation



A50: Respiratory responses to irritants

The lungs are protected from damage by:

- presence of hair (vibrissae) in the nasal cavity (traps dust particles)
- presence of ciliary epithelium covered with mucus (cilia moving mucus in one direction - into the pharynx)
- pulmonary alveolar macrophages
- presence of antibodies in bronchial secretion (IgA)

Reflexes:

- Herring-Breuer reflexes (inflation/deflation)
- Sneeze reflex
- Cough reflex
- Hiccup
- Yawn

A50: Respiratory responses to irritants

Cough Reflex	Sneeze Reflex	Hiccup
<p>Cough is an expulsive reflex that protects the lungs and respiratory passage from foreign bodies.</p> <p><i>Causes of cough:</i></p> <ul style="list-style-type: none"> - Irritants-smokes, fumes, dusts, etc. - Diseased conditions like COPD, tumors of thorax, etc. <p><i>Pathway for cough reflex:</i></p> <ul style="list-style-type: none"> - Receptors in nose, paranasal sinuses, pharynx, trachea, pleura, diaphragm, perichondrium, stomach, ex.auditory canal and tympanic membrane - V,IX,X cranial nerves and phrenic nerves - medulla - X cranial nerve, phrenic nerve, spinal motor nerve - primary and accessory respiratory muscles 	<p>Sneeze is defined as the involuntary expulsion of air containing irritants from nose.</p> <p><i>Causes of sneeze:</i></p> <ul style="list-style-type: none"> - Irritation of nasal mucosa - Excess fluid in airway <p><i>Pathway for sneeze reflex:</i></p> <ul style="list-style-type: none"> - Olfactory receptors or V cranial nerve endings - I and V cranial nerve - medulla – nucleus solitarius and reticular formation - V, VII, IX, X cranial nerves and intercostal muscles - pharyngeal, tracheal and respiratory muscles 	<p>Hiccup is spasmodic contraction of the diaphragm which causes a sudden intake of breath that is involuntarily cut off by closure of the glottis, thus producing a characteristic sound.</p> <p><i>Causes of hiccup:</i></p> <ul style="list-style-type: none"> - Eating too fast or too much - Strokes, brain tumors, damage to the vagus or phrenic nerve - Anxiety and stress <p><i>Pathway for sneeze reflex:</i></p> <ul style="list-style-type: none"> - Phrenic, vagus, and sympathetic nerves - Midbrain - Motor fibers of phrenic nerve and accessory nerves - Diaphragm and intercostal muscles

A50: Respiratory responses to irritants

Herring-Breuer reflexes (inflation/deflation)

- *a. keeps the lungs from over-inflating with inspired air*
 - *pulmonary stretch R – vagus nerve – medulla – inhibition of inspiration and initiation of expiration*
- *b. serves to shorten exhalation when the lung is deflated*
 - *pulmonary stretch R – vagus nerve – the pontine center*