


# **Regional Circulations**

**(pulmonary, skin, muscle,  
cerebral, splanchnic, renal, fetal)**

Assoc. Prof. MUDr. Markéta Bébarová, Ph.D.

Department of Physiology, Faculty of Medicine, Masaryk University



# Regional Circulation

- an organ may be supplied by two blood inflows:
  - the nutrient circulation
  - the functional circulation
- various ways of **anatomical and functional adaptation** of an organ-specific circulation to provide the optimal function of the organ
- varying impact of particular ways of regulation of the blood flow (~ vasal tone) in various organs

# Pulmonary Circulation

# Pulmonary Circulation

- Blood flow through lungs is virtually equal to the blood flow through all other organs.
- Functions:
  - provide the gas exchange
  - blood reservoir
  - mechanical, chemical and immunological filter


# Pulmonary Circulation

- **Arteries** (differences compared to the arteries in the systemic circulation)
  - bigger total cross-section of all pulmonary arteries
  - smaller thickness of the vessel walls
  - high compliance
- **Capillaries**
  - wide, abundant anastomoses form a net surrounding alveoles
  - time of passage, area of perfused capillaries at rest and intensive exertion
- **Veins**
  - high compliance (blood reservoir, orthopnoe)

Blood pressure in pulmonary circulation

# Pulmonary Circulation

- Nutrient circulation
  - physiological arteriovenous shunt



Ganong's Review of Medical Physiology, 23<sup>rd</sup> edition

- Lymphatic vessels
  - fast transport of proteins and various particles from the peribronchial and perivascular tissue → ↓ formation of the tissue fluid ~ prevention of the pulmonary edema

No filtration in pulmonary capillaries physiologically!

1. pressures in interstitium and pulmonary capillaries
2. permeability of pulmonary capillaries

# Pulmonary Circulation


- Regulation of blood flow in lungs
  - A. Systemic mechanisms
    - 1) Neural regulation (sympathicus, parasympathicus)
    - 2) Humoral regulation (circulating substances)
  - B. Local mechanisms
    - chemical (metabolic) autoregulation
      - opposite reaction compared to systemic circulation (vasoconstriction)
  - C. Passive factors
    - cardiac output
    - gravity (blood distribution in lungs)

# Pulmonary Circulation

- Ratio of ventilation and perfusion

- kept constant (local metabolic autoregulation)
  - non-ventilated alveolus - vasoconstriction
  - non-perfused alveolus - bronchoconstriction
- decreased ratio -
  - most often cause of hypoxic hypoxia in clinical practise (right-left shunt) → ↓ arterial blood saturation with O<sub>2</sub>

Ganong's Review of Medical Physiology, 23<sup>rd</sup> edition



- content of CO<sub>2</sub> usually not changed (compensatory hyperventilation in other alveoles)

# Skin Circulation

# Skin Circulation

- Skin blood flow considerably varies (0.02-5 l/min).

## Function:

- Metabolic demands of skin – small (*decubitus*)
- **Maintenance of body temperature**


poikilothermic tissue

## Arteriovenous anastomoses

- Protection against environment
- Maintenance of mean blood pressure

# Skin Circulation

- Arteriovenous anastomoses
  - convoluted muscle vessels directly connecting arteriols and venules (low-resistance shunt)



Honzíková N - Poznámky k přednáškám z fysiologie (1992)

- regulated by sympathetic vasoconstrictive nerve fibers

# Skin Circulation

- Regulation of skin blood flow:
  - Sympathetic nerve fibers
  - Humoral – local factors  
(histamine, serotonin)

# Skin Circulation


- Reaction on a temperature change:
  - 1) direct impact of a temperature change on the vessel tone
  - 2) excitation of skin thermoreceptors
  - 3) excitation of thermoreceptors in brain



reflex modulation of  
sympathetic vasoconstrictive activity

# Skin Circulation

- Axon reflex



Ganong's Review of Medical Physiology, 23<sup>rd</sup> edition.

# Muscle Circulation

# Muscle Circulation

- Function:
  - 1) Blood supply of muscles

the resting blood flow – 18% of the cardiac output vs. even 90% at intensive exertion (the local blood flow ↑ even 20times)
  - 2) Regulation of blood pressure

skeletal muscles – 40% of the body weight → resistance of the muscle bloodstream has a high impact on the total peripheral resistance
- The blood flow during muscle activity is intermittent, during the tetanic contraction even zero (oxygen debt).

# Muscle Circulation

- Regulation of the muscle blood flow:

## 1) Neural regulation

dominates at rest (vasoconstriction through sympathetic - big dilation reserve)

## 2) Local chemical regulation

dominates at physical exertion (metabolic vasodilation)

almost linear increase of the flow with increasing metabolic activity

increased blood flow + increased O<sub>2</sub> extraction

↑ capillary pressure + ↑ osmolarity → ↑ filtration → edema in active muscles

# Cerebral Circulation

# Cerebral Circulation

**TABLE 34–1** Resting blood flow and O<sub>2</sub> consumption of various organs in a 63-kg adult man with a mean arterial blood pressure of 90 mm Hg and an O<sub>2</sub> consumption of 250 mL/min.

Region	Mass (kg)	Blood Flow		Arteriovenous Oxygen Difference (mL/L)	Oxygen Consumption		Resistance (R units) <sup>a</sup>		Percentage of Total	
		mL/min	mL/100 g/min		mL/min	mL/100 g/min	Absolute	per kg	Cardiac Output	Oxygen Consumption
Liver	2.6	1500	57.7	34	51	2.0	3.6	9.4	27.8	20.4
Kidneys	0.3	1260	420.0	14	18	6.0	4.3	1.3	23.3	7.2
Brain	1.4	750	54.0	62	46	3.3	7.2	10.1	13.9	18.4
Skin	3.6	462	12.8	25	12	0.3	11.7	42.1	8.6	4.8
Skeletal muscle	31.0	840	2.7	60	50	0.2	6.4	198.4	15.6	20.0
Heart muscle	0.3	250	84.0	114	29	9.7	21.4	6.4	4.7	11.6
Rest of body	23.8	336	1.4	129	44	0.2	16.1	383.2	6.2	17.6
Whole body	63.0	5400	8.6	46	250	0.4	1.0	63.0	100.0	100.0

<sup>a</sup>R units are pressure (mm Hg) divided by blood flow (mL/s).

Reproduced with permission from Bard P (editor): *Medical Physiology*, 11th ed. Mosby, 1961.

# Cerebral Circulation

- provides:
  - 1) constant sufficient blood supply  
(black-out during several seconds of the brain ischemia,  
irreversible damage during several minutes)
  - 2) dynamic blood redistribution  
(metabolic hyperaemia)

# Cerebral Circulation

- Anatomical specialities of cerebral circulation:
  - 1) *circulus arteriosus cerebri*  
(interconnection of main cerebral arteries by anastomoses)
  - 2) very high density of capillaries  
(3000 – 4000 capillaries / mm<sup>2</sup> od the grey matter)  
~ minimalization of diffuse distance for gases and other substances
  - 3) very short arteriols  
(almost 1/2 of the vasa resistance falls on arteries which are abundantly innervated)

# Cerebral Circulation

- Functional adaptation of cerebral circulation:
  - 1) high and stable blood flow
  - 2) high O<sub>2</sub> extraction
  - 3) well developed autoregulation (myogenic and metabolic)
  - 4) high reactivity on changes of CO<sub>2</sub> concentration
  - 5) local vs. total hypoxia
  - 6) innervation



# Cerebral Circulation

- Special physical conditions of cerebral circulation:
  - 1) solid cover of brain by skull
    - Monro-Kelli theory
      - flow may be increased only by acceleration of the blood flow, not by an increase of capacity of the bloodstream
      - Cushing reflex
  - 2) gravity
    - orthostatic reaction (postural syncope)

# Cerebral Circulation

- Blood-brain barrier

cerebral capillaries – tight inter-endothelial connections



Ganong's Review of Medical Physiology, 23<sup>rd</sup> edition

# Cerebral Circulation

- Blood-brain barrier

**By free diffusion:**

- lipophilic substances ( $O_2$ ,  $CO_2$ , xenon; unbound forms of steroid hormones)
- water (aquaporins; osmolality of blood and cerebrospinal fluid is identical!)
- glucose – the main source of energy for neurons (free diffusion would be slow – accelerated by GLUT)

**By transcellular transport (regulated):**

- ions (e.g.  $H^+$ ,  $HCO^{3-}$  vs.  $CO_2$  !)
- transporters for thyroid hormones, some organic acids, choline, precursors of nucleic acids, aminoacids, ...

# Cerebral Circulation

- Blood-brain barrier
- Functions:
  - maintenance of constant composition of the neuron environment
  - protection of brain against endogenic and exogenic toxins
  - prevention of loss of neurotransmitters to the bloodstream

# Cerebral Circulation

- Cerebrospinal fluid

- localization
- composition
- volume ~150 ml,  
rate of production ~550 ml/d  
(exchange 3.7times/day)


Substance		CSF	Plasma	Ratio CSF/Plasma
Na <sup>+</sup>	(meq/kg H <sub>2</sub> O)	147.0	150.0	0.98
K <sup>+</sup>	(meq/kg H <sub>2</sub> O)	2.9	4.6	0.62
Mg <sup>2+</sup>	(meq/kg H <sub>2</sub> O)	2.2	1.6	1.39
Ca <sup>2+</sup>	(meq/kg H <sub>2</sub> O)	2.3	4.7	0.49
Cl <sup>-</sup>	(meq/kg H <sub>2</sub> O)	113.0	99.0	1.14
HCO <sub>3</sub> <sup>-</sup>	(meq/L)	25.1	24.8	1.01
PCO <sub>2</sub>	(mm Hg)	50.2	39.5	1.28
pH		7.33	7.40	...
Osmolality	(mosm/kg H <sub>2</sub> O)	289.0	289.0	1.00
Protein	(mg/dL)	20.0	6000.0	0.003
Glucose	(mg/dL)	64.0	100.0	0.64
Inorganic P	(mg/dL)	3.4	4.7	0.73
Urea	(mg/dL)	12.0	15.0	0.80
Creatinine	(mg/dL)	1.5	1.2	1.25
Uric acid	(mg/dL)	1.5	5.0	0.30
Cholesterol	(mg/dL)	0.2	175.0	0.001

# Cerebral Circulation

- Cerebrospinal fluid

Function:


- protection of brain  
(together with meninges)



Ganong's Review of Medical Physiology,  
23<sup>rd</sup> edition

# Cerebral Circulation


- Cerebrospinal fluid



Ganong's Review of Medical Physiology, 23<sup>rd</sup> edition

# Cerebral Circulation

- **Paraventricular organs**
  - ~ brain regions where the blood-brain barrier is missing (fenestrated capillaries)
  - secretion of **polypeptides** (oxytocin, vasopressin, ...),
  - **chemoreceptive zones (AP)**
  - **osmoreceptive zones (OVLT)**



Ganong's Review of Medical Physiology, 23<sup>rd</sup> edition

# Splanchnic Circulation

# Splanchnic Circulation


- blood flow through GIT including liver and pancreas
- blood flow through spleen
- Main functional roles:
  - metabolic function of GIT
  - blood reservoir
  - special (e.g. spleen – removal and degradation of old/altered erythrocytes)

# Splanchnic Circulation

- Blood reservoir
- at rest ~20% of the total blood volume
- rich innervation with sympathetic vasoconstrictive fibers - a rec. (even 350 ml of the blood emptied into the systemic circulation during several minutes!)

# Splanchnic Circulation

- **Intestinal circulation**  
(*a. coeliaca, a. mesenterica superior and inferior*)
- submucous plexus, branches enter musculature and intestinal villi
- countercurrent exchange of substances



Ganong's Review of Medical Physiology, 23<sup>rd</sup> edition

# Splanchnic Circulation



- **Intestinal circulation**  
(*a. coeliaca, a. mesenterica superior and inferior*)
- Regulation of blood flow:
  - metabolic vasodilation (mediators: adenosine, ↓  $[K^+]$ <sub>e</sub> and ↑ osmolarity)
  - neural regulation – almost exclusively sympathetic,  $\alpha > \beta$  rec. → **vasoconstriction**

# Splanchnic Circulation

- **Hepatic circulation (*v. portae, a. hepatica*)**
  - 25% of the cardiac output (~1.5 l/min)
    - $\frac{3}{4} v. \text{ portae}, \frac{1}{4} a. \text{ hepatica}$ 
  - **portal circulation**
    - 2 capillary bloodstreams in series (intestinal villi, liver sinusoids)
    - $\downarrow O_2$  content  $\rightarrow a. \text{ hepatica}$  represents the nutritive hepatic circulation
- Regarding  $O_2$  supply, the ratio is opposite!

# Splanchnic Circulation


- Hepatic circulation (*v. portae, a. hepatica*)
- functional unit - acinus



Ganong's Review of  
Medical Physiology,  
23<sup>rd</sup> edition

# Splanchnic Circulation

- **Hepatic circulation (*v. portae*, *a. hepatica*)**
- pressures:
  - *a. hepatica*: 90 mmHg
  - *v. hepatica*: 5 mmHg
  - *v. portae*: 10 mmHg
  - sinusoids: 2.25 mmHg



Ganong's Review of  
Medical Physiology,  
23<sup>rd</sup> edition

# Splanchnic Circulation

- **Hepatic circulation (*v. portae, a. hepatica*)**
- inverse regulation of blood flow in *v. portae* and *a. hepatica*:
  - between meals: many sinusoids collapsed, **flow in *v. portae* low**, adenosine formed constantly and washed less → **dilation of terminal hepatic arterioles**
  - after a meal: **flow in *v. portae* ↑**, adenosine washed faster → **constriction of hepatic arterioles**, higher flow in *v. portae* opens so far collapsed sinusoids
- increased hepatic pressure (cirrhosis) → ascites

# Splanchnic Circulation

- **Hepatic circulation** (*v. portae, a. hepatica*)
- Regulation of blood flow:
  - **neural:** symp. vasoconstrictive fibers – α rec.
  - **metabolic:** adenosine → **vasodilation**
  - **passive:** ↑ BP → passive dilation of portal vein radicles → ↑ liver blood amount  
*congestive heart failure*  
*diffuse noradrenergic discharge due to ↓ BP*
- **sufficient O<sub>2</sub> supply is essential for liver function!** - ↓ flow → ↑ O<sub>2</sub> extraction


# Splanchnic Circulation

- **Hepatic circulation (*v. portae, a. hepatica*)**
- hepatic lymphatic circulation
  - formation of almost  $\frac{3}{4}$  of the body lymph
  - lymph rich on proteins (many plasmatic proteins are formed in hepatocytes + proteins from plasma due to the high permeability of sinusoids)

# Fetal Circulation


# Fetal Circulation

- placenta, umbilical vein
- liver, *ductus venosus*
- *crista dividens, foramen ovale*
- blood supply of the head and upper limbs
- *v. cava superior and inferior*
- the right ventricle
- *ductus arteriosus*
- aorta – the blood supply of the lower part of body + 60% of the cardiac output is directed to placenta



# Fetal Circulation


- fetal haemoglobin
- short-period hypoxia
- longer hypoxia
- thick muscle wall of umbilical vessels



Ganong's Review of Medical Physiology, 23<sup>rd</sup> edition


# Fetal Circulation

- **Changes after birth**
- **Closure of umbilical vein**
  - sudden ↑ of peripheral resistance and blood pressure
  - contraction of musculature of *ductus venosus* and its closure
- **The first inspiration** (due to asphyxia and cooling of the body)
  - ↓ resistance of the lung bloodstream
  - much more blood into lungs



# Fetal Circulation

- **Changes after birth**
- Decrease of pressure in right atrium and its increase in left atrium due to:
  - ↑ filling of left atrium by the blood from lungs
  - ↓ venous return to right atrium due to closure of umbilical vein
  - left ventricle works against ↑ pressure in aorta
- **Closure of *foramen ovale***
- **Closure of *ductus arteriosus***




# Renal Circulation

# Renal Circulation


- main functions of kidneys
- High filtration rate requires an adequate blood supply!
  - kidneys form only ~0.4 % of the body weight
  - blood flow 1.2 l/min, ~25% of cardiac output
- distribution of blood flow is **irregular**, the most flows through cortex (glomeruli – filtration)

# Renal Circulation




(a) Frontal section illustrating major blood vessels

© 2013 Pearson Education, Inc.



(b) Path of blood flow through renal blood vessels

<http://classes.midlandstech.edu/carterp/Courses/bio211/chap25/chap25.htm>



© 2013 Pearson Education, Inc.

<http://classes.midlandstech.edu/carterp/Courses/bio211/chap25/chap25.htm>


# Renal Circul

- v. aff., v. eff.

- glomerular blood flow = 
$$\frac{P_{v.a.} - P_{v.e.}}{R_{v.a.} + R_{v.e.} + \cancel{R_{g.k.}}}$$
- ↑ resistance in vas aff. or vas eff. → ↓ the renal blood flow (if the arterial pressure is stable)
- regulate the glomerular filtration pressure:

constriction of vas aff. → ↓ glomerular pressure → ↓ filtration

constriction of vas eff. → ↑ glomerular pressure → ↑ filtration



Guyton &  
Hall.  
Textbook  
of Medical  
Physiology

# Renal Circulation

- **Regulation of renal blood flow:**
  - 1) Myogenic autoregulation
  - 2) Neural regulation
  - 3) Humoral regulation

# Renal Circulation

- **Regulation of renal blood flow:**
  - 1) Myogenic autoregulation
    - dominates
    - provides stable renal filtration activity by maintaining stable blood flow at varying systemic blood pressure

# Renal Circulation

- **Regulation of renal blood flow:**

- 2) Neural regulation

- conformed to demands of systemic circulation
- **sympathetic system - norepinephrine**

light exertion/upright body posture → ↑ sympathetic tone → ↑ tone of v. aff. and eff. → ↓ renal blood flow but without ↓ GFR (↑ FF)

higher ↑ of sympathetic tone - during anesthesia and pain - GFR may already ↓

# Renal Circulation

- **Regulation of renal blood flow:**

## 3) Humoral Regulation

- contribute to regulation of systemic BP and regulation of body fluids
- **NE, E** (from the adrenal medulla)  
constriction of aff. and eff. arterioles → ↓ renal blood flow and GFR

(small impact with the exception of serious conditions, for example serious bleeding)

# Renal Circulation

- **Regulation of renal blood flow:**

## 3) Humoral Regulation

- contribute to regulation of systemic BP and regulation of body fluids
- **endothelin**
  - constriction of aff. and eff. arterioles → ↓ renal blood flow and GFR
  - released locally from the impaired endothel (physiological impact - hemostasis; pathologically increased levels at the toxemia of pregnancy, acute renal failure, chronic uremia)

# Renal Circulation

- **Regulation of renal blood flow:**

## 3) Humoral Regulation


- contribute to regulation of systemic BP and regulation of body fluids
- **NO**
  - continual basal production → vasodilation → stable renal blood flow and GFR
- **prostaglandins (PGE<sub>2</sub>, PGI<sub>2</sub>), bradykinin**
  - vasodilation
  - minor impact under physiological conditions
  - non-steroidal anti-inflammatory agents during stress!

# Renal Circulation

- **Regulation of renal blood flow:**

## 3) Humoral regulation

- contribute to regulation of systemic BP and regulation of body fluids
- **Renin-angiotensine system**



Ganong's Review of Medical Physiology, 23<sup>rd</sup> edition