

Regulation of Blood Flow

Assoc. Prof. MUDr. Markéta Bébarová, Ph.D.

Dept. of Physiology, Faculty of Medicine, Masaryk University



Definition of Blood Flow

mathematical formulation – analogy with the electric current

Ohm's law

$$I = U / R \longrightarrow$$

$$Q = \Delta P / R$$

Q blood flow

ΔP difference of pressure at the beginning and at the end of a vessel

R resistance of the vessel (peripheral resistance)

Definition of Blood Flow

$$Q = \Delta P / R$$

$$R = 8\eta l / \pi r^4$$



Poiseuille – Hagen formula

$$Q = \Delta P \cdot \pi r^4 / 8\eta l$$

r radius of the vessel

η viscosity of the blood

l length of the vessel

This formula applies to the steady laminar flow in a rigid tube!

Blood viscosity is not constant, *plasma skimming*, turbulent flow, elastic vessels!

Definition of Blood Flow


$$Q = \Delta P / R$$

$$R = 8\eta l / \pi r^4$$

r radius of the vessel
 η viscosity of the blood
l length of the vessel

Poiseuille – Hagen formula

$$Q = \Delta P \cdot \pi r^4 / 8\eta l$$




Methods for Measuring Blood Flow

- A. with a cannula inserted into a vessel
- B. without direct contact with the blood flow
 1. Electrical Induction Principle
 2. Doppler Effect
 3. Plethysmography
 4. Fick Principle

Methods for Measuring Blood Flow

1. Electrical Induction Principle

- ❖ the electromagnetic flowmeter



Guyton and Hall.
Textbook of Medical Physiology, 11th edition


- ❖ the generated electromotive force is proportional to the velocity of blood flow
- ❖ can detect changes in the velocity < 0.01 s \rightarrow recording of both steady blood flow and its pulsatile changes

Methods for Measuring Blood Flow

2. Doppler Effect

- ❖ the ultrasonic Doppler flowmeter; most common
 - ultrasonic waves of a known wave length (frequency)
 - waves reflect from the red and white blood cells → a change (↑) of the wave length (↓ frequency)
 - reflected waves are picked up by a sensor

- ❖ change of the wave length (frequency) is proportional to the velocity of blood flow




- ❖ both steady blood flow and its pulsatile changes can be measured

Methods for Measuring Blood Flow

3. Plethysmography


- ❖ usually as the venous occlusion plethysmography
- ❖ can be used on limbs
- venous drainage of the limb is stopped (e.g. with an arm cuff)
- increasing volume of the limb is lineary proportional to the arterial inflow of blood



Methods for Measuring Blood Flow

3. Plethysmography

- ❖ usually as the venous occlusion plethysmography
- ❖ can be used on limbs
- venous drainage of the limb is stopped (e.g. with an arm cuff)
- increasing volume of the limb is linearly proportional to the arterial inflow of blood



Methods for Measuring Blood Flow

4. Fick Principle - Direct Fick Method

$$Q = \frac{A / \text{time}}{AV \text{ diff}}$$

- blood flowing from the right heart to the lungs – about 150 ml O₂ / 1 l
- blood flowing from the lungs to the left heart – about 200 ml O₂ / 1 l




The blood catches 50 ml O₂ / 1 l during passage through the lungs.

- The total O₂ consumption is 250 ml / 1 min.



$$CO = \frac{250 \text{ ml O}_2 / \text{min}}{50 \text{ ml O}_2 / \text{l}} = 5 \text{ l / min}$$



Methods for Measuring Blood Flow

4. Fick Principle – Method of Indicatory Gas

- ❖ to determine the instantaneous blood flow through a specific tissue
- ❖ for example the cerebral or coronary blood flow using inhaled nitrous oxide N_2O – **Kety method**

$$\text{cerebral blood flow} = \frac{\text{N}_2\text{O removed from blood by brain / time}}{\text{averaged arteriovenous difference of N}_2\text{O}}$$

N₂O concentration in the venous blood
↓


Methods for Measuring Blood Flow

4. Fick Principle - Indicator Dilution Technique

- known amount of an indicator (dye or radioactive isotope) is injected into a peripheral (an arm) vein (A , [mg])
- concentration of the indicator in serial samples of the arterial blood is determined
- estimation of the averaged concentration of the indicator in the arterial blood after a single circulation (C , [mg/ml])

$$CO = \frac{A}{C (t_2 - t_1)} \quad \begin{matrix} [\text{mg}] \\ [\text{mg.ml}^{-1}.\text{s}] \end{matrix}$$

❖ thermodilution



Ganong's Review of Medical Physiology, 23rd edition.

Regulation of Blood Flow

$$Q = \Delta P \cdot \pi r^4 / 8 \eta l$$

Resting Tone

- ❖ tonic activity of vasoconstrictive sympathetic fibres
- ❖ a role might play also: myogenic response of vessels to the blood pressure (later), high concentration of O₂ in the arterial blood, Ca²⁺

Basal Tone


- ❖ in response to denervation; due to spontaneous depolarizations of the vascular smooth muscles

Regulation

Local

Systemic

B. Vasoconstriction and vasodilatation



Regulation of Blood Flow - Local

A. Acute

seconds to minutes, but incomplete (about $\frac{3}{4}$ of the desired effect)

1. Metabolic Autoregulation
2. Myogenic Autoregulation
3. Regulation Mediated by Endothelium

B. Chronic

hours, days to weeks , even months

Regulation of Blood Flow - Local

Metabolic Autoregulation

insufficient blood flow $\begin{cases} \nearrow \uparrow \text{ metabolic demands of a tissue} \\ \searrow \downarrow \text{ or stopped blood supply} \end{cases}$

→ \uparrow concentration of metabolites, \downarrow pH, \uparrow osmolarity in the interstitium, \uparrow tissue temperature; \downarrow pO₂, nutrients

→ **vasodilatation**

Preferred to the systemic regulation in case of hypoxia (to preserve the adequate tissue perfusion).

It plays the key role in e.g. brain, heart and skeletal muscles.

Regulation of Blood Flow - Local

Metabolic Autoregulation

active hyperemia

reactive hyperemia

Regulation of Blood Flow - Local

Myogenic Autoregulation (Bayliss effect)

↑ blood pressure

→ ↑ blood flow and ↑ tension in the vascular wall

$$Q = \Delta P / R$$

Law of Laplace

$$T = P \cdot r$$

→ mechanical stimulation, depolarization and subsequent contraction of the smooth muscle cells in the vascular wall → vasoconstriction

→ return of the blood flow back on the original level

It plays an important role in the brain and kidneys.

Regulation of Blood Flow - Local


Regulation Mediated by Endothelium

endothelial-derived relaxing factor (EDRF) – NO

→ vasodilatation

- ❖ synthesized in the **endothelial cells** of arteriols and small arteries due to the **shear stress** induced by the flowing blood
- ❖ **synthesis stimulated by** the products of thrombocyte aggregation and also by many primary vasoconstrictive substances


B. Vasoconstriction and vasodilatation



Regulation of Blood Flow - Local

Regulation Mediated by Endothelium

endothelial-derived relaxing factor (EDRF) – NO



Regulation of Blood Flow - Local

Regulation Mediated by Endothelium

prostacyclin

- ❖ synthesized in the endothelial cells from the arachidonic acid
- ❖ inhibition of thrombocyte aggregation and **vasodilation**

thromboxane A_2

- ❖ synthesized from the arachidonic acid by thrombocytes
- ❖ support of thrombocyte aggregation and **vasoconstriction**

A balance between them is crucial for formation of the localized clot and preservation of the blood flow.

Regulation of Blood Flow - Local

Regulation Mediated by Endothelium

endothelins

- ❖ polypeptides synthesized by endothelial cells (ET-1, ET-2, ET-3)
- ❖ 2 endothelin **receptors**:
 - ET_A – specific for ET-1, in many tissue vessels, → **vasoconstriction**
 - ET_B – ET-1 to ET-3, function?
- ❖ **ET-1 – one of the most potent vasoconstrictive substances**
- ❖ the exact physiological role not known
- ❖ restricts bleeding, play a role in closing *ductus arteriosus* at birth

Regulation of Blood Flow - Local

Serotonin (5-OH tryptamine)

❖ vasoconstrictive effect

- in a damaged tissue
- direct local effect
- released from thrombocytes

❖ vasodilatory effect

- in an undamaged tissue
- through increased activity of NO synthase

Regulation of Blood Flow - Local

A. Acute

seconds to minutes, but incomplete (about $\frac{3}{4}$ of the desired effect)


1. Metabolic Autoregulation
2. Myogenic Autoregulation
3. Regulation Mediated by Endothelium

B. Chronic

hours, days to weeks , even months

Regulation of Blood Flow - Local

Chronic regulation



Guyton and Hall.
Textbook of Medical
Physiology, 11th edition

Especially important in case of the long-term change of metabolic demands of a tissue - to provide sufficient blood flow without circulation overload.

Regulation of Blood Flow - Local


Chronic regulation

- ❖ mediated by changes of the tissue vascularity
- ❖ the key role – lack of O_2 , also nutrients
- ❖ **angiogenic or vascular growth factors** - small peptides, best characterized: vascular endothelial growth factor (VEGF), fibroblast growth factor, and angiogenin
- ❖ fast in young individuals and in newly formed tissue

Regulation of Blood Flow - Local

Chronic regulation

unstimulated muscle



regularly stimulated muscle

