

BLOOD AND HEMATOPOIESIS


Petr Vaňhara

Department of Histology and Embryology
LF MU

BLOOD

Blood is body fluid

- transport medium (O_2 , CO_2 , metabolites, hormones, nutrients...)
- homeostasis of inner body environment (thermoregulation, acidobasic equilibrium, oncotic pressure)
- integrity of cardiovascular system (clotting cascade)
- immune reactions



formed blood elements


- erythrocytes
- leukocytes
- thrombocytes

plasma

- ions, proteins, low mass organic compounds
- fluid ECM





Blood can be considered as a specialized connective tissue



BLOOD PLAMA AND TISSUE FLUID

plasma

- 2,8-3,5 l
- pH 7.4 (± 0.05)
- ~ 92% **water**
- ~ 1% **ions** (Na^+ , K^+ , Ca^+ , Mg^+ , Cl^- , HCO_3^-), **low mass organic compounds** (glucose, aminoacids, cholesterol, lipids, waste products of metabolism), **respiration gases**
- ~ 7% **proteins** (albumins, globulins, fibrinogen)



IONS AND LOW MASS MOLECULES OF BLOOD PLASMA (~1%)

- ~ 1% **ions** (Na^+ , K^+ , Ca^+ , Mg^+ , Cl^- , HCO_3^-), **low mass organic compounds** (glucose, aminoacids, cholesterol, lipids, waste products of metabolism), **respiration gases**

Cations	Sodium	136–148 mmol/l	Osmotic pressure, volume, pH
	Potassium	3,7–5,0 mmol/l	Membrane potential of cells (nerve, muscle)
	Calcium	2,15–2,61 mmol/l	Permeability of membranes, blood clotting, neuromuscular junctions
	Magnesium	0,66–0,94 mmol/l	Cofactor of enzymes, neuronal conduction
	Iron ♂	12–27 $\mu\text{mol/l}$	Cofactor of enzymes, in hem of hemoglobin
	Iron ♀	10–24 $\mu\text{mol/l}$	
	Copper	12–22 $\mu\text{mol/l}$	Cofactor of enzymes
Anions	Chlorides	95–110 mmol/l	Osmotic pressure, volume, pH
	Bicarbonates $[\text{HCO}_3^-]$	22–26 mmol/l	Transport of CO_2 , buffer - pH
	P_i	0,6–1,4 mmol/l	Buffer - pH
	Iodide	276–630 $\mu\text{mol/l}$	Hormones of thyroid gland

Memorizing of this table is not necessary for completing our course 😊

IONS AND LOW MASS MOLECULES OF BLOOD PLASMA (~1%)

- ~ 1% **ions** (Na^+ , K^+ , Ca^+ , Mg^+ , Cl^- , HCO_3^-), **low mass organic compounds** (glucose, aminoacids, cholesterol, lipids, waste products of metabolism), **respiration gases**

Glucose	3,3–6,1 mmol/l
Aminoacids	2,3–3,9 mmol/l
Urea	3,0–7,6 mmol/l
Lipids	4–9 g/l
Triacylglycerols	0,5–1,8 mmol/l
Phospholipids	1,8–2,5 g/l
Creatinine	55–110 $\mu\text{mol/l}$
Cholesterol (total)	3,5–5,2 mmol/l
Bilirubin	3,3–18,0 $\mu\text{mol/l}$
Lactate	0,55–2,22 mmol/l


But what is important to know is that:



COMPOSITION OF BLOOD PLASMA IS CONSTANT
and regulated in narrow range → essential for **clinical medicine**


PROTEINS OF BLOOD PLASMA (7%)

- oncotic blood pressure
- transport
- coagulation
- immune response
- regulatory proteins



PROTEINS OF BLOOD PLASMA (7%)

- prealbumin
- transport
- albumin
- 68kDa
- transport
- osmotic pressure
- α_1 region
- α_1 lipoprotein (HDL)
- α_1 acid glycoprotein
- α_1 antitrypsin
- (α_1 fetoprotein)
- α_2 region
- α_2 macroglobulin
- haptoglobin
- β_1 region
- transferrin
- hemopexin
- β lipoprotein (LDL)
- C4 (complement)
- β_2 region
- CRP
- fibrinogen
- β_2 microglobulin
- C3 (complement)
- γ region
- IgA, IgG, IgM



Why it is useful? [Click here.](#)




Electrophoretic separation of serum proteins


- A. Electrophoretogram of normal serum on cellulose acetate strip
- B. Densitometric scanning from cellulose acetate strip converts bands to characteristic peaks of albumin, α_1 -globulin, α_2 -globulin, β -globulin and γ -globulin


Reference ranges for blood tests


sorted by mass (upper) and molarity (lower diagram)

- Sources differ in the faded interval
- Limit is 0 or not available











BLOOD PLASMA AND SERUM

- serum ≠ plasma



FORMED BLOOD ELEMENTS



HEMATOCRIT

Ratio of erythrocyte mass volume to volume of full blood

Erythrocytes
 $4.2\text{-}6.2 \times 10^6/\mu\text{l}$

HEMATOCRIT



47±5%




42±4%

Norm





Anemia

Polycythemia



ERYTHROCYTES


Size depends on osmotic pressure of environment



ERYTHROCYTES



Deviations from normal size

- **anisocytosis**
 - macrocytes ($>9 \mu\text{m}$)
 - microcytes ($<6 \mu\text{m}$)



ERYTHROCYTES




Erythrocyte is amazingly flexible cell



ERYTHROCYTES

Shape of erythrocytes

- **integral proteins**
 - band 3, glycoprotein A (ion transporters)
- **spectrin**
- **ankyrin**
- **actin and actin-associated proteins**
 - tropomodulin, tropomyosin
- **hemoglobin**





Courtesy of Dr. Paul Biering, Tufts University

ERYTHROCYTES

Deviations from biconcave shape


- **poikilocytosis**
- **acantocytes** (irregular spikes)
- **codocytes** („tyre“)
- **echinocytes** (spiked membrane)
- **eliptocytes** (elliptic)
- **spherocytes** (spheroidal)
- **stomatocytes** (some parts missing or other irregularities)
- **drepanocytes** (sickle)
- **dacrocytes** (tear drop)




ERYTHROCYTES

Deviations from biconcave shape

Normal




Codocyte





(b)

Spherocyte




(d)



Echinocyte


ERYTHROCYTES

Hereditary elliptocytosis




ERYTHROCYTES

Hereditary elliptocytosis












Hereditary spherocytosis



ERYTHROCYTES

Sickle cell anemia


- Abnormal hemoglobin (hemoglobin S)





ERYTHROCYTES

Sickle cell anemia

- pathological genotype (heterozygote HbS/HbA) is beneficial





Malaria



ERYTHROCYTES


- Erythrocytes lack nucleus and other organelles
- Anaerobic glycolysis
- Hemoglobin




Heme




Hemoglobin



Erythrocyte






ERYTHROCYTES





ERYTHROCYTES

- Life span 120 days
- Constant abrasion
- No regeneration
- Removal of aged or damaged erythrocytes in spleen




LEUKOCYTES

- immune response
- morphological classification – **cytoplasmic granules**
(does not follow hematopoiesis)






GRANULOCYTES

- Lysosomes (primary, azurophilic, nonspecific granules)
- Specific (secondary) granules
- Polymorphic nucleus
- Terminally differentiated
- Short lifespan (hours)
- Reduced ER, GA, mitochondria (anaerobic glycolysis)
- Apoptosis





NEUTROPHILIC GRANULOCYTES

- **Neutrophils**
 - 50-70% of leukocytes in circulation
 - $\varnothing > 12 \mu\text{m}$
 - Segmented nucleus
 - Barr's body in females
 - **Azurophilic (primary) granules**
 - myeloperoxidase, lysozyme, proteases, defensins
 - **Neutrophilic (secondary) granules**
 - collagenase, bactericidal enzymes
 - Chemotaxis of other leukocytes
 - Microphages
-
- **Neutrophilic band**
 - **Neutrophilic segment**




NEUTROPHILIC GRANULOCYTES






NEUTROPHILIC GRANULOCYTES

- **Extravasation (diapedesis)**





Leukocyte
Extravasation






EOSINOPHILIC GRANULOCYTES

- **Eosinophils**
- 1-4% of leukocytes in circulation
- \varnothing 12-15 μm
- Irregular, characteristic bi-segmented nucleus
- **Azurophilic (primary) granules**
 - myeloperoxidase, lysozyme, proteases, defensins
- **Eosinophilic (secondary) granules**
 - bright red (eosinophilic)
 - major acidic protein
 - peroxidase
 - cytokines, chemokines
- Chemotaxis of other leukocytes
- Phagocytosis of antibody-antigen complexes
- Parasitic infections, allergic reaction
- Chronic inflammation



BASOPHILIC GRANULOCYTES


- **Basophils**
 - <1% of leukocytes in circulation
 - $\varnothing 12 \mu\text{m}$
 - Irregular, bisegmented nucleus, masked by granules
- **Azurophilic (primary) granules**
 - myeloperoxidase, lysozyme, proteases, defensins
- **Basophilic (secondary) granules**
 - $0.5 \mu\text{m}$
 - large, dark (basophilic)
 - heparin, histamin - vasodilatation
 - phospholipase A
- Analogs of mast cells
- Receptors for IgE
- Allergy, anaphylaxis, inflammation





AGRANULOCYTES

- Lysosomes only (azurophilic, nonspecific granules)
- Specific granules absent
- Non-segmented nucleus

Lymphocytes



Monocytes




LYMPHOCYTES

Lymphocytes


CD8 T Cell



B Cell



Regulatory T Cell






CD4 T Cell






LYMPHOCYTES

- Plasma cells



MONOCYTES

- Ø 12-15 µm
- Circulating precursors of macrophages, osteoclasts, microglia, Kupfer cells and dendritic cells
- Mononuclear phagocytic system
- Large, oval (bean, kidney) nucleus with less condensed chromatin and 2-3 nucleoli
- Basophilic cytoplasm
- Azurophilic granules



https://www.youtube.com/watch?v=R279Cs1_ltg

THROMBOCYTES

- Cell fragments without nucleus
- \varnothing 2-3 μm , discoid shape
- hyalomere, granulomere
- $150-400 \times 10^3/\mu\text{l}$
- blood clotting, repair of vessel wall

α -granules
300-500 nm





fibrinogen, PDGF

δ -granules
250-300 nm

serotonin, Ca^{++}
pyrophosphate
ADP, ATP

λ -granules
175-200 nm

lysosomal enzymes



THROMBOCYTES

1. Primary aggregation of platelets



- collagen fibers exposed by endothelial rupture
- platelet clot

2. Secondary aggregation of platelets

- clotting factors, ADP from thrombocytes attracts other platelets – **white thrombus**

3. Coagulation – blood clotting

- fibrin mesh capturing erythrocytes – **red thrombus**



4. Thrombus retraction


- contraction of thrombus (platelet actin and myosin)

5. Thrombolysis

- dissolving of thrombus (plasmin) and tissue regeneration



THROMBOCYTES



You needn't to know the clotting cascade in full details for our course, leave it for biochemistry



DIFFERENTIAL WHITE BLOOD CELL COUNT

THIS SLIDE IS REALLY IMPORTANT

Neutrophil band 4 %

segment 67 %

Norm

1:17

shift to the left

more bands

shift to the right

more segments

Eosinophils 3 %

Basophils 1 %

Lymphocytes 20 %

Monocytes 5 %

$$\sum = 100 \%$$

DIFFERENTIAL WHITE BLOOD CELL COUNT

Deviations from norm

	↑ Increased	↓ Decreased
Neutrophils	neutrophil granulocytosis	neutrophil granulocytopenia
Eosinophils	eosinophil granulocytosis	eosinophil granulocytopenia
Basophils	basophil granulocytosis	basophil granulocytopenia
Lymphocytes	lymphocytosis	lymphocytopenia
Monocytes	monocytosis	monocytopenia


DIFFERENTIAL WHITE BLOOD CELL COUNT

Example of population variability

Neutrophils	bands	0-5 %
	segments	35-85 %
Eosinopils		0-4 %
Basophils		0-1 %
Lymphocytes		20-50 %
Monocytes		2-6 %

According to: Haferlach et al. Kapesní atlas hematologie. Grada 2014

DIFFERENTIAL WHITE BLOOD CELL COUNT




DIFFERENTIAL WHITE BLOOD CELL COUNT

Age dependence

Age	Leukocytes ($\times 10^3$)	Neutrophils (%)	Lymphocytes (%)	Monocytes (%)	Eosinophils (%)
Birth	18	61	31	6	2
1 week	12.2	45	41	9	4
1 mo	10.8	35	56	7	3
6 mo	11.9	32	61	5	3
1 yr	11.4	31	61	5	3
4 yr	9.1	42	50	5	3
10 yr	8.1	54	38	4	2
16 yr	7.8	57	35	4	3


WBC, White blood cell.

HEMATOPOIESIS






ERYTHROPOIESIS

- **2×10^{11} of new erythrocytes daily**
- **proerythroblast** (~14-19 μm)
 - mitotically active
 - dominant, round nucleus with 1-2 nucleoli
 - mildly basophilic cytoplasm
- **basophilic erythroblast** (~13-16 μm)
 - mitotically active
 - heterochromatic nucleus with inconspicuous nucleoli
 - basophilic cytoplasm (sometimes more than in proerythroblast)
- **polychromatophilic erythroblast** (~13-16 μm)
 - mitotically active
 - **production of hemoglobin**
 - blue-gray cytoplasm due to combined basophilic (polyribosomes) and acidophilic aspects (hemoglobin)
 - heterochromatic nucleus (checkerboard appearance)
- **orthochromatophilic erythroblast** (~8-10 μm)
 - mitotically inactive
 - small, compact, eccentric, pyknotic nucleus → **extrusion**
 - mildly acidophilic cytoplasm with basophilic residues
- **reticulocyte** (polychromatophilic erythrocyte, ~ 7-8 μm)
 - **lacks nucleus, still spheroid shape**
 - acidophilic cytoplasm
 - *substantia reticulofilamentosa* visible by supravital staining (brilliant cresyl blue)
- **erythrocyte** (~7-8 μm)
 - anucleate, biconcave disc
 - acidophilic cytoplasm



ERYTHROPOIESIS



GRANULOPOIESIS

- **myeloblast** (~15 µm)
 - mitotically active
 - round-oval, euchromatic nucleus
 - 2-6 apparent nucleoli
 - weakly basophilic cytoplasm without granules
- **promyelocyte** (~15-24 µm)
 - mitotically active
 - round-oval nucleus with partly condensed chromatin
 - basophilic cytoplasm with azurophilic granules
- neutrophilic, eosinophilic or basophilic **myelocyte** (~10-16 µm)
 - mitotically active
 - oval or bean-shaped nucleus with condensed chromatin
 - increasing number of specific granules in cytoplasm
- neutrophilic, eosinophilic or basophilic **metamyelocyte** (~10-12 µm)
 - mitotically inactive
 - horseshoe-like nucleus with condensed chromatin
- neutrophilic, eosinophilic or basophilic **granulocyte** (~10-12 µm)
 - segmentation of nucleus
 - cytoplasm rich in specific and azurophilic granules

myeloblast




promyelocyte



myelocyte



metamyelocyte



granulocyte - band




granulocyte - segment



GRANULOPOIESIS

myeloblast



polychromatophilic erythroblast

promyelocyte

neutrophilic segment

neutrophilic
metamyelocyte

eosinophilic
myelocyte

neutrophilic myelocyte

promyelocyte

neutrophilic myelocyte

promyelocyte




myeloblast

promyelocyte

metamyelocyte




myelocyte

granulocyte



THROMBOPOIESIS

- **megakaryoblast** (up to 30 µm)
 - large oval, nonlobed nucleus with prominent nucleoli
 - basophilic cytoplasm
 - successive endomitoses without karyokinesis and cytokinesis
- **promegakaryocyte** (up to 100 µm)
 - large cell with polyploid nucleus (8n-64n)
- **megakaryocyte** (80-150 µm)
 - polyploid, multilobed nucleus (8n-64n)
 - azurophilic and platelet granules
 - multiple centrioles, ER and Golgi apparatus
 - numerous peripheral invaginations of plasma membrane—platelet demarcation channels defining individual thrombocytes
 - release of **thrombocytes** into bone marrow sinusoids



MONOCYTOPOIESIS AND LYMPHOPOIESIS


MONOCYTOPOIESIS

- **monoblast** (~16 µm)
 - round, bean shaped nucleus with 2-6 nucleoli
 - mildly basophilic cytoplasm
- **promonocyte** (~16-20 µm)
 - mitotically active (1-2 divisions)
 - large nucleus with mild indentation, unapparent nucleoli
 - basophilic cytoplasm
 - azurophilic granules
- **monocyte**
 - short-time in circulation, then extravasation and differentiation to tissue macrophages

LYMPHOPOIESIS

- **lymphoblast** (~18-20 µm)
 - round-oval nucleus with several nucleoli
 - mildly-basophilic cytoplasm without azurophilic granules
- **prolymphocyte** (~12-15 µm)
 - morphological transition and maturation to lymphocytes
- **lymphocyte**
 - further maturation and differentiation outside bone marrow

OVERVIEW OF ADULT HEMATOPOIESIS





HEMATOPOIETIC STEM CELLS AND PROGENITORS

- **Hematopoietic stem cell**

- Quiescent, slow cell cycle
- Transmembrane phosphoglycoprotein CD34⁺ - adhesion within niche
- No expression of lineage surface markers (Lineage negative or Lin⁻)
- Transplantations




- **Colony/Burst – Forming Unit – CFU/BFU**

- Progenitors of individual lines
- Colonies in vitro




EMBRYONIC HEMATOPOIESIS

- **Extraembryonic mesoblastic period (day 16-20 – week 8)**
 - yolk sac
 - classical model – hemangioblasts (bipotent cells)
 - large, nucleated erythroid cells
- **aorta-gonad-mesonephros (day 28 – week 4)**
- **hepatolienal period (month 1 – birth)**
 - colonization of fetal liver and spleen
- **medullary period (month 4-6. – rest of life)**
 - bone marrow




HEMATOPOIETIC ISLANDS



- hepatolienal and bone marrow hematopoiesis
- erythroblast islands



A





B





- Pro-erythroblast
- Basophilic erythroblast
- Polychromatic erythroblast
- Orthochromatic erythroblast
- Erythrocyte
- Pyknotic nucleus
- Central macrophage

INTRAEMBRYONIC HEMATOPOIESIS

- **Aorta-gonad-mesonephros (day 28 – week 4)**
 - para-aortic clusters in mesoderm of splanchnopleura
 - source of embryonic HSCs




- **Placenta**



SUMMARY OF HEMATOPOIESIS

Embryonic

- yolk sac
- AGM
- liver and spleen
- bone marrow



Adult

- bone marrow (yellow, red)
- extramedullar hematopoiesis rare (pathology)

Thank you for attention



pvanhara@med.muni.cz
www.histology.med.muni.cz