

**Topic P09: Diagnostics of spirochetal infections**

To study: *Borrelia, Leptospira, Treponema* (from textbooks, WWW et  
 From spring term: Microscopy, PCR, methods of detection of antibodies

**Lyme borreliosis**

Common table for Task 1, 2 and 3.

Patient Letter	Short clinical description (1-3 words characterizing the situation)	ELISA (Task 1)				W. blotting (T2)		PCR (T3) (+/-)	Conclusion: final interpretation, recommendation for event. therapy
		IgM		IgG		IgM (+/-)	IgG (+/-)		
		Abs.	(+/-)	Abs.	(+/-)				
J									
K									
L									
M									
N									

**Task 1: Proof of antibodies to *Borrelia garinii* using ELISA**

Read the results of patients with suspect Lyme borreliosis. Both IgG and IgM antibodies are assessed. In A1 field (corresponding to A1 well of the microtitration plate) you can see CAL level (borderline level; all absorbance levels above CAL are positive, all absorbance levels beneath CAL are negative). There are controls in B1 and C1. Patients J to N are in fields with coloured squares.

Write CAL level here, check, whether negative control is really negative and positive control really positive. Then read and interpret ELISA results for patients J, K, L, M, N (do not write them here, use main table above).

CAL level (well A1):		K+ absorbance level (well B1):		<input type="checkbox"/> K+ is OK <input type="checkbox"/> K+ is not OK	← tick what is correct
<b>IgM</b>		K- absorbance level (well C1):		<input type="checkbox"/> K- is OK <input type="checkbox"/> K- is not OK	
CAL level (well A1):		K+ absorbance level (well B1):		<input type="checkbox"/> K+ is OK <input type="checkbox"/> K+ is not OK	← tick what is correct
<b>IgG</b>		K- absorbance level (well C1):		<input type="checkbox"/> K- is OK <input type="checkbox"/> K- is not OK	

**Task 2: Proof of antibodies to *Borrelia garinii* using Western blotting**

In patients diagnosed in the task No.1, the serum samples or CSF were performed by Western blotting. Read results according to instructions. Use the given pattern for evaluation of the reaction. A diagnostic scheme is always the same – ELISA is used for screening, whereas Western blotting is performed as a confirmation of ELISA results. Read the Western blot results of patients J to N and write the results to the main table.

**Task 3: Diagnostics of Lyme borreliosis using polymerase chain reaction (PCR)**

According to the given photos of PCR product on the agarose gel, draw and record which of the tested samples is positive. Remark, that with regard to anamnesis, PCR reaction was performed only in two of our five patients. After that, interpret finally the total of all three examinations and write down a conclusion.

**Syphilis**

**Task 4: Direct proof of syphilis**

Direct proof of syphilis is only possible if suitable samples are sent to the laboratory. In some stages of the disease it is not possible to take anything for this purpose.



**a) Rabbit infectivity testing – RIT**

Write down the name of the rabbit used for the test.

(It is derived from these islands: →→→→→→→→.)

Exsudate from a suspect ulcer uses to be evaluated by darkfield microscopy and inoculated to rabbits testes. The tested animal starts to suffer from orchitis after 10 days after inoculation. Rabbit name:



**b) Darkfield microscopy**

Look at the microphotography of treponemas taken from a darkfield microscope, draw the principle of darkfield microscopy, and also record your observation.

**c) Direct immunofluorescence**

Look at the microphotography of treponemas taken from a fluorescent microscope and record your observation.

4b) principle	4b) result	4c)
---------------	------------	-----

The causative agent of syphilis, *Treponema pallidum*, is NOT a culturable microorganism. The diagnostics is dependent on a stage of disease.

**Indirect diagnostics of syphilis**

Common table for Task 5 and 6.

Patient Letter	Short clinical characterization	Task 5 screening		Task 6 confirmation				Conclusion: final interpretation, eventually recommended therapy	
		RRR	MHA-TP	ELISA		WB			
				FTA-ABS	IgM Absorbance (+/-)	IgG Absorbance (+/-)	IgM (+/-)		IgG (+/-)
A									
B									
C									
D									
E									

**Task 5: Screening of syphilis – RRR and MHA-TP**

Pregnant women and blood donors undergo a screening using rapid reagin reaction (RRR) and *Treponema pallidum* microhemagglutination (MHA-TP). Read results of screening in a given group of persons and assess who of them need to be tested using confirmation tests. Record your results directly into the table.

RRR: flocculation in the well is positive; MHA-TP: agglutinate formation positive (see practical J07).

**Task 6: Confirmation of syphilis – FTA-ABS, ELISA and Western blotting**

Evaluate the results of FTA-ABS, ELISA a western blotting (WB) in patients who are suspect of syphilis (see the previous task). In ELISA reaction, count the cut-off and compare K-, + and patient values with it.


A1 field (A1 well) represents the blank.

Cut off level (C1 + D1) / 2		K- absorbance level (B1 value):	<input type="checkbox"/> K- is OK <input type="checkbox"/> K- is not OK	← tick what is correct
<b>IgM</b>		K+ absorbance level (E1 value):	<input type="checkbox"/> K+ is OK <input type="checkbox"/> K+ is not OK	
Cut off level (C1 + D1) / 2		K- absorbance level (B1 value):	<input type="checkbox"/> K- is OK <input type="checkbox"/> K- is not OK	← tick what is correct
<b>IgG</b>		K+ absorbance level (E1 value):	<input type="checkbox"/> K+ is OK <input type="checkbox"/> K+ is not OK	

## Leptospirosis

### Task 7: Direct proof of *Leptospira* sp.

According to a given picture, describe and draw morphology of leptospires cultivated in the liquid Korthoff's medium for 2 weeks. Urine of a patient who is suspect of leptospirosis was used for the test.



### Check-up questions:

1. What is the basic limitation in diagnostics of spirochetal infections in comparison with common bacterial infections?
2. A patient had a tick two days in his fossa poplitea. The tick was extracted and the doctor decided to take blood "for *Borrelia*". What type of diagnostics was meant by this and what finding can be supposed in this case?
3. Why materials with findings of antiborrelial antibodies are usually re-examined by Western blotting?
4. What are possible specimens for direct diagnostics of syphilis?
5. Does have a sense to use PCR method for testing seronegative patients (not having anti-borrelial antibodies)?
6. Do we perform antibiotic susceptibility testing in agents of spirochetal infections?
7. In a pregnant woman, RRR was found to be positive. What would be the interpretation, eventually what more tests would be required?
8. Should be syphilitics at control testing allways re-tested by all serological tests?
9. What reaction is the best to assess the activity of syphilis?