

Institute of Microbiology shows:




## TRACING THE CRIMINAL

Part three: More G+ criminals

# Survey of G+ bacteria

Story	Shape	In police evidence written as
<i>P01</i>	Medically important Cocci	Staphylococci ( <i>S. aureus</i> , CONS)
<i>P02</i>		Streptococci (viridating, haemolytical)
1.		Enterococci ( <i>E. faecalis</i> , <i>E. faecium</i> )
2.	Medically important Rods	Listeriae ( <i>L. monocytogenes</i> )
3.		Corynebacteria <a href="http://web.fccj.org">http://web.fccj.org</a>
4.		Bacilli
		<a href="http://vietsciences.free.fr">http://vietsciences.free.fr</a>



*Listeriae a coryneforms do not sporulate, unlike Bacilli*

# Contents

Clinical characteristics – enterococci

Clinical characteristics – G+ rods

Enterococci and G+ rods: therapy

Diagnostics of enterococci and G+ rods (+ pictures)


Differential diagnostics of enterococci and G+ rods

# Clinical characteristics - enterococci

# Story 1

- Lucy has problems with urination. Doctor prescribed Zinnat, but problems did not change. On the next visit, he let Lucy to urinate in a glass and sent her urine sample to microbiology. But the specimen could not be examined: urine was contaminated. Finally, it was possible to take urine aseptically and to change the therapy.


# Criminal No 1



<http://www.lbl.gov>

- *Enterococcus faecalis*
- As the „entero-“ in his name tells us, it is a bug **normally present in the intestine**. Nevertheless, it is also a **common UTI pathogen**.
- The doctor is guilty, too – prescribed antibiotics before knowing microbial susceptibility. **Enterococci are resistant to all cephalosporin atb**. And he did not perform aseptic urine sampling in the first phase.
- *UTI (urinary tract infections) are mostly bacterial, and many pathogens are primarily or secondarily resistant to some atb-s. So bacteriological examination of urine is recommended, although often not done in practice.*

# More about enterococci



- There are **tens of species** of them today
- All of them may be found
  - **in stool** (as a normal flora)
  - in the **urinary bladder** (as pathogen)
  - in the **vagina** (both symptomatically and asymptotically)
  - sometimes in **other sites** (wounds, bloodstream)
- Among two most common species, *E. faecalis* is slightly more often a pathogen, *E. faecium* is more often part of normal intestinal flora
- **Vancomycin-resistant enterococci (VRE)** are dangerous
- One of Enterococci, found in Brno, is named *Enterococcus moraviensis*

Clinical  
characteristics –  
G+ rods



# Story 2

- European commission had to solve one more problem. French agriculturiers protested againsts several EU-members, that do not want to import some **delicatess French cheese specialities** to their area.
- German officials stated, that one pregnant woman, Mrs. Hildegarda Messerschmidt, after having eaten the cheese had **elevated body temperature** and after delivery, her baby suffered **newborn meningitis** that needed prolonged and complicated treatment


<http://www.leighday.co.uk>



<http://womansday.ninemsn.com.au>

# Criminal No 2

<http://www.leighday.co.uk>



- *Listeria monocytogenes* is a G + rod, able to grow at low temperatures and high NaCl concentrations, typically in cheese storehouses. Nevertheless, people may get infected also by eating other food (salads, sausages, delicatesses, not properly cleaned vegetables)

# Listeria – more info

- In adults, symptomatic infections are rare. In pregnant women there is a risk of congenital infection of the fetus through placenta with abortus or fetus infection (in the last three months) as a result, or also perinatal infection during delivery (infection by contaminated vaginal secretion). In newborns, meningitis or septicemia are common
- Infection is not very frequent, but it has high lethality (% of dead people of all infected people)
- Sometimes it is used as a pretence for import limitation – it should be always decided individually according to real risk in a specific case

# Classification des aliments selon le risque lié à *Listeria monocytogenes*




<http://www.territoire-belfort.gouv.fr>

# Stroy




- Mr. Ulcerous, chronic diabetic, treated for ulcerations on his legs. He is often infected by various pathogens. What will be the newest one?

## Criminal No 3



- *Corynebacterium jejkeium*, the worst among non-diphtheria coryneforms. Its name is derived from „corynebakterium group JK“.
- Corynebacteria are grampositive rods club-shaped (*koryné* = club), sometimes pleomorphic.
- In the same genus, we have also diphtheria causative agent, rare in Europe, because of vaccination – *C. diphtheriae*.

# Diphtheria



Diphtheria - notice the pseudomembrane in the posterior pharynx. It can become very large and may obstruct the airway.



# More about non-diphtherical corynebacteria

- Part of **normal flora of skin**, together with staphylococci and yeast. Pathogens in woundsIn microscopy, they form „**palisades**“ – like the early medieval wooden fortifications


File:St Fagans Celtic village palisade.jpg, From Wikipedia, the free encyclopedia, available at [http://en.wikipedia.org/wiki/File:St\\_Fagans\\_Celtic\\_village\\_palisade.jpg](http://en.wikipedia.org/wiki/File:St_Fagans_Celtic_village_palisade.jpg), visited 2012-10-02



# What are „coryneform rods“

- „Coryneform rods“ (eventually „diphtheroids“) are various rods that share similar morphology (although size of rods may vary considerably).
- All of them are rare causative agents of various human infections.
- *Arcanobacterium haemolyticum* is a rare causative agent of pharyngitis
- Other genera: *Dermatophilus*, *Rhodococcus*\*, *Turicella* etc.
- Similar is also *Erysipelothrix rhusiopathiae* – causative agent of a zoonosis (erysipeloid)


*Rhodococcus jostii* was found on the body of the Moravian Marques and uncrowned Roman Emperor Jodocus (Jošt), that died 1411. The body is buried in St. Thomas church in Brno.



## Story 4

- Nurse Eileen was shocked: microbiology examination of ward furniture, week ago taken by hospital epidemiologists, were released, and in halfht of them contained some bacteria, even **BACILLI!** Yes, it is here – *Bacillus* sp. Poor Eileen was worrying all the night about it and slept badly. In the morning, she asked microbiologists, what type of bacterium the „*Bacillus*“ is...

# And she was very glad:





- it is no criminal! Usual members of *Bacillus* genus are harmless microbes from external environment. When found in clinical material, it is usually a contamination. So, the finding was not a problem – problem would be only when a *Bacillus* would be found from a site that is supposed to be sterile.

# But some Bacilli are important

- *Bacillus anthracis* causes a veterinary disease – anthrax. It was one of first diseases where vaccination was attempted (already by Louis Pasteur). Its spores are abusable for biological war or bioterrorism (about a case of leak of anthrax spores in the Soviet Union in 1979 see: [http://en.wikipedia.org/wiki/Sverdlovsk\\_anthrax\\_leak](http://en.wikipedia.org/wiki/Sverdlovsk_anthrax_leak))
- *Bacillus cereus* is causative agent of intoxications coming from cereals
- *Geobacillus* (formerly *Bacillus*) *stearothermophilus* & *Bacillus subtilis* are able to survive hot temperatures → we use them as control organisms for hot air and steam sterilisers.

# *Bacillus* and its spores



[www.cropsoil.uga.edu](http://www.cropsoil.uga.edu)

Spores of *Bacillus* sp. are sometimes larger than the vegetative cell, sometimes not; sometimes they are terminal, in other species they are subterminal or central


Enterococci and  
G+ rods: therapy

# Therapy of infections caused by enterococci and G+ rods

- No cephalosporins to fight enterococci and listeriae! In *E. faecalis*, ampicillin, is good, but in *E. faecium* there is a primary resistance. More atb's are co-trimoxazol, doxycylin, and, as a reserve, vankomycin. In haematooncological patients we can find epidemiologically serious vancomycin resistant strains – VRE. In such strains, only a new atb – linezolid – is effective



# Antibiotics used for enterococci and G+ rods



- Enterococci are tested on MH. G+ rods are tested on MH + red blood cells.

Antibiotic	Abbr.	Reference zone
Ampicilin (aminopenicilin)	AMP	17 mm
Co-amoxicilin (aminopnc*)	AMC	18 mm
Co-trimoxazol (mixt. 2)	SXT	16 mm
Tetracycline	TE	15 mm
Chloramphenicol	C	21 mm
Vancomycin (glykopeptid)	VA	17 mm

\*potentiated by a beta-lactamase inhibitor

Diagnosatics of  
enterococci and  
G+ rods  
(+ pictures)

# Description of criminals (diagnostics 1)

	Enterococ.	Listeria	Coryneform	Bacillus
Microscopy	G+ cocci short chains  <a href="http://textbookofbacteriology.net">http://textbookofbacteriology.net</a>	G+ rods chains or palisades	G+ rods palisades	G+ robust rods, sporulating (sometimes non visible)
Cultivation	greyish, as large as that of <i>S.</i> <i>agalactiae</i> , various types of haemolysis	like enterococci of <i>Strep.</i> <i>agalactiae</i>	very tiny colonies, like flour	large colonies, sometimes intensive haemolysis

# Enterococci – colonies

<http://microbiology.mtsinai.on.ca>



# Description of criminals (diagnostics 2)

## Enterococci

<http://www.morgenwelt.de>



- **Biochemical tests:** catalase negative, possible biochemical determination, **arabinose splitting** (*E. faecalis* does not split, green medium, *E. faecium* makes it yellow)
- **Antigen analysis** used rarely. (Originally „group D streptococi“ according to Lancefield, as genus *Enterococcus* did not exist in time of Lancefield research)
- **Atb testing** on common MH agar. There exist also screening media for VRE.

# Description of criminals (diagnostics 3)

## G+ rods

- Biochemical tests: catalase positive in all three of our genera. But e. g. genus *Arcanobacterium* (not member of *Corynebacterium* genus, but nevertheless a coryneform) is CAT neg. Biochemical detection possible (API Coryne, Remel)
- Growth at low temperatures, high NaCl concentrations etc. used in *Listeria* dg.
- Biochemical dg. and atb testing are also a part of the diagnostics
- Antigen analysis – e. g. searching diphtheria toxin

# Photos of criminal database 1

## Enterococci

Microscopy

Bile-aesculin

Slanetz-Bartley




Photo: archive of the Inst. both left pictures made by Prof. MVDr. Boris Skalka, DrSc.

# Photos of criminal database 2




1, 2, 3 archive of the Inst.

4 <http://medinfo.ufl.edu>

## Rods I

## Corynebacterium Gram



Listeria – BA, Gram





# Photos of criminal database 3

## Rods II – corynebacteria, forms



# Photos of criminal database 4



Rods III

*Bacillus cereus*



*Arcanobacterium haemolyticum*

*Bacillus subtilis*



Differential  
diagnostics of  
enterococci and  
G+ rods

# Differential diagnostics: enterococci

- Gram staining differentiates Gram + cocci, Gram + rods and other bacteria.
- Catalase of NaCl differentiates staphylococci
- Slanetz-Bartley / Bile-aesculin, PYR test differentiates enterococci from streptococci
- Arabinose test/other biochemical tests – mutual differentiation of Enterococci

# Differentiation of *Enterococcus*



- **Arabinose test:** colonies are mixed with arabinose and indicator, and let to incubate

Green	negative	<i>E. faecalis</i>
Yellow	positive	<i>E. faecium</i>

- ENCOCCUStest has only 8 reactions, but otherwise it is like other similar tests

# Differential diagnostics – *Bacillus*

- **Gram staining** differentiates **G+ rods** from other bacteria
- *Bacillus* has typical **Gram staining result** – long and large rods; sometimes (not always) also endospore formation may be seen (empty places in the rod)
- **Cultivation** is also characteristic (large, felt-like colonies)
- **Species determination** available by biochemical tests, susceptibility to antibiotics etc.

*There is no clear algorithm for G+ rods!*

# Differential diagnostics – *Listeria* and coryneforms

- Gram staining differentiates G+ rods from other bacteria
- In case of spore-non forming, non-robust rods the microbe is likely *Listeria* or one of coryneform rods (mere absence of endospore is not sure!) Further diagnostics is available by means of biochemical methods, growth at various temperatures, tests of haemolytic interactions (synergisms, antagonisms) etc.

*There is no clear algorithm for G+ rods!*

# Bile-aesculin agar

<http://www.geocities.com>



2. AESCULIN BILE AZIDE AGAR

Colonie di *E. faecalis*, nere, esculinasi positive.



# *Listeria* growth at 4 °C

- Among Gram positive rods, only *Listeria* is able to grow in low temperatures. This enables it to spread in cheese factories
- Among other bacteria (not being G+ rod), there are some more species able to grow at such low temperatures (*Yersinia*, some *Pseudomonas* sp.)

# Chromogenic medium for Listeria


www.oxoid.com



There exist various chromogenic media to Listeria diagnostics. The one on the picture is called ALOA and it is typical by **o blue colour of all Listeria colonies**, furthermore, **pathogenic species have also halo around them** (halo = differently coloured surrounding).


# Elek test

It is a detection of a toxin of *Corynebacterium diphtheriae*. We use a paper with specific antitoxin, that is put on the surface of the agar, then tested strains are inoculated. Positive result = precipitation lines.



# The end

<http://www.cdphe.state.co.us>



*Bacillus anthracis*

