


Burns

Z. Rozkydal



Epidermis

Dermis – blood vessels, nerves, sebaceous glands, sweat glands, hair roots

Fatty tissue

Burns

Dry burn- flames, contact with hot objects

Scald- steam, hot liquids

Electrical burns- low- voltage current
high- voltage current
lightening strikes


Cold injury- frostbite, contact with freezing metal

Chemical burns- industrial chemicals, acid, alkali

Radiation burns- sunburn, exposure to radioactive source

Depth of burn

1. Superficial- outermost layer of the skin
redness, swelling, tenderness
healing complete in 1-2 weeks
2. Partial thickness burn- epidermis
blisters, redness, raw skin
 - a/ superficial- healing in 2 weeks without consequences
 - b/ deep- healing in 3-6 weeks, scars develop, risk of infection
transplants are needed
3. Full- thickness burn- all the layers of the skin are affected
damage to the nerves, fat tissue, muscles, blood vessels
dark, black skin, infection, necrectomy, transplants, sequels



Superficial burn

Partial thickness burn

Full thickness burn

Extent of burns: rule of 9

Head	9 %
Upper limbs	9 + 9 %
Trunk	18 % + 18 %
Lower limbs	18 + 18 %
Perineum	1 %

Risk of shock in burns

Children up to 10 years of age- 10 % of body surface

Adults- above 20 % of body surface

Signs

Severe pain, local signs of a burn

Oedema, escape of fluids and proteins from injured area

Hypovolemic shock

Damage to the airways- bronchospasm, dyspnoea

First aid

Stop the burning as soon as possible

Maintain open airways

Remove him from the site of injury

Start cooling the injury as soon as possible

- tap water, ice packs – at least 10 minutes

Remove any rings, watches, belts, shoes

Cover the area with a kitchen film, plastic bag
or apply sterile dressing

Do not remove sticking cloths

Do not burst any blisters

Do not use adhesive dressing, ointments and powders

Do not to allow him to eat and drink

Treat him as a shocked casualty

Monitor vital signs

Chemicals

Acids (dark skin)

Alkali (yellow- brown, green skin)

First aid

Cooling with a plain water

Acids

soap water, soda water

Alkali

Lemon water, vinegar water

Sunburn

Overexposure to the sun or sunlamp

High altitudes

Some medicines can trigger high sensitivity to sunlight

Skin is red, blistered, painful

First aid

Remove him from the sun

Cold water on the skin

Sun lotions

Frequent sips of cold water

Heat exhaustion

Loss of salt and water from the body through excessive sweating

Develops gradually

In humid conditions, dancing in warm environment etc.

Signs: fatigue, exhaustion, vomiting, diarrhoea

First aid

Put him to a cool, shady place

Plenty of water to drink

Monitor vital signs

Heatstroke

The body is overheated

Prolonged exposure to heat or due to a high fever

Drugs

Signs

Headache, dizziness, confusion

Hot, flush and dry skin

Rapid pulse

Body temperature over 40° C

Impaired consciousness

First aid

Remove him to a cool place

Cold wet sheet, cold water

Cold drinks

Hypothermia

Body temperature falls below 35° C

Severe hypothermia- below 30° C- is often fatal

Prolonged exposure to cold

Wind- chill factor, immersion to cold water

Poorly heated houses, homeless people

Thin and frail persons, alcohol, chronic illness

Signs

Shivering, pale, dry skin

Apathy, desorientation

Slow breathing and pulse

Impaired consciousness

Hypothermia

First aid

Take the casualty to a sheltered place

Remove any wet cold clothing

Rewarm him

Warm room, warm blankets

Insulating material, dry sleeping bag

Survival bag

Monitor vital signs

In children- mechanism for regulating body temperature is underdeveloped – first aid and medical advice

Frostbite

1. Pale skin, less sensation
2. Yellowish skin with blisters, no sensation
3. Necrotic skin

Frostbite

Occurs in freezing or cold and windy conditions

Pale skin, numbness

Hardening of the skin

Stiffening of the skin

Change of colour – white, mottled, blue, red

painful, blistered, black in gangrene

First aid

Put him to a warm place

Remove gloves, rings etc, sterile dressing

Rewarm him slowly

Avoid direct heat

Transport to hospital