
#### 14


# **Motor system III**

# Hierarchic organization of motor system




#### Basal ganglia

- Corpus striatum
  - Nucleus caudatus
  - Putamen
- Globus pallidus (Pallidum)
  - Externum
  - Internum
- Nucleus subthalamicus
- Substantia nigra
  - Pars compacta
  - Pars reticulata
- Thalamic motor nuclei



# Basal ganglia


- Corpus striatum
  - Nucleus caudatus
  - Putamen
- Globus pallidus (Pallidum)
  - Externum
  - Internum
- Nucleus subthalamicus
- Substantia nigra
  - Pars compacta
  - Pars reticulata
- Thalamic motor nuclei



## **Basal ganglia - inputs**

#### **Corpus striatum**

- Connections from all cortical areas with two exceptions – primary visual and primary auditory cortex
- The most of connections from
  - Frontal and parietal association areas
  - Motor areas




# **Basal ganglia - inputs**

#### **Corpus striatum**

- Connections from all cortical areas with two exceptions – primary visual and primary auditory cortex
- The most of connections from
  - Frontal and parietal association areas
  - Motor areas


Motor gating – initiation of wanted and inhibition of unwanted movements




## Basal ganglia


- Motor control realized by two circuits
  - Direct pathway
    - Motor cortex activation

- Indirect pathway
  - Motor cortex inhibition







 Thalamic motor nuclei activate motor cortex




- Thalamic motor nuclei activate motor cortex
- Tonic inhibitions of thalamic motor nuclei by GPi




- Thalamic motor nuclei activate motor cortex
- Tonic inhibitions of thalamic motor nuclei by GPi
- Activated corpus striatum transiently inhibits Gpi, resulting in transient disinhibition of thalamic motor nuclei







NS activates GPi



- NS activates GPi
- GPe tonically inhibits NS




- NS activates GPi
- GPe tonically inhibits NS
- Corpus striatum transiently inhibits
  GPe



- NS activates GPi
- GPe tonically inhibits NS
- Corpus striatum transiently inhibits
  GPe


NS disinhibition



- NS activates GPi
- GPe tonically inhibits NS
- Corpus striatum transiently inhibits GPe

NS disinhibition


Gpi activation



- NS activates GPi
- GPe tonically inhibits NS
- Corpus striatum transiently inhibits GPe

NS disinhibition


Gpi activation



Less important is a direct inhibition of Gpi by GPe

## Direct and indirect pathway differences


- Direct pathway
- Motor cortex activation
- Indirect pathway
- Motor cortex inhibition



# Direct and indirect pathway differences


- Direct pathway
- Motor cortex activation
- Indirect pathway
- Motor cortex inhibition

Indirect pathway may be considered as a "handbrake" of the direct pathway




# **Dopaminergic projections**

- Dopaminergic projections are crucial for the function of corpus striatum
- S. nigra pars compacta



# Dopaminergic projections

- Dopaminergic projections are crucial for the function of corpus striatum
- S. nigra pars compacta
- Direct pathway activation
  - D1 receptors
- Indirect pathway inhibition
  - D2 receptors




#### Basal ganglia

- Beside motor loop there are other loops associated with other thalamic nuclei
- "Gating" of the other sort of information
- Association loop
- Limbic loop
- Basal ganglia play an important role in information processing in general and this is crucial for thinking process
- Connections of corpus striatum are plastic what allows learning and this was very important during evolution

#### Cerebellum

#### Coordination



#### Cerebellum

Cerebellum plays an important role not only in the coordination of movement, but also in the "coordination" of thoughts