

## Topic P08: Laboratory diagnostics of tuberculosis, actinomycetes and nocardiae

### Task 1: Microscopy of acid-fast and partially acid fast microorganisms


While entirely acid-fast microorganisms (*Mycobacterium*) cannot be stained at all according to Gram, only partially acid-fast ones (*Nocardia*) or not at all (*Actinomyces*) can be Gram-stained, but they stain irregularly; they also tend to form branched filaments.

#### a) Staining of (negative) clinical material using Ziehl-Neelsen staining method

Ziehl-Neelsen staining is used for mycobacteria (*M. tuberculosis*, *M. leprae*), but also for some parasites (*Cryptosporidium parvum*, *Cyclospora cayetanensis*). The acid-fast organisms are stained only when heated during staining\*, but then they are not decolorized even by so called „acid alcohol“ (solution of alcohol with HCl or H<sub>2</sub>SO<sub>4</sub>). Decolorized background is then counterstained.

Stain the negative sputum according to the Ziehl-Neelsen method (methylene blue variant). It is not likely that acid-fast rods would be present. Observe in microscope (immersion). Draw the results; at least, you will see the background, e. g. leucocytes, epithelia and other objects. Do not forget to **describe** your picture (use lines)!

Describe also the staining procedure – fill in the following table with names of used reagents.



1.	During the staining the preparation is _____ until _____	
2.	This reagent is made of _____	and _____
3.	Instead of this reagent, it is also possible to use _____	_____


#### b) Microscopy of a mycobacterial culture

Examine microscopically (immersion oil, immersion 100× objective) the preparation of mycobacterial culture stained by Ziehl-Neelsen staining method.

Evaluate presence of red acid-fast rods.

Draw observed structures.


Do not forget to **describe** your picture (use lines)!



#### c) Microscopic examination of actinomycetes and nocardia strains

Examine microscopically the slide stained by Gram. Describe and draw observed formations. Observe high polymorphism of the microorganisms (from coccal shape, through rods to fibre/strings, often branched; Gram-positive, but often staining half Gram-negative).

Do not forget to **describe** your picture (use lines)!



### Task 2: Culture of mycobacteria, *Actinomyces* and *Nocardia*.

The culture requests of acid fast and partially bacteria are very different.

- ❖ For *Mycobacterium tuberculosis* we use special media: liquid media (Šula, Banič) and solid media (Ogawa, Löwenstein-Jenssen). The solid media are different from majority of other solid media used in medical microbiology; they do not contain agar, they are „solid“ because of coagulated egg proteins. Before culture, the medium should be specially treated.
- ❖ For *Nocardia* a current blood agar is sufficient.
- ❖ For *Actinomyces* we need VL-agar and culture in anaerostat/anaerobic jar (see P07), as this organism is microaerophilic with so low need for oxygen that anaerobic conditions are optimal for it.

\*Heating may be eventually substituted by use of highly concentrated carbolfuchsin and highly concentrated phenol; this modification of Ziehl Neelsen staining (Kinyoun modification) does not require heating.

**a) Describe media for mycobacterial cultivation**

Medium name	liquid/solid	colour	notes

**b) Describe and draw the growth of *Mycobacterium*, *Actinomyces* and *Nocardia* on (in) given media**

Bacterium	Medium name	Presence/absence of growth, eventually growth character (use your own words to characterize the growth)
<i>Mycobacterium</i>		
<i>Actinomyces</i>	blood agar	
	VL agar	
<i>Nocardia</i>	blood agar	
	VL agar	

**Task 3: Assessment of antimicrobial drugs susceptibility**

For treatment of mycobacterial infections, it is necessary to use special drugs, called antitubercotics. The way of testing is different from other bacteria, too: antitubercotics are added directly to the culture media. On the other hand, *Actinomyces* and *Nocardia* are treated by „normal“ antibiotics and also „normal“ diffusion disc test is used for testing.

**a) Assessment of mycobacterial susceptibility to antitubercotics**

By comparing with a control test-tube, read the results of antitubercotic susceptibility tests of *Mycobacterium tuberculosis* strain.

Antitubercotic				Growth control
Growth Y/N				
Interpretation				

**b) Antibiotic susceptibility of *Nocardia* and *Actinomyces***

Perform in vitro susceptibility testing of *Nocardia* and *Actinomyces* to suitable antibiotics.

Complete the table with the abbreviations of the antibiotics according to the card and for all the tested strains, measure the diameter of the susceptibility zones. On your card, you have limit zones – according these, interpret the zones as susceptible (S) or resistant (R). There are no “intermediate” interpretations this time.

Strain →				
Antibiotic (full name)	Zone Ø (mm)	Interpretation	Zone Ø (mm)	Interpretation


**Task 4: PCR in diagnostics of TB**

As the culture of mycobacteria is complicated, PCR becomes a very important method in its diagnostics. Read a result of PCR TB diagnostics (from slideshow), write the results and interpret them.

Patient No.	Sample band	Control band	Interpretation
1			
2			
3			
4			

**Task 5: Diagnostics of leprosy**

Leprosy is a disease that still affects millions of people in underdeveloped countries. Its laboratory diagnostics is difficult because *Mycobacterium leprae* does not grow on artificial media. Fill in the following table.

	The name of this animal is	
	It is used to produce	
	and this substance is used for	

Picture source: [http://www.1-costaricalink.com/costa\\_rica\\_fauna/nine\\_banded\\_armadillo.htm](http://www.1-costaricalink.com/costa_rica_fauna/nine_banded_armadillo.htm)

**Task 6: Indirect TB detection by means of QUANTIFERON®-TB Gold test**

It is a test of induced interferon gamma release checking and by means of this, checking of the cell-mediated immunity. **Test principle:** It was proven that in TB, including latent TB, tuberculosis antigens activate T-lymphocytes and they produce big amounts of interferon gamma. Similarly those T-lymphocytes may be activated non-specifically by so called mitogenem; that is why mitogene is used as a positive control (MIT). As a negative control we use a test tube containing nothing (NIL). The test tube with proper TB antigen is labeled "TB". Interferon itself is detected by ELISA reaction.

Interpret the Quantiferon-TB Gold examination in four patients with use of interpretation table.

Anna: MIT = 4.8 TB = 1.2 NIL = 1.1 Your interpretation: \_\_\_\_\_

Berta: MIT = 5.3 TB = 4.8 NIL = 2.1 Your interpretation: \_\_\_\_\_

Cecil: MIT = 0.9 TB = 0.9 NIL = 0.8 Your interpretation: \_\_\_\_\_

Dimos: MIT = 8.4 TB = 8.3 NIL = 8.2 Your interpretation: \_\_\_\_\_

(all values are in IU/ml)

**Interpretation table (according to test recommendations; simplified!)**

NIL	TB minus NIL	MIT minus NIL	Final test interpretation	Presence of infection <i>M. tuberculosis</i>
≤ 8,0	< 0.35	≥ 0.5	negative	Not likely
	≥ 0.35	any value	positive	Likely
> 8,0	< 0.35	< 0.5	unsure	Cannot be determined
	any value	any value		