








# Respiratory system












Bucket-handle and water-pump handle effects







páka  $A - B < A' - C'$  → zvedání žebere



páka  $A - B' > A' - C$  → klesání žebere

PLEURA  
pulmonalis      parietalis







$$P = \frac{2 T}{r}$$

Hollow spherical organ





# TRANSPORT O<sub>2</sub>




# Regulation of breathing

# Control of ventilation







- **Breathing is an automatic process that takes place unconsciously. Automaticity of breathing comes from regular (rhythmic) activity of groups of neurons anatomically localized in the medulla and its vicinity.**


- They can be divided into three **main groups**:
  - *dorsal respiratory group* – placed bilaterally on the dorsal side of the medulla oblongata, only inspiratory neurons, sending axons to motoneurons of inspiratory muscles (diaphragm, external intercostal muscles; their activation=inspiration, their relaxation=expiration; participates on inspiration at rest and forced inspiration
  - *ventral respiratory group* - located on the ventrolateral part of the medulla oblongata, the upper part: neurons whose axons of motor neurons activate the main and auxiliary inspiratory muscles; the lower part: expiratory neurons which innervate expiratory muscles (internal intercostal muscles). Neurons in this group operate only during forced inspiration and forced expiration.
  - *Pontine respiratory group* - *pneumotaxic center* - dorsally placed on top of the pont, contributes to the frequency and depth of breathing; affects the activity of respiratory neurons in the medulla oblongata.

# Chemical factors affecting the respiratory center:


## Central chemoreceptors

- on the front side of the medulla
- sensitive only to increase of arterial  $p\text{CO}_2$  (by increasing  $\text{H}^+$ )

- Notice:
- central chemoreceptor are stimulated by other types of acidosis (lactate acidosis, ketoacidosis)








## Peripheral chemoreceptors


– located in the aortic and carotid bodies  
-primarily sensitive to decrease in arterial  $pO_2$ , particularly to decrease of  $O_2$  under 10-13 kPa in the arterial blood.

They convey their sensory information to the medulla via the vagus nerve and glossopharyngeal nerve.


Mechanism of action: Decreased ATP production in mitochondria leads to depolarization of receptors membrane and to excitation of chemoreceptor




### A LOCATION OF CAROTID AND AORTIC BODIES



### B MICROSCOPIC ANATOMY OF CAROTID BODY





## Modulation of respiratory output

Major parameters for feedback control – classical gases:  $pO_2$ ,  $pCO_2$ , pH

In addition to these, the respiratory system receives input from two other major sources:

1. **variety of stretch and chemical/irritant receptors** that monitor the size of airways and the presence of noxious agents/receptors in respiratory system
2. **Higher CNS centers** that modulate respiratory activity for the sake of nonrespiratory activities

**Irritants receptors** on mucosa of respiratory system – rapidly adapting

Stimulus: agents - chemical substances (histamine, serotonin, prostaglandins, ammonia, cigarette smoke).

Response: increase mucus secretion, constriction of larynx and bronchus

**C-fibre receptors** (juxtacapillary=J receptors)– free nerve ending of n.vagus (unmyelinated axon) in interstitium of bronchus and alveolus;

Stimulus: Mechanical irritants (pulmonary hypertension, pulmonary oedema)+chemical

Response: hypopnoea, rapid shallow breathing, bronchoconstriction, cough

**Stretch receptors** slowly adapting (mechanoreceptors in tracheobronchial tree that detect the changes in lung volume by sensing the stretch receptors of the airway wall), inform to brain about the lung volume to optimize respiratory; its irritants triggered decrease activity of respiratory centre – **Hering-Breuer's reflexes**. (protecting the lungs from overinflation/deflation)


**Baroreceptors** – suppresses activity of respiratory centre


Irritants of **proprioceptors of muscles, tendons** during active and pasive movements of limbs  
Influenced activity of respiratory neurons (increase minute ventilation during work load)

**Limbic system, hypothalamus** – strong pain, emotion

Tractus corticospinalis =cortex – activated RC during work load

**temperature**






## ENDOTRACHEAL CANNULA




# HERING-BREUER REFLEX

REFLEX STOP BREATHING









# VAGOTOMY



# Periodic breathing





- It is not regular, rhythmic, but respiration occurs in periods ("a moment to breathe, take a moment to not breathe,,")
- **CHEYNE-STOKES**
- **BIOT'S**
- „gaspings“
- **KUSSMAUL**

**A INTEGRATED PHRENIC NERVE ACTIVITY**



0 0.2 0.4  
Time (min)

**B LUNG VOLUME**



0 0.5 1.0  
Time (min)