

Institute for microbiology shows



TRACING THE CRIMINAL

Part fourteen:
Repeating

Organisation of practical exam

- Usually student picks **one of 60 tasks; nobod says that all of them are always available**
- One topic of practical sessions corresponds to **2 to 4 tasks**; some tasks are related to more than one practical sessions (e. g. ASO – related to both neutralization and streptococci)
- Some tasks are mostly practical (like Gram staining), some are rather discussion with practical parts. **Students may be asked also to topics not directly placed in the tasks**
- **Students may be told to fill in a protocol to their tasks (not necesarilly always)**

Přehled témat

Methods

Special microbiology

Clinical microbiology

Methods

J01 Microscopy

- **Wet mount** – for large and/or motile microbes (parasites, fungi, motile bacteria)
 - **To know also: Dark field wet mount** (mainly spirochets)
- **Gram staining** – how to do it, + survey of other staining methods (Giemsa, Ziehl Neelsen, Burri...)
- **Observation of already stained preparations** – mainly interpretation
- Also some more tasks from other topics are microscopical!

Wet mount – do it practically

- Do not forget to cover the preparation by a coverslip and to use **non immersion** objective lenses, magnifying e. g. 4×, 10× or 40×.
- We use no immersion oil
- After having it done, observe the objects in the microscope
- **Attention! Students that put into contact a non-immersion objective and immersion oil, finish immediately with F!!!**

Wet mount – procedure

10x
40x
100x
100x

Stained preparation

DRY 10
X 10

Gram, Lugol, Alcohol, Safranin
Tap water use
Drying with filtration paper

V C ROSCO 2V

V V 25 0 0

100 x 0 3. C V

Gram staining – procedure

- Gentian/crystalline violet = Sol. Gram-Nowy (20 –) 30 sec.
- Lugol (20 –) 30 sec.
- Alcohol 15 (– 20) sec.
- rinse by tap water!!! important!
- Safranin 60 – 120 sec.
- rinse by tap water
- dry by filtration paper
- microscopy as in Task One

Specimen microscopy





Photo O. Z.



Strain microscopy

In observation of slides...

- ...it is important to have basic knowledge of size and morphology (yeast × staphylococci etc.) and to know something about interpretation (= knowledge of clinical microbiology, not just J01!)
- For example:
 - WBCs = inflammation
 - no WBCs in sputum = not properly taken specimen
 - G- diplococci inside WBC, urethral swab – suspicion for gonorrhoea

J02 – Culture.

Reinoculation of a strain

Sterilize your loop

Take the strain

Inoculate first phase

Sterilize your loop

Do not take the strain again

Inoculate second phase

Sterilize your loop

Do not take the strain again


Inoculate third phase

Sterilize your loop

Do not take the strain again

Inoculate the „serpent“

crossing of
previous lines!



Attention!

- The task is more difficult than it seems (student forget to **sterilise the loop** between the steps, or they forget to **cross the lines**)
- On the other hand, **technical excellence of the lines is not rated** (we understand that you are neither experienced microbiologists or laboratory assistants)
- So it is about understanding (maybe also explaining) of **principle of cross inoculation**

One more sub-topic: Explain the function of given media

1. Broth
2. VL-broth
3. selenite broth
4. Sabouraud
5. Löwenstein-Jenssen
6. Blood agar
7. Endo agar
8. MH
9. 10 % NaCl
10. VLA
11. XLD (+ MAL)
12. CHA
13. Levinthal
14. Slanetz-Bartley

Some more media at special bacteriology tasks.

J03 – biochemical identification tests

One only task, but other stuff is in other topics!

- In special bacteriology, you might meet:
 - Catalase test
 - Tests with diagnostic strips (oxidase, PYR, INAC)
 - Hajna medium (red = G-NF, other = ENT/VIB)
 - Eventually also MIU (not necessary to know details)
- The only „pure biochemical“ task
 - ENTEROtest 16 or something similar (STAPHYtest etc.) – you would get everything that you need to read the test

Do not forget the ONPG test

it should be read before the other tests

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
ONPG	H	G	F	E	D	C	B	A	H	G	F	E	D	C	B	A	
	First row of the plate								2 nd row of the plate								
+																	
-																	
?																	
?	+	-	+	+	+	-	-	-	-	-	-	-	-	+	+	+	+
	1	2	4	1	2	4	1	2	4	1	2	4	1	2	4	1	2
	5			3			0			0			6			3	

Decontamination tests – J04

- Important basic information:
- If we want to find **survival limit** of bacteria, we have to remove the tested extreme parameters to the conditions and to let them then in optimal conditions for a sufficient time.
 - In testing of disinfection effect, bacteria are treated by a disinfectant and then cultured on a medium without disinfectant
 - In testing of sterilisation, bacteria are placed to the sterilizer and then cultured in normal conditions

Task: Assessment of effectivity of disinfection

- It is not sufficient to say how-much-percent disinfection is the result, it is also important to say (and to explain!) that **it is bactericidal, not bacteriostatic concentration of the given disinfectant.**
- (In the agar, where the bacterium is cultured, it is no disinfection anymore.)

Sterilisation parameters

Sometimes the action of factor **combines**

The factor always important is the **time**


A resistant, spore forming bacterium	160 °C	170 °C	180 °C
20 min	survives	survives	dies
30 min	survives	dies	dies
60 min	dies	dies	dies

Plus (to both tasks):

- An extra sub-task to both tasks: Make pairs of cards with names of methods/disinfectants and cards of characterisation of methods
- The cards are on a working table, it is possible to see it


J05: Atb susceptibility

Diffusion disc test: to read it, to interpret it



It is not a separate task, but a part of tasks of special microbiology

Microdilution test – reading



For interpretation, comparison with breakpoints is necessary

○ **MIC** ○ **growth control – kontrola rústu**

- In case of columns 1, 3, 4 & 5, $MIC > \text{the highest value}$
- *In wells 8 and 11, $MIC \leq \text{the lowest value}$*

E-test – it is not a separate task, but you should know it


www.uniklinik-ulm.de

We can read the MIC value directly on the strip – in place, where the margins cross the strip




Beta-lactamases detection: limited knowledge is sufficient

They also may be a part of a task




J06–J08 (but also related with many others): serology

Antigen detection: laboratory (animal origin) antibodies + patient's sample or microbial strain.



Direct method

Antibody detection: laboratory antigen (microbial) + patient's serum (or saliva).




Indirect method

Interpretation – important to know

- **Antigen detection:** (including antigen analysis): it is a direct method. Positive result means presence of the microbe in the patient's body
- **Antibody detection:** it is an indirect method. Nevertheless, there are some ways how to get the information – when the microbe met the body:
 - Amount of antibodies (relative – titre + titre dynamics; agglutination, CFT, neutralisation)
 - Class of antibodies: IgM/IgG (reactions with labelled components – mostly ELISA and immunoblotting)
 - (*Avidity of antibodies*)


Precipitation




Two examples of precipitation

1) Microprecipitation in agar according to Ouchterlony – not a task, but might come as a sub-question

2) RRR/RPR reaction for syphilis diagnostics (flocculation)




Agglutination




Agglutination for antigen detection or antigen analysis

- **EPEC detection:** in what situation we perform it, how it is performed... (Practically, you obtain a strain and you have to know what to do with it)
- **CSF agglutination** (task „Comment a videoclip“). Important: besides microscopy this is the second way what to do as quick diagnostics of purulent meningitis

Agglutination reaction for detection of antibodies




Aglutination on carriers




Example
MHA-TP

(Inst. for Microbiology)



CFT – principle



Neutralisation schematically


- Antibody (Ig) prevents an effect of a toxin/virus to a cell / red blood cell



Cell in a tissue culture or a red blood cell

Antibody

Toxin or virus



Cell in a tissue culture or a red blood cell

Toxin or virus

Examples of neutralisation reactions

	Neutralised	Object	Reaction
	Bacterial toxin (haemolysin)	RBC haemolysis	ASO
	Virus	RBC agglutination	HIT
	<i>Virus</i>	<i>Cell metabolic effect</i>	<i>VNT</i>

Important: What is the antistreptolysin O


and why we attempt to detect it

- After every streptococcal infection antibodies are produced, often including antibodies against streptococcal toxin – streptolysin O.
- Nevertheless, sometimes after infection the antibodies increase instead of decreasing. Antibodies are bound to some structures of the host organism (autoimmunity), so a „circulus vitiosus“ starts to run
- In such a situation, paradoxically the antibodies are worse than the pathogen that challenged the antibody response to protect us.


HIT

- **H**aemagglutination **I**nhibition **T**est: Pay attention, it is NOT an agglutination reaction, it is a neutralisation! Antibody neutralises the aggregation of RBCs due to viruses.
- **So: Potato-like shape = negative response. Dense round target = positive response**
- HIT differs from ASO reaction mostly by the fact, that **the RBCs are not haemolyzed, but agglutinated**. But the fact, that a specific antibody blockages the reaction is valid in both of the

Reactions with labelled components



ELISA – an example (Inst. for Microbiol.)



One task: HBsAg / anti-HBS puzzle

- HBsAg testing – positive
- HBsAg testing – negative
- anti-HBs testing – positive
- anti-HBs testing – negative

Reading of ELISA

BL	4	BL	4
K-	5	K-	5
K-	6	K-	6
K+	7	K+	7
K+	8	K+	8
1	9	1	9
2	10	2	10
3	11	3	11
IgA		IgG	

$$c. o. (IgA) = (0.107 + 0.137)/2 + 0.320$$

$$c. o. (IgA) = 0.122 + 0.320 = 0.442$$

$$90 \% c. o. = 0.398 \quad 110 \% c. o. = 0.486$$

all values bellow 0.398 are negative

all values above 0.486 are positive

$$c. o. (IgG) = (0.034 + 0.029)/2 + 0.320$$

$$c. o. (IgG) = 0.032 + 0.320 = 0.352$$

$$90 \% c. o. = 0.317 \quad 110 \% c. o. = 0.387$$

all values below 0.317 are negative


all values above 0.387 are positive

**FIND POSITIVE AND BORDERLINE
WELLS FOR BOTH IgA and IgG!**

Western blotting – principle


- 1: original antigen (mixed)
- 2: decomposition of antigen by a detergent
- 3: electrophoresis division of antigen

- 4: „blotting“ of divided antigen to a nitrocelulose membrane
- 5: ELISA reaction (only some antibodies present)




Western blot – example

(picture from Inst. for Microbiology)



Immunochromatography



Important!

- It is necessary to know not only how to read the test, but also how to interpret it.
- There are three special tasks (concerning toxoplasmosis, Lyme disease and hepatitis A/B/C) based on complex interpretation of all results including anamnesis!
 - E. g. pregnant woman with IgG anti-toxoplasmosis is NOT ill, but protected!

J09: DNA detection (PCR)


Mostly use of PCR in medical microbiology

- The methods are used mostly in situations, where **microscopic and culture diagnostic is difficult or impossible**
- It is not very useful for common, ubiquitous **pathogens**. Because of its sensitivity they would detect accidental molecules coming from environment
- The methods are **neither useless**, as some people think, **nor all-problems-solving**, as some other people suppose.

Survey of interpretation

Proper reaction	Internal control	Interpretation
negative	positive	negative
negative	negative	inhibition of reaction
positive	positive	positive
positive	negative	(highly) positive

An example of a gel



proper reactions

internal controls

Patients 1 and 4 – positive, patient 2 – negative, patient 3 – inhibition of reaction. 5 – positive control, 6 – negative control, 7 – ladder

Special

medical


microbiology

J10+11: viruses

- Majority of viral tasks are serological examinations (HBsAg, anti-HBs)
- Two extra virology tasks concern isolation of viruses
 - Fertilized egg – parts of fertilised egg, used for isolation
 - Cytopathic effect – what is it, how to find it

Fertilized egg

(+ how to get the info that virus is there)




SH – shell
membrane
(paper
membrane)

AB – albumen
<http://www.scielo.cl/fbpe/img/bres/v38n4/fig02.gif>


AM – amniotic sac, YS – yolk sac, AL – allantois

CH – chorioallantoic membrane (CAM)



http://cmir.mgh.harvard.edu/cellbio/cellculture.php?menuID_=122

HSV Growing in Tissue Culture



www.herpesdiagnosis.com/diagnose.html

(HSV is Herpes Simplex virus – HSV 1 causing mostly herpes labialis, HSV 2 herpes genitalis)

J12 Parasitology

- As a basis, we use methods based on **modified wet mount**:
 - In **Kato method** counterstain with malachite green is used, to make parasites better visible
 - **Faust method** is a concentration one (see later)
- **Graham method** is used in pinworms only (and as one task you can do it practically!)
- *Wet mount „sensu stricto“ and stained preparations (e. g. trichrom) are used in increased suspicion for intestinal protozoa (either primarily, or after seeing Faust and Kato)*

Morphology of eggs of intestinal parasites


You should know at least these shapes to the examination – another task



Pinworm

Enterobius

Trichuris




Roundworm

Ascaris

Tapeworm

Taenia



Pictures taken from CD-ROM „Parasite-Tutor“ – Department of Laboratory Medicine, University of Washington, Seattle, WA

Toxoplasmosis – another task, including definition of patients

- P: healthy pregnant woman, cats at home
- Q: another healthy pregnant woman, no cats
- R: a young lady trekking in forest; no cats, but contact with objects contaminated by faeces of wild animals
- S: a senior, working in garden, cats use to walk through the garden, symptoms of retinitis + enlarged lymph nodes

J13 – mycology

- Yeasts are examined like bacteria from P01–P06 (see later)
- In filamentous fungi you are supposed to determine the fungus according to the microscopical preparation by comparison with pictures of fungi from the practical session

J14: biofilm

Comparison of methods for blood catheter examination

- Classical cultivation in broth
- Maki semiquantitative method
- Sonification (also semiquantitative!)

Reading + discussin: advantages and disadvantages of all methods, possible interpretation


J14: biofilm

Influence of saccharides presence to dental plaque formation: like in practical, including 3D-graph making

- Assess the influence of uptake of various amounts of saccharides in food on rate of biofilm formation in a cariogenic *Streptococcus mutans*.


What are the conclusions of this experiment, as to amounts of saccharides in food, how long they stay in oral cavity etc.?

Wells in the panel



Another task: MBEC assessment

MBEC ... minimal
biofilm eradicating
concentration



P01 to P06, J13

- There is a universal task to those topics:
„Among given strains, find strain(s) of Xxxxx, perform more detailed diagnostics (and perform antibiotic susceptibility testing).
- It is necessary to follow the logical algorithm, for example like this:
Gram staining → catalase test → plasmacoagulase test → STAPHYtest 16 etc.

Exceptions:

- ASO is examined as other serological tasks (but important to know the meaning of the test for clinical practice)
- G+ rods are not examined in this algorithmic way, but you get pictures and you have to say „this looks like *Corynebacterium*, this does not look like *Corynebacterium*, because it is spore forming“ etc.
- Very similar is also a task to *Clostridium tetani* (to P07)

Survey of diagnostics (simplified)





(or other tests)

Enterococcus or



Corynebacteria, forms



J07 Anaerobic jar

description, explaining function

air-proof lid

palladium catalyser
(beneath the lid)

construction for placing
of Petri dishes

Anaerobiose generator
(packet with chemicals)



Anaerobic box (again)



source of anaerobic gases

space for entering culture plates

entrances for hands of personel

Clostridium tetani morphology (practical recognition from other bacteria)

<http://www.geocities.com>



Detection of lecithinase

- Lecithinase production is detected as **strain precipitation on the yolk agar**. Nevertheless, there are many lecithinases, and one only, that of *Clostridium perfringens* is interesting for us, we have to test, whether the lecithinase may be inhibited by a specific antitoxin.


„Negative I“ no lecithinase production.

„Negative II“ a lecithinase is produced, but not the tested one



Survey of other tests, e. g. animal experiment and immunochromatography

- Look at the picture of tetanic mouse



Drawing by Petr Ondrovčík (1959–2007)

Graphically adapted.

Background counterstained using not malachit green, but „Paint“ programme by Microsoft

www.biotox.cz




Opisthotonus is typical both for mice and humans

P08: Ziehl-Neelsen stain:


principle and knowledge about results (red bacteria on blue or green background)




1 Cover smear with carbol-fuchsin. Steam over boiling water for 8 minutes. Add additional stain if stain boils off.




2 After slide has cooled decolorize with acid-alcohol for 15 to 20 seconds.




3 Stop decolorization action of acid-rinsing briefly with water.



4 Counterstain with methylene blue for 30 seconds.



5 Rinse briefly with water to remove excess methylene blue.




6 Blot dry with bibulous paper. Examine directly under oil immersion.

How it is examined

- You have to **know how to do it + to know how a positive Ziehl-Neelsen looks like** and according to that knowledge to be able to differentiate its picture from other (e. g. Gram-stained) pictures
- Plus (like in the next task) **knowledge of other methods of TB diagnostics**, basic diagnostics of actinomycetes and nocardias for sub-questions

Ziehl-Neelsen stain



Another task: Culture of mycobacteria

- Hydroxide should be used before culture
- We use liquid Šula or Banić media and egg Ogawa or Löwenstein-Jenssen media. Egg media are solid because of egg white coagulation, they do not contain agar
- Results are read after 1 (check for contamination) 3, 6 and for sure after 9 weeks of culture. (Positive results are mostly found after 6 weeks of culture.)
- *+ knowledge of more methods (PCR etc.)*

Appearance of mycobacterial colonies

<http://www.stockmedicalart.com/>



Indirect diagnostics of TB

- **Cell mediated immunity**
- Quantiferon testing including understanding of importance of „MIT“ and „NIL“ test tubes
- Explaining the role of indirect TB diagnostics

P09: Serology of syphilis + serology of Lyme disease (see Serology part)

Historical	BWR – Bordet Wassermann	Nontr.
Screening	RRR – Rapid Reagin Test or RPR or VDRL test	
	MHA-TP (TPHA)	Treponema
Confirmatory	ELISA	
	FTA-ABS (indir. imunofluor.)	
	Western Blotting	
<i>Historical, or superconfirmation</i>	<i>TPIT (Treponema Pallidum Immobilisation Test) = Nelson</i>	



Basically you should also know direct tests!

Clinical

microbiology

„sensu stricto“

P10–P13: Clinical microbiology I-IV

Sampling and decision making globally (P10):

- Several tasks, all of them the same:
- „For three minicasuistics, find suitable sampling methods and vessels/swabs for sampling“
- Knowledge of swabs and vessels necessary (you are supposed to find practically the corresponding sampling kit)

Some types of swabs



Plain (dry) swab

www.calgarylabservices.com

Today its use is for PCR and antigene detection only, not for culture!




Amies medium with charcoal www.herenz.de

Universal transport medium for bacteriology (all types of swabs). The wire variant important, if we want to go „behind the corner“

More swabs

Fungi Quick (for yeast and molds)

www.copanswabs.com



C. A. T. swab (for Candida And Trichomonas,
from genitals only www.copanswabs.com)




Virus swab

www.copanswabs.com


Chlamydia swab

www.copanswabs.com


Vessels



Common test tube. Universal use: clotted blood (serology), urine, CSF, pus, punctate etc.; blood and urinary cathethers, parts of tissue...



Sputum vessel. Not only for sputum, but also larger parts of tissue etc.



Stool vessel, for parasitology. Only this one does not have to be sterile!




Vessel for urine sampling. It is better, if the patient urinates just into a test-tube, but especially for women it is difficult (except if they are in shower). So they can urinate into this vessel, and then a nurse removes the urine into a test-tube.

Clinical microbiology (continuing)

- Several more tasks. One: Find a pathogen in oropharyngeal flora
- Normal flora consists of greyish, viridating colonies (oral streptococci) and yellowish, usually non-haemolytical colonies (oral neisseriae). Possible pathogens are:
 - Haemolytic streptococci (and also *Staphylococcus aureus*) are visible by a strong haemolysis on blood agar
 - For haemophili detection we use antibiotic disc with bacitracin – higher concentrations than in bacitracin test (to decline the normal microflora) – plus *Staphylococcus* line
 - For meningococcal detection we use another disk, with mixture of vancomycin and colistin

Detection of pathogen in throat/sputum



1 swab inoculation

2 loop inoculation

3 staphylococcus line

4 bacitracin disc (for hemophili)

5 V + K disc (colistine and vancomycine) for meningococci


In all parts of inoculated area we search for colonies with haemolysis. They could be streptococci (rather colourless) or goldish)

Cultivation result of throat swab with common flora


In these sites we search for haemophili



Photo: InstMicr



Stool cultivation



Negative result is ready after 48 h

Positive in 72 h or more

*If not written differently, culture runs at 37 °C

Urine specimen cultivation

- **Task: Perform semiquantitative and qualitative examination of urine**
- As **likely contamination** (or accidental finding) is counted everything below 10^4 / ml, everything below 10^5 / ml in finding of two various bacteria and everything in three/more bacterial strains
- It is not necessary to count exactly the colonies, just to say what is the approximate amount
- **Among pathogens**, the most common are enterobacteria, enterococci, *S. agalactiae*, staphylococci etc.

Semiquantitative urine evaluation

Number of colonies	Number of CFU (bacteria) in 1 μ l of urine	Number of CFU (bacteria) in 1 ml of urine	Evaluation (valid for 1 bacterium)
Less than 10	Less than 10	Less than 10^4	Contamination
10 – 100	10 – 100	10^4 - 10^5	Borderline
More than 100	More than 100	More than 10^5	Infection

Nugent score counting

Due to the fact that it is a microscopic and not the culture proof, we work with so-called morphotypes. For example, bacteria belonging to the "morphotype Lactobacillus" may not be Lactobacillus, but it is very likely



- **Morphotype Gardnerella/Bacteroides:** not present = zero point, + = one point, ++ = two points, +++ = three points, ++++ = four points
- **Morphotype Lactobacillus:** the contrary: not present = four points, positivity ++++ = zero points
- **Curvet Gram-labile rods:** none = 0 points, + or ++ = one point, +++ or ++++ = two points

Vaginal culture evaluation

Vaginal swabs are usually cultured on following media:

- blood agar (for common bacterial pathogens)
- Endo agar (or McConkey agar)
- Agar with 10 % NaCl (for staphylococci)
- Special blood agar variant for *Gardnerella vaginalis* (GVA agar)
- WCHA agar (anaerobic culture) – only sometimes

Swab and imprint



Pictures of imprint methods are taken from a slideshow by MUDr. Zdeněk Chovanec from I. Surgical Clinic of Medical Faculty of Masaryk University and St. Anna Faculty Hospital in Brno

Wound swabs (without anaerobes):

Possible diagnostic scheme

- (Different in different types of wound etc.)
- Day 0: start of culture only
- Day 1: result of primary culture of specimen on blood agar (BA), Endo agar, NaCl and BA + amikacin. If all solid media are negative, broth is observed; if turbid, a subcultivation to solid media is performed
- Day 2: expedition of negative and some positive results; too resistant bacteria → more tests
- Days 3, 4: expedition of remaining results

Blood cultures – sampling

- It means **not clotted blood**, principally very different from serological examinations (*it is neither antibody nor antigen detection, microbe should remain alive and is detected by cultivation*)
- Today we usually sample into **special vessels with transport-cultivation medium** for automatic culture (*sooner just not-clotted blood without any medium was sent*)
- We have to ensure **minimalisation of risk of pseudobacteraemia** (see more)
- **In adults we take 10 to 20 ml of blood, in children usually 1–5 ml according to age** (sampling is more difficult in adults, and also in children also less bacteria are important)

Function of cultivators

- **Cultivator, connected to a computer**, keeps automatically optimal conditions of cultivation, and also evaluates status of the vessel and indicates eventual growth (e. g. change of reflectance, i. e. optical characteristics of the vial)
- The growth is **signalized optically and by a sound. When nothing is growing even after a week**, the apparatus signalizes it too (it is time to give out a negative result)

See you at the examination!



Photo: InstMicr

