

Periapical diseases.

Markéta Hermanová

Periapical periodontitis - aetiology

- Pulpitis and pulp necrosis
- Trauma
- Endodontic treatment

Periapical periodontitis

- Acute periapical periodontitis
- Chronic periapical periodontitis (periapical or apical granuloma)
- Acute periapical abscess and spread of inflammation


Periapical periodontitis

- Dynamic process; inflammation can vary with time
- Outcome reflects the balance between the nature, duration, and severity of the irritant and the effectiveness of the host defences
- Bacterial infection of the root canals is the major cause of clinically significant periodontitis
- Can follow acute traumatic injury to periapical tissues without pulp necrosis
- Endodontic treatment, instrumentation of infected root canal

Acute periapical periodontitis

- Acute inflammatory exudate in the periodontal ligament (between root apex and alveolar bone)
- Pain elicited by external pressure (pulpitis well located)
- Hot or cold stimulation does not induce pain, as it would in pulpitis
- Radiography usually normal; no bone resorption yet
- Acute periapical or alveolar abscess can develop directly

Periapical abscess





Chronic periapical periodontitis (periapical or apical granuloma)

- Persistent irritation, bacteria (anaerobes predominate) in the pulp chamber and root canals results in chronic periapical periodontitis
- Resorption of periapical alveolar bone, replacement by granulation tissue, periapical granuloma
- Root with attached periapical granuloma

Apical and periapical granulomas

- Asymptomatic (may remain quiescent for long periods) or symptomatic
- Chronically inflamed granulation tissue around apex of a non-vital tooth
- Infection and antigenic challenge from endodontic flora
- Stimulation of proliferation of rests of Malassez within the lesion (=radicular cysts)


Periapical granuloma



Radiological findings in apical and periapical granulomas.


- Margins reflects the dynamics of the lesion
- Widening of periodontal ligament space at the beginning
- Active bone resorption = margins ill defined
- Static lesions = bone apposition and the formation of a zone of sclerosis (=osteosclerosis)

Periapical Lucency




Apical granuloma

< 1.5 cm
corticated margin



Radicular cyst

> 1.5 cm
corticated margin




Abscess

less well defined
margin

Sequelae of chronic periapical periodontitis

- Antigenic challenge and host's immunological response in balance = quiescent granuloma
- The balance disturbed = enlargement of the granuloma, bone resorption
- Bacteria invading the granuloma from the root canal = acute exacerbation = acute symptoms = enlargement of granuloma = abscess formation
- Suppuration in the granuloma
- Development of radicular cyst
- Osteosclerosis (=bone apposition)
- Hypercementosis (=apposition of cementum)

Radicular cyst



牙根囊腫

Aetiology and microbiology

Frequently detected bacterial species in periapical abscesses

Microaerophilic streptococci

S. Milleri group e. g. *S. anginosus*

Anaerobic streptococci

Peptostreptococcus species,
e. g. *P. anaerobius*

Gram-positive anaerobic rods

Actinomyces species, e. g. *A. israelii*
Eubacterium species, e. g. *E. lentum*

Gram-negative anaerobic rods

Porphyromonas species, e. g. *P. gingivalis*
Prevotella species, e. g. *P. intermedia*
Bacteroides species
Campylobacter species
Fusobacterium species, e. g. *F. nucleatum*

Routes of spread

- Increase in hydrostatic pressure causes pus to track along lines of least resistance
- Pus directly into oral cavity through a sinus following penetration of periostium and mucosa
- Palatal mucoperiostium resistant = palatal abscess
- Abscesses in molar region penetrate the buccal cortical plate spreading into soft tissues = cellulitis
- Abscesses at anterior maxillary teeth = perforation of the labial bone = spreading to inner canthus of the eye and lower eyelid, obliteration of nasolabial fold, into upper lip
- Abscesses at maxillary molars and premolars = into the maxillary sinus
- Abscesses at mandibular premolars and molars = involvement of submandibular, sublingual and lateral pharyngeal spaces, and anteroposteriorly under the skin surface
- Abscesses at mandibular incisors and canine = labially, perforate the bone, subcutaneous abscess in the midline between attachments of mentalis muscles

Cellulitis

- Rapidly spreading inflammation of the soft tissues
- Usually associated with streptococcal infections (related to the release „spreading enzymes, e. g. hyaluronidase, stroptokinase)
- diffuse, tense, painful swelling of soft tissues; malaise, elevated temperature
- Risk: cavernous sinus thrombosis; extension into submandibular and cervical tissues = respiratory embarrassment; pain, trismus

Ludwig's angina

- Severe cellulitis involving the submandibular, sublingual, and submental spaces
- May involve the pharynx and larynx
- Oedema of the glottis; risk of death by suffocation

Ludwig's angina



Figure—Submandibular and sublingual erythema and swelling typical of Ludwig's angina.

