

# MUNI MED

# Physiology of reproduction.























Summary of the key events and associations during growth and puberty.


	What happens	When	Factors involved
Mini-puberty	Gn and sex steroid levels rise after nadir at birth. Facilitates testicular descent in M and gonadal cell populations mature	2–3 months of age	T levels in M reach mid-pubertal levels. Oestradiol levels similar in M and F.
Adrenarche	'Awakening of adrenal gland' causes development of body odour, oily skin and hair, pubic hair (pubarche)	Usually begins at 6–8 years of age and is independent of true central puberty- usually precedes gonadarche by approx. 2 years	Production of androgens by adrenal cortex including DHEA, DHEAS and androstenedione
Thelarche (F)	Onset of female breast development, or budding (Tanner stage 2), usually first sign of puberty in girls	Usually occurs after 8 years of age (mean 10–11, range 8–13 years)	Rising oestrogen levels
Gonadarche	Growth of ovaries and testes and increased sex steroid production (true central puberty).	Testicular enlargement in M usually signals pubertal development. Ovarian growth can't be directly seen but usually coincides with thelarche	Activation of gonads by LH and FSH to increase T and oestrogen levels
Pubarche	Development of first pubic hair	First pubic hair occurs at adrenarche (can be transient) and then again at Tanner stage 3. In F usually 6 m after thelarche.	Rising levels of androgens (during adrenarche) and sex steroids during central puberty
Growth spurt	Peak growth velocity seen in childhood after infancy. Occurs shortly before final height is reached.	Usually occurs at Tanner stage 2 in F and 3–4 in M. Always occurs before menarche in F.	Multiple hormones involved. Rising Oestrogen levels cause GH secretion and act directly at GP. Occurs later in M as T needed in higher conc to convert to oestradiol via aromatase
Spermarche (M)	Development of sperm in the testicle and first ejaculation	Usually coincides with development of secondary sexual characteristics in mid- puberty	Increased T levels from Leydig cells and nocturnal LH surge
Menarche (F)	Onset of menstruation (Often thought of as culmination of pubertal development in F)	Usually occurs 2 years after thelarche and soon after growth spurt	Oestrogen stimulated growth of uterus and vascularity of endometrium, leading to sloughing of part of the lining. Most menstrual cycles are initially anovulatory.

## Menstrual cycle






# **Negative feedback**






## **Positive feedback**






# **Ovarian cycle**





**Endometrial cycle** 



# **Contraceptives**

## BARRIER CONTRACEPTIVES:

- condoms
- contraceptive sponges with spermicide
- diaphragms
- cervical caps

#### HORMONAL:

- oral pills
- implants under the skin
- injections
- patches
- IUDs
- · vaginal ring

## INTRAUTERINE DEVICES:

- copper IUDs
- IUD with levonorgestrel

#### **BEHAVIORAL**:

- fertility awareness methods
- · coitus interruptus
- lactation




#### STERILIZATION:

- tubal ligation
- vasektomy



# **Oral pills**





## **Oral pills**

### – Progestational effects include:


- Inhibition of ovulation by suppressing luteinizing hormone (LH);
- Thickening of cervical mucus, thus hampering the transport of sperm;
- Possible inhibition of sperm capacitation;
- Hampered implantation by the production of decidualized endometrium with exhausted and atrophic glands

### – Estrogenic effects include:

- Partial inhibition of ovulation in part by the suppression of follicle-stimulating hormone (FSH) and luteinizing hormone (LH), depending on dose;
- Alteration of secretions and cellular structures of the endometrium within the uterus



## **Benefits and riscs of HC**





## **Functions of the PLACENTA**

#### TRANSPOR function

- Respiratory gases
- transport and metabolism of sacharides
- transport and metabolism of aminoacods
- transport and metabolism of fat
- transport of H2O, minerals and vitamines

#### ENDOCRIN function


- Estrogens
- Progesteron
- HCG
- HPL
- Growth factors (epidermal and insulin-like growth factors)

#### PROTECTIV function

- Cytochrom P450
- Pinocytosis (IgG)
- Barier against bacterias, virus etc.




## **HCG**






## **HCG**






## RX





## RX






## **HCS/HPL**

#### **HCS (Human Chorionic Somatomammotropin)**


placental growth hormone - Human placental lactogen (hPL)

- has weak actions similar of growth hormone
- causes retention of nitrogen, potassium, calcium
- causes decreased insulin sensitivity and decreased utilization of glucose in the mother
- secretion of this hormone increases progressively in direct proportion to the weight of the placenta






## **PROL**







## OX






## OX








## **Progesteron**






### **Estroen**











## Fetoplacental unit






# Fetoplacental unit






### Lactation



## Physiological changes during pregnancy





### THANK YOU FOR YOUR ATENTION

