

M U N I
M E D

M U N I
M E D

10

Vision II

Image formation




Table 26-1 Differences Between Rods and Cones and Their Neural Systems

Rods

- High sensitivity to light, specialized for night vision
- More photopigment, capture more light
- High amplification, single photon detection
- Low temporal resolution: slow response, long integration time
- More sensitive to scattered light

Rod system

- Low acuity: not present in central fovea, highly convergent retinal pathways
- Achromatic: one type of rod pigment

Cones


- Lower sensitivity, specialized for day vision
- Less photopigment
- Lower amplification
- High temporal resolution: fast response, short integration time
- Most sensitive to direct axial rays

Cone system

- High acuity: concentrated in fovea, dispersed retinal pathways
- Chromatic: three types of cones, each with a distinct pigment that is most sensitive to a different part of the visible light spectrum


Retina

- Photoreceptors
- Interneurons
 - Horizontal cells
 - Horizontal interconnection
 - Bipolar cells
 - Vertical interconnection
 - Amacrine cells
 - Both horizontal and vertical interconnection
- Ganglion cells
 - AP generation
 - Transmission of AP to the brain





Retina







- Fovea
 - Low convergence
 - Small receptive field
 - High resolution
 - Lower sensitivity to light
- Periphery of retina
 - High degree of convergence
 - Large receptive field
 - Low resolution
 - High sensitivity to light








Receptive field





Receptive field



Receptive field




Receptive field




Receptive field

- Magnocellular system
 - Large receptive field
 - Rods and cones
 - **M ganglion cells (10%)**
 - High speed of velocity
 - Brightness/low contrast sensitivity
 - Minimal sensitivity to color



Receptive field



- Magnocellular system
 - Large receptive field
 - Rods and cones
 - **M ganglion cells (10%)**
 - High speed of velocity
 - Brightness/low contrast sensitivity
 - Minimal sensitivity to color
- Parvocellular system
 - Small receptive field
 - Cones and rods
 - **P ganglion cells (80%)**
 - Low speed of velocity
 - Low sensitivity in low contrast
 - Good sensitivity to color



<http://www.slideshare.net/drpsdebs/presentations>


Receptive field

- Magnocellular system
 - Large receptive field
 - Rods and cones
 - **M ganglion cells (10%)**
 - High speed of velocity
 - Brightness/low contrast sensitivity
 - Minimal sensitivity to color
- Parvocellular system
 - Small receptive field
 - Cones and rods
 - **P ganglion cells (80%)**
 - Low speed of velocity
 - Low sensitivity in low contrast
 - Good sensitivity to color



Optic nerve and optic tract


- Optic nerve
 - Signal from one eye
 - Signal from „whole“ visual field
- Optic tract
 - Signal from both eyes
 - Signal from half of visual field





Copyright © 2008, 2004, 1998, 1993, 1988, 1983 by Moebius, Inc., an affiliate of Elsevier Inc.

Optic nerve and optic tract

- Optic nerve
 - Signal from one eye
 - Signal from „whole“ visual field
- Optic tract
 - Signal from both eyes
 - Signal from half of visual field



<https://www.aafp.org/afp/2013/0901/p319.pdf>




Copyright © 2008, 2004, 1998, 1993, 1988, 1983 by Moebius, Inc., an affiliate of Elsevier Inc.

<http://www.slideshare.net/CsillaEgri/presentations>

Visual pathways


- Nucleus corporis geniculati lateralis
 - Thalamus
 - Majority of projections
 - Via optic radiation to neocortex
- Hypothalamus
 - Regulation circadian activity
- Prepectum
 - Pupillary reflex
- Colliculi superiores
 - Reflex movement of eyes and head



<http://www.slideshare.net/drpsdeb/presentations>





Nucleus corporis geniculati lateralis


- Six nuclear layers
- Retinotopic organization
- Each layer receives input from only one eye
- Layers 1-2
 - Magnocellular system
 - M ganglion cells
 - Large receptive field/brightness sensitivity
 - Information about localization and movement
- Layers 3-6
 - Parvocellular system
 - P ganglion cells
 - Small receptive field / color sensitivity
 - Information about form and color





Primary visual cortex

Retinotopic organization







<http://www.slideshare.net/CsillaEgri/presentations>




Primary visual cortex

Retinotopic organization





Copyright © 2008, 2004, 1998, 1993, 1988, 1983 by Mosby, Inc., an affiliate of Elsevier Inc.



MUNI
MED


Primary visual cortex

- ✓ Retinotopic organization
- ✓ Columnar organization
 - Orientation columns
 - Respond to specific orientation
 - Blobs
 - Respond to color
 - Ocular dominance column
 - Information from left or right eye
- ✓ Horizontal connections




Spatial vision

Binocular




Spatial vision


Binocular



Monocular – based on previous experience




Motion perception




Vestibuloocular reflex

Rotational VOR



Translational VOR



Nystagmus


- Involuntary rhythmic eye movement
- Physiological
 - Postrotational
 - Optokinetic
- Pathologic
 - ✓ Peripheral
 - Vestibular system pathologies
 - ✓ Central
 - CNS damage (cerebellum, midbrain...)

Classifying nystagmus


The various types of jerk and pendular nystagmus are illustrated below.

JERK NYSTAGMUS

Convergence-retraction nystagmus refers to the irregular jerking of the eyes back into the orbit during upward gaze. It can indicate midbrain tegmental damage.



Downbeat nystagmus refers to the irregular downward jerking of the eyes during downward gaze. It can signal lower medullary damage.




Vestibular nystagmus, the horizontal or rotary movement of the eyes, suggests vestibular disease or cochlear dysfunction.




PENDULAR NYSTAGMUS

Horizontal, or pendular, nystagmus refers to oscillations of equal velocity around a center point. It can indicate congenital loss of visual acuity or multiple sclerosis.



Vertical, or seesaw, nystagmus is the rapid, seesaw movement of the eyes: One eye appears to rise while the other appears to fall. It suggests an optic chiasm lesion.



http://dxline.info/img/new_ai/nystagmus.jpg

Saccadic eye movements




<https://en.wikipedia.org/wiki/Saccade#/media/File:Szakkad.jpg>

Saccadic eye movements



<https://en.wikipedia.org/wiki/Saccade#/media/File:Szakkad.jpg>




<https://s-media-cache-ak0.pinimg.com/564x/51/f7/26/51f7267e7c8a59caa90f904cd4f965eb.jpg>

26TH PRAGUE INTERNATIONAL FILM FESTIVAL


21.-29.03.2019



MUNI
MED



Pupillary reflex



http://www.ubooks.pub/Books/B0/E27R7642/MAIN/images/1509_Pupillary_Reflex_Pathways.jpg

78. The basic physiology of visual system – rods and cones function, on/off receptive field, nervus opticus vs. tractus opticus

- Rods and cons function
 - Characterization and comparison
 - Phototransduction mechanism and adaptation
- Brief overview of retina organization (retina process receptor potential – analog, AP is generated in ganglion cells)
- Receptive field organization
 - On/off receptive fields
 - Magnocellular system (BW)
 - Parvocellular system (Color)
- Nervus opticus vs. tractus opticus
- Projections from tractus opticus (Main centers in the brain involved in visual signals processing)

M U N I
M E D