


Epiphyseal disorders

Z. Rozkydal

Epiphyseal disorders

Idiopathic avascular
necrosis of epiphysis
of long bones

Etiology unknown



Obr. 1

Epiphysis

Metaphysis

Diaphysis

Metaphysis

Epiphysis

Perthes disease

It is a complication of the necrosis of proximal epiphysis of the femur

4 -12 years

10 % bilateral


More often in boys

Symptoms: limping, pain
limited ROM (rotation, abduction)




Obr. 2

Perthes disease- stages





Obr. 4




Obr. 5

Frejka classification

1. st. latency 6 - 12 months
2. st. necrosis
3. st. decalcination
4. st. recovery- fragmentation
5. st. consequences





Ischemic necrosis of epiphysis

Loss of vascularity of epiphysis

Necrosis of cartilage

Microdamage in osseous part- resorption


Diminished mechanical strength

Damage of the physeal plate- coxa vara, coxa brevis

Radiolucent lesions in metaphysis

Perthes disease

Ischemia of the whole epiphysis
Articular cartilage continues to grow
Bone is resorbed and replaced
by woven bone
The bone is soft and vulnerable
Subchondral fracture
- shows the extent of damage
New bone is gradually revascularised
New bone is plastic-
can be deformed



Obr. 6
Subchondral fracture
of femoral epiphysis

M. Perthes

1. Ischemic stage: avascular necrosis
growth arrest of epiphysis
revascularisation from periphery
ossification
2. Ischemic stage: trauma, subchondral fracture
resorption under the fracture
replacement by plastic woven bone
subluxation, deformity

Catterall classification


I. 25 %

II. 50 %

med.- lateral column

III. 75 %

IV. 100 %



Obr. 7



Catterall I



Catterall II




Catterall III





Catterall IV

Dr. SURESH K. N. S.




Subchondral fx
less than one half

Subchondral fx
more than one half



Catterall I

Obr. 8



Catterall II


Obr. 9



Obr. 10



Catterall III



Obr. 11



Catterall IV

Examination

X-ray

MRI

Scintigraphy

Ultrasonography



X-ray



MRI



MRI

Prognosis

I. and II. stage	good prognosis
III. and IV. stage	wrong prognosis


Risk factors:

Older age


Loss of containment, subluxation

Large extent

Limited movements



Types of deformity in Perthes disease



Stulberg classification of deformity
of the femoral head in Perthes disease

Management

- containment of the head in the acetabulum
- good range of motion

Conservative methods

- Atlanta orthosis, no weightbearing

Operative methods

Osteotomy of the pelvis (Salter, Steel, Sutherland, Dungal)

Osteotomy of the femur

Conservative methods

Rest in bed


Crutches

Atlanta orthosis




Obr. 16

Atlanta orthosis



Experiment: Ibandronate + BMP




Experiment: Osteoprotegerin

Operative methods


Salter pelvic osteotomy

Obr. 17





Varus osteotomy of the femur


Obr. 18



Salter osteotomy



Obr. 19



Obr. 20



Perthes disease on the right hip
after Salter osteotomy
Almost normal hip in 18 years of age

Consequences of Perthes disease

Coxa plana

Shortening of the leg

Limited movements



Early osteoarthritis

Better prognosis

Younger age

Less extent of damage

No subluxation



Tibia vara Blount


Disorder of proximal epiphysis
of the tibia

Early arrest of growth plate on medial
side with smaller epiphysis

Infantile – up to 3 years

Juvenile - up to 10 years


Th: orthosis, osteotomy



Tibia vara Blount



Obr. 24



Obr. 25

Slipped upper femoral epiphysis

Growth plate of proximal epiphysis

of the femur is weak and soft

Imbalance of growth hormon and
sexual hormones


Obese patients

Fröhlich syndrom

Adiposogenital syndrom

9-15 years

Bilateral in one third





Obr. 26

Slipped upper femoral epiphysis

Slipping of epiphysis
down and backwards
to varus and to retroversion

Metaphysis goes proximally
and to external rotation



Symptoms


Pain in groin and in the thigh

Limping

Shortening of the leg

Limited abduction and external rotation


Positive Trendelenburg sign



Obr. 28


Types

1. Preslip (6%)
2. Acute slip (11%)
3. Chronic slip (after two weeks, 60 %)
4. Acute slip on chronic slipping (23%)



Obr. 29

Stages



1.

2.

3.

Obr. 30

Stages

1. Slight: slip up to 30%
2. Moderate : slip 30-60 %
3. Severe: slip above 60 %

Management


Fixation in situ (K wires, Knowles pins)

Closed reduction and K wires

Open reduction

Osteotomy of proximal femur -
Southwick, Imhäuser-Weber

Fixation in situ




Obr. 31




Obr. 32

Open reduction





Obr. 33

Southwick osteotomy



Pertrochanteric osteotomy



Obr. 35

Complication of slipped upper femoral epiphysis

Avascular necrosis of the femoral head

Chondrolysis of the femoral head

Osteoarthritis of the hip

Necrosis of os lunatum m. Kienböck

Therapy

Rest

Immobilisation

Removal and replacement
by tendon, by os pisiforme
or by artificial material




M. Köhler I. - necrosis of navicular bone

Therapy

Rest

Immobilisation

Arthrodesis



Obr. 37

M. Köhler II.
M. Freiberg-Köhler
Necrosis of metatarsal head

Therapy

Rest, padding

Surgery:
Removal of necrotic part
Osteotomy



Obr. 38

Avascular necrosis of femoral head in adults



Etiology unknown


Pain

Limited movements

Limping

Obr. 39

Avascular necrosis of femoral head



Obr. 40

Etiology unknown

72 % bilateral

Without management- 85 % progress into colaps
of the femoral head

5-12 % indications to THA

Genetic background

Risk factors

Table 1 Conditions that may cause or are related to ONFH

Trauma

Femoral neck fracture

Hip dislocation

Extensive burns

Direct vessel trauma

Hypercoagulation

Deficit of antithrombin III

Deficit of protein C

Deficit of protein S

Resistance to activated protein C

Deficit of plasminogen activator inhibitor

Surplus of inhibitor for plasminogen activator

Factor V Leiden mutation

Secondary conditions of hypercoagulation

Corticosteroids

Alcoholism

Hemoglobinopathy

Trombophilia

Corticosteroids

Haemoglobinopathies (sickle-cell disease)

Polycythemia

Metabolic diseases

Hyperparathyroidism

Gout

Cushing's disease

Gaucher's disease

Alimentary system diseases

Pancreatitis

Ulcerative colitis

Chrohn's disease

Other risk factors

Smoking

Decompression disease

Radiation

Chemotherapy

Hemodialysis

HIV infection

Secondary conditions of hypercoagulation

Corticosteroids

Alcoholism

Myelodysplastic syndromes

Pregnancy

Oral contraceptive use

Hyperlipidaemia

Collagen diseases

Ehler–Danlos syndrome

Raynaud's disease

Diabetes mellitus

Antiphospholipidaemic antibodies (APLA)

Diagnosis

Bone infarction at the onset is asymptomatic


Groin pain, around the hip, limping

X-ray


MRI








X-ray
Subchondral changes



MRI



Specimen



Subchondral fracture

Management

Cons: crutches, bisphosphonates

physiotherapy, drugs for promotion of vascularity

Oper.:

Forrage, decompression, drilling, bone grafting

Long cylindrical bone graft


Osteotomy –varus, valgus, rotation

Free vascularized fibular graft - stage II , III.

Nonvascularised bone grafts


Drilling + stem cells + BMP

THA




Preop.

Vascularised fibular graft
5 y.




Trabecular metal Tantal rods
4 y. post op

11 y. postop.



10 y postop
Asymptomatic.

Vascularised fibular graft



LED, percutaneous drilling – Steinman pin

Necrosis after fracture
of the neck of the femur



Necrosis of the femoral head
after coxitis



M. Ahlbäck – necrosis of medial condyle of the femur

m. Osgood- Schlatter – proximal apophysis
of the tibia

Osteochondrosis dissecans

Necrosis of sesamoid bone

M. Panner – osteonecrosis of humeral head

Vertebra plana Calvé

Necrosis of apophysis of calcaneus

