

**MUNI PHARM**

**Molecular Biophysics**

Biophysics

23

**Water**

- in the human body, the proportion of water is ca 60%
- water content in the blood is 90%, in the bodies of about 80% (depending on the fat content), 76% of muscles, skeleton 22%
- compound of fundamental importance: loss of 10% of water is a serious complication; the loss of 25% = death
- **polar compound**, bonding angle = 104,5°
- formation of **hydrogen bonds** (cluster structure of water): influence of physico chemical properties (e.g. boiling point, heat of vaporization, heat capacity)
- the hydrogen bond is the electrostatic attraction between polar molecules that occurs when a hydrogen atom bound to a highly electronegative atom: H<sub>2</sub>O, HF, HCl

24

**Water**

- in the human body, the proportion of water in the bodies of ca 60%
- water content in the blood is 83%, in the bodies of about 80% (depending on the fat content), 76% of muscles, skeleton 22%
- compound of fundamental importance: loss of 10% of water is a serious complication; the loss of 25% = death
- **polar compound**, bonding angle = 104,5°
- formation of **hydrogen bonds** (cluster structure of water): influence of physico chemical properties (e.g. freezing point, heat of vaporization, heat capacity)
- the hydrogen bond is the electrostatic attraction between polar molecules that occurs when a hydrogen atom bound to a highly electronegative atom: H<sub>2</sub>O, HF, HCl

25

**Water and living organisms**

1. **Very good solvent** (due to its polarity and H-bond formation)
2. **Transport functions** (transport of gases, nutrients, heat)
3. **Acid – base properties** (dissociation, chem. reaction)
4. **Mechanical function** - fills cavities (joints) and body structures
5. **Thermoregulatory functions**
  - a) good thermal conductivity: heat exchange between the inside of the body and the environment (heat loss by conduction)
  - b) large heat capacity: heat storage - preventing climate change and reduces temperature fluctuation throughout the day
  - c) large latent heat of vaporization: cool the body through perspiration (evaporation of sweat)
6. **Density anomalies of water** (max. ρ at 3.98 °C)
  - life under the ice

26

**Colligative properties of solutions**

- Lowering the vapor pressure (Raoult's law  $p_A = x_A \cdot p^*_A$ )
- Elevation of the boiling point
- Depression of the freezing point
- Osmotic pressure

The solution has higher  $T_b$  than the pure solvent

The solution has a lower freezing point than the pure solvent (coniferous trees - metabolism even at low temperatures, sap is saturated solution of biopolymers)

$\pi = iRT \cdot C_{solute}$

$i$  - the correction factor on the number of dissociation particles

**For intravenous administration must be used isotonic solutions (have the same osmotic pressure as blood plasma); e.g. 0.9% NaCl or 5% glucose**

27

**Colligative properties**

**Osmotic pressure: infusion preparations**

The osmotic pressure of parenteral preparations directly affects the balance of body fluids in the extracellular and intracellular space. Osmotic concentration (**osmolality**) is expressed in osmoles:

1 mol/kg NaCl ≈ 2 osmol/kg NaCl (NaCl dissociates into 2 ions)

The basic osmotically active particles are plasma ions (mainly Na<sup>+</sup> and Cl<sup>-</sup>).

Osmotic concentration of solutions is expressed as **osmolality** (osmol/l), can be converted from osmolality. Osmolality of blood plasma 280-296 mosmol/L.

28

### Colligative properties

**Osmotic pressure: The diagnostic significance**

- **Hyperosmolality of blood plasma**

**Cause:** water loss, diabetic coma, burns, kidney failure, sepsis, drowning in salt water, acute intoxication by small molecules (e.g. ethylene glycol, methanol).

**Clinical manifestations:** from mild neuropsychiatric disorders to coma

- **Hypoosmolality of blood plasma**

**Cause:** excess of water, the metabolic response to trauma, drowning in fresh water, inappropriate ADH secretion.

**Clinical manifestations:** weakness, nausea, lethargy, headache, brain edema.

The osmotic pressure in the body is **controlled by the hypothalamus** (antidiuretic hormone, ADH). Increased resorption of water by the kidney is the result of ADH.


29

### Colligative properties

**Oncotic pressure**

Osmotic pressure, which is caused by solutions containing particles with a high molecular weight (e.g. **proteins**).

- Oncotic pressure contributes to maintaining a **sufficient circulating blood volume**.
- Decreased protein content (hypoproteinemia) in the blood plasma cause **edema**.
- Endothelium is permeable to ions but poorly permeable to proteins => oncotic pressure is applied instead of osmotic pressure.



30

### Dissociation: Influence of pH on passage through biological membranes

- Many drugs are **weak acids or weak bases**
- These compounds are **more water soluble in their ionized form**, and vice versa **more liposoluble in their non-ionized form**
- **pH in the GIT determines** the degree of ionization of weak acids and bases, and thus determines **their solubility and absorption of drug into the body**

$$HA \leftrightarrow A^- + H^+ \quad K_A = \frac{[A^-][H^+]}{[HA]}$$

$$[H^+] = K_A \frac{[HA]}{[A^-]} \quad -\log [H^+] = -\log K_A - \log \frac{[HA]}{[A^-]}$$

$pH = pK_A - \log \frac{[HA]}{[A^-]}$  **Henderson-Hasselbach equation**

$pK_A - pH = \log \frac{[\text{nonionized form}]}{[\text{ionized form}]}$  **weak acid**

$pH - pK_A = \log \frac{[\text{nonionized form}]}{[\text{ionized form}]}$  **weak base**

31

### Dissociation: Influence of pH on passage through biological membranes

- **Stomach (pH = 1-3)**  
Facilitate absorption of **weak acids (nondissociated)**; **bases are not absorbed**.

**Sample problem (pH = 3 or 1; pKa = 3):**

50% of the drug is nonionised

$$3 - 3 = \log \frac{[HA]}{[A^-]} \Rightarrow \log \frac{[HA]}{[A^-]} = 0 \Rightarrow \frac{[HA]}{[A^-]} = 1$$

$$3 - 1 = \log \frac{[HA]}{[A^-]} \Rightarrow \log \frac{[HA]}{[A^-]} = 2 \Rightarrow \frac{[HA]}{[A^-]} = 100$$

- **Small intestine (pH = 5-7.5)**  
**A weak acids and bases are good absorbed; a strong acid and a strong base are absorbed less.**

**Sample problem (pH = 6; pKa = 4 or 5):**

99% of the drug is nonionised

$$4 - 6 = \log \frac{[HA]}{[A^-]} \Rightarrow \log \frac{[HA]}{[A^-]} = -2 \Rightarrow \frac{[HA]}{[A^-]} = 0.01$$

$$5 - 6 = \log \frac{[HA]}{[A^-]} \Rightarrow \log \frac{[HA]}{[A^-]} = -1 \Rightarrow \frac{[HA]}{[A^-]} = 0.1$$

1% of the drug is nonionised

10% of the drug is nonionised

32

### Dissociation: Acid-base balance (ABB)


- dynamic balance between acidic and alkaline substances inside the body

- Precise adjustment of pH is important since pH changes influence the properties of proteins (including the activity enzymes), transport mechanisms, properties of the membrane channels etc.
- Due to metabolism pH is changed => **regulation of ABB by buffers** (substances capable of releasing or binding H<sup>+</sup>).

$$H_2PO_4 \leftrightarrow HPO_4^{2-} + H^+ \quad \text{equilibrium} \Rightarrow \text{partial dissociation}$$

☠ **Blood plasma pH = 7.4 (7.35-7.45); pH values below 7.0 and above 7.8 are incompatible with life.**

• Blood plasma proteins act as buffers, mainly due to carboxyl groups and amino groups.



33

### Dissociation: Acid-base balance

**Buffers of boold plasma**

- Phosphates
- Plasma proteins

} **immediate effect but limited buffer capacity**

**Bicarbonate buffer**

The most important system for controlling the acid-base balance, because the body is able to actively change the concentration of [HCO<sub>3</sub><sup>-</sup>] (by **metabolic process**) and the concentration of [CO<sub>2</sub>] (by **respiration**) = the **delayed but long-term effect, due to connection with metabolism**

- **Bicarbonate**  $H_2O + CO_2 \rightleftharpoons H_2CO_3 \rightleftharpoons H^+ + HCO_3^-$

$pH = pK_A - \log \frac{[CO_2]}{[HCO_3^-]}$  **affected by metabolism**

- **Hemoglobin** (Hb released O<sub>2</sub> and bound H<sup>+</sup>; Hb bound O<sub>2</sub> in the lungs and released H<sup>+</sup>)

$$HbH \rightleftharpoons Hb^- + H^+ \quad \text{oxy-HbH} \rightleftharpoons \text{oxy-Hb}^- + H^+$$

34

### Acid–base imbalance

**Acidosis:** decrease of blood pH below the physiological value  
**Alkalosis:** increase of blood pH above the physiological value


- **Acid-base imbalance caused by inappropriate regulation of respiration**  
 Acidosis ( $\uparrow pCO_2$ ): hypoventilation (e.g. neuropathy), obstruction in the airway (inflammation, tumor), drugs (sedatives, hypnotics)  
 Alkalosis ( $\downarrow pCO_2$ ): hyperventilation (reflexively occurring in lung diseases), irritation of the respiratory center (e.g. stress, infection)
- **Acid-base imbalance caused by inappropriate metabolic regulation**  
 Acidosis ( $\downarrow [HCO_3^-]$ ): kidney damage, diabetic ketoacidosis, diarrhea  
 Alkalosis ( $\uparrow [HCO_3^-]$ ): high intake of alkaline substances, loss acids (e.g. vomiting)

The state of bicarbonate buffer (determination of pH,  $[HCO_3^-]$  and  $pCO_2$ ) can be used to clinically assess the state of the patient AB $\bar{B}$ .

35

### Hydrogen bond



- **Hydrogen bonds play an important role in physiological systems.**
- H-bonds have a much shorter lifetime than the covalent bonds (which are stable under physiological conditions); the lifetime of H-bonds between water molecules is about  $10^{-11}$  s.
- H-bonds are about 10 times weaker than a covalent bonds.
- **Some structures, such as DNA, have many hydrogen bonds, which are mutually supportive => long-term stability of the whole structure.**



37

### Hydrogen bond

- Hydrogen bonds influence the structure of biopolymers.
- H-bonds may be **part of the internal structure** of the molecule, or contribute to **intermolecular binding**.
- The intermolecular H-bonds also contribute to the formation and **stability of the molecular complexes** (e.g. vitamin C and noradrenaline form a bimolecular complex).




38

### Non-bonded interactions

The motions of the electron around any atom produce instantaneous dipoles that in turn induce the formation of other attraction dipoles in neighboring atoms.

**Dipole-dipole interaction:** formed by the interaction of **polar molecules**



#### Van der Waals forces

**Dipole-induced dipole interaction:** formed by the interaction of **polar and nonpolar molecules**




**London dispersion force:** formed by the interaction of **nonpolar molecules**. London interactions are caused by random fluctuations in electron density in an electron cloud, which induces dipoles in other neighboring molecules.


40

### Non-bonded interactions

**Hydrophobic interaction:** formed by the interaction **between the nonpolar parts of macromolecules in aqueous environments**. The importance for the stability of the conformation of biopolymers - e.g. **biological membrane**, stabilizing pairs of aromatic amino acids.





**$\pi$ - $\pi$  interaction:** the **ring structures** have delocalized  $\pi$ -electrons produced by conjugated bond systems in the rings. When the rings are parallel to one another,  $\pi$ - $\pi$  interactions occur between the p-orbitals. These interactions occur in supramolecular structures, like **DNA** (they provide substantial structural stability).



41

### Non-bonded interactions




**DNA**  
 $\pi$ - $\pi$  interactions (also called stacking forces) and H-bonds are important in nucleobase stacking within DNA molecules.


42

### Non-bonded interactions: Cell Membrane

**Non-bonded interactions between phospholipids in cell membrane** (hydrophobic interaction; partially Van der Waals forces)



**Cholesterol (stability of membranes)**




The hydroxyl group will associate with the phosphate head groups of phospholipids. The steroid rings and hydrocarbon chain will associate with the fatty acid tails of phospholipids with the same London attraction forces that hold the fatty acids together. The presence of cholesterol in our membranes = **protection of membranes from rupture.**

43

### Non-bonded interactions: Protein structure

- **The primary structure** – is the exact specification of its atomic composition and the **chemical (covalent) bonds** connecting those atoms.
- **The secondary structure** – the pattern of hydrogen bonds in a biopolymer (three-dimensional form of local segments of the biopolymers)
- **The tertiary structure** – the three-dimensional structure of the macromolecule.
- **The quaternary structure** – the arrangement of multiple molecules in a multi-subunit complex (e.g. proteins, DNA).



44

### Biophysical interfaces


In physiological systems, interfaces occur in two areas, at the **air-tissue surface in the lungs**, and at **cell membranes**.

**Surface tension**


At liquid-air interfaces, surface tension results from the **greater attraction of liquid molecules to each other** (due to cohesion) than to the molecules in the air (due to adhesion).

$$\sigma = F / l \quad [N.m^{-1}]$$

$\sigma$  tends to reduce the surface of phase (droplet formation, surface wetting).





The **interfacial tension** of the solid-liquid interface: affects the wettability of the solid surface.



46

### Biophysical interfaces: The capillary action

**The capillary elevation** (good wettability of the solid surface) allows the water to rise into the upper parts of plants.

$$h = \frac{2\sigma}{r\rho g}$$

$\sigma$  – surface tension  
 $\rho$  – density  
 $r$  – capillary radius  
 $g$  – gravitational constant



$r = 0,5 \text{ mm} \Rightarrow h = \frac{2 \cdot 76 \cdot 10^{-3}}{0,5 \cdot 10^{-3} \cdot 1000 \cdot 10} = 3 \text{ cm}$

$r = 0,5 \mu\text{m} \Rightarrow h = \frac{2 \cdot 76 \cdot 10^{-3}}{0,5 \cdot 10^{-6} \cdot 1000 \cdot 10} = 30 \text{ m}$

47

### Biophysical interfaces: The micelles and lipid bilayer formation

**The critical micellar concentration:** Exceeding the the critical concentration of surfactant leads to formation of **micelles or lipid bilayer** (due to hydrophobic interactions).


The phospholipid bilayer form the building blocks of cell membranes

48


### Biophysical interfaces: The pulmonary surfactant

- Pressure in alveoli is directly proportional to surface tension; and inversely proportional to radius of alveoli (**law of Laplace**).
- Pressure in smaller alveolus would be greater than in larger alveolus, if surface tension were the same in both => **Small alveoli are unstable.**
- How can they be stabilized?  
**The pulmonary surfactants stabilize the alveoli.**
- Premature babies not yet created a surfactant in the lungs - they need help to inflate the lungs (respiratory equipment).

**Law of Laplace**  
 $P = 2 \cdot T / r$



$r = 2 \Rightarrow P = 2 \cdot T / 2 \mid P = T$   
 $r = 1 \Rightarrow P = 2 \cdot T / 1 \mid P = 2T$



50

### Biophysical interfaces: The pulmonary surfactant

Recall from Laplace's Law:  
 $P = \frac{2T}{r}$   
 $P_1 = \frac{2T}{r_1} = 2T$   
 $P_2 = \frac{2T}{r_2} = T$   
 $P_1 > P_2 \Rightarrow$  Small alveoli are unstable.

Alveoli without pulmonary surfactant:  
 $P_1 > P_2 \Rightarrow$  Small alveoli are unstable.

Recall from Laplace's Law:  
 $P = \frac{2T}{r}$   
 $P_1 = \frac{2 \cdot \frac{1}{2}T}{r_1} = T$   
 $P_2 = \frac{2 \cdot \frac{1}{2}T}{r_2} = T$   
 $P_1 = P_2 \Rightarrow$  Alveoli are stable.

Alveoli with pulmonary surfactant:  
 $P_1 = P_2 \Rightarrow$  Alveoli are stable.

51

### Dispersion

- a system in which particles (solid, liquid or gas) are dispersed in a continuous phase.
- any aqueous environment in the human body (e.g. blood) are dispersions (the extracellular and intracellular environment)

**Classification of disperse systems according to the particle size:**

- coarse dispersion (particle size 1  $\mu$ m to 1 mm)
- colloid dispersion (particle size 1  $\mu$ m to 1 nm)

Continuous medium	Dispersed phase	Coarse dispersion	Colloid dispersion
gas	gas	-	-
	liquid	Fog	aerosol
	solid	dust	aerosol
liquid	gas	foam	foam
	liquid	emulsion	sol
	solid	suspension	sol
solid	gas	solid foam	solid foam (aerogel)
	liquid	wet sponge	gel
	solid	solid mixture	solid sol

53

### Dispersion

Dispersed phase / Continuous medium	Coarse dispersion (1 $\mu$ m to 1 mm)	Colloid dispersion (1 nm to 1 $\mu$ m)	Analytical dispersion (less than 1 nm)
solid / liquid	blood (blood cells)	blood plasma	true solutions (electrolytes, non-electrolytes)
liquid/ liquid	milk (fat globules in solution of proteins)	solutions of macromolecules	mixtures of miscible liquids
<b>The properties</b>			
observability	light microscope	electron microscope	you can not observe individual particles
osmotic effect	no	small	big
diffusion	no	slow	fast
sedimentation	in the gravitational field	ultracentrifuge	no
optical properties	opaque	opalescence - due to the diffraction of light	clear

54

### Dispersion

Dispersed phase / Continuous medium	Coarse dispersion (1 $\mu$ m to 1 mm)	Colloid dispersion (1 nm to 1 $\mu$ m)	Analytical dispersion (less than 1 nm)
solid / liquid	<b>Oncotic pressure</b> is a form of osmotic pressure exerted by proteins (albumin, globulins, fibrinogen) – is relatively small but <b>significant</b> (basic physiological parameter of blood).		
liquid/ liquid	The diffusion of small molecules and ions from the colloidal solution through a <b>dialysis membrane</b> – is relatively small but <b>significant</b> (basic physiological parameter of blood). <b>Hemodialysis</b> – replacement of kidney function.		
<b>The properties</b>			
observability	<b>Sedimentation rate</b> of blood (erythrocytes) is basic laboratory tests - the first warning signal indicating some disease (blood sedimentation is accelerated especially in inflammations or infectious diseases).		
osmotic effect			
diffusion	Determination of <b>particle size</b> (laser diffraction) and <b>particle concentration</b> (turbidimetry, nefelometry).		
sedimentation			
optical properties	opaque	opalescence - due to the diffraction of light	clear

55

### Electrokinetic properties of colloid dispersion

**The electrical double layer**  
 The colloids particles carry on their surface **electric charge**, which affect the **stability** of the sol. The charge is formed e.g. by **dissociation** of groups on the particle surface.

**Zeta potential:** around the charged colloidal particles are grouped ions of opposite sign - charge neutralization. Part of the ion atmosphere is fixed to the particle (compact layer), part continuously passes into solution (diffuse part). There is a potential difference (zeta potential) the compact part and the diffuse part. **Zeta potential affects the stability of colloidal solutions.**

59

### Electrokinetic properties of colloid dispersion

**The isoelectric point**  
 The pH of the solution, in which the same number of basic and acidic groups is dissociated (positive and negative charges are equalized; the electrical double layer is canceled; zeta potential = 0; the colloidal system is not stable - **coagulation occurs**).

The graph shows the concentration of different species as a function of pH. At the isoelectric point (pI), the net charge is zero, and the system becomes unstable, leading to coagulation. The photo shows test tubes where the solution becomes turbid at the isoelectric point.

60