



MG 100 Archaeobotany seminar Handout: Archaeobotanical Taphonomy, for charred seed assemblages

Institute of Archaeology 2006
Dorian Q Fuller

This hand-out includes selected diagrams on crop processing

Minnis' taphonomic diagram, the 'catch-all' model

Dennell's general processing assemblage formation model

Hillman's activities and assemblages relationship diagram

cereal disarticulation (for wheats and barley).

G. Jones summary of processing affects of weed seed types, and statistical summary of separation of products and by-products

C. Stevens pictorial summary of key stages in crop-processing (glume wheats) [based on Hillman 1984 and G. Jones 1987]

Rice processing model from Harvey and Fuller (2005) based on Thompson (1996)

Hulled milled processing model from Harvey and Fuller (2005) based on Reddy (1997)

Pulse processing model from Fuller and Harvey (2006)

Diagram of potential variables affecting pre-depositional, depositional, post-depositional stages of charred assemblages

A synopsis of some of the key publications in the development of thinking on archaeobotanical formation processes with bibliography is included at the end.

SEED SOURCES




Figure 1. Sources of seeds recovered from archaeological sites.

Minnis (1981) the catch-all model of potential seed sources. Important distinction between intrusive modern and ancient material.




Figure 1. Suggested relationship between economic status of a plant, its archaeological context and sample composition. Width of line indicates degree of association between economic status and context.

Dennel (1976): common sense model of use type and archaeological context in relation to sample size and heterogeneity. Note the suggested relationship between larger samples and staples.




Figure 2. Two sets of relationships encountered on archaeological sites (from Hillman 1973)

Hillman's (1973) diagram of the relationships between source, context and assemblage composition.





Figure 3. The same relationships on present-day settlements: all are observable and measurable.

Hillman's (1973) diagram of observable relationships in ethnographic contexts: the role of ethnoarchaeology for archaeobotany.



The formation of an archaeobotanical (carbonized) sample in terms of patterning composition. The importance of considering routine activities and regular inputs to fire for preservation. (From Fuller, Stevens and McClatchie, in press).







Diagram of main categories of weed seeds expected in by-products of main processing stages (after G. Jones 1984; 1987).



Cartoon of the main crop-processing stages for glume wheats (from Stevens 2003): 1. Threshing. 2. Raking. 3. Winnowing. 4. Coarse sieving [note return of some by-products to threshing]. 5. Fine-sieving. 6. Pounding (de-husking). 7. Winnowing. 8. Coarse sieving (to return undehusked spikelet to previous step). 9. Fine sieving. 10. Hand-picking.



Schematic representation of main quantitative patterns through the course of crop-processing (from Fullerm Stevens and McClatchie, in press).



Simplified schema of rice processing (from Harvey and Fuller 2005) indicating products (top) and by-products (bottom). Potential macro-remains shown black, potential phytolith outputs shown in outline. Also below:




Fig. 2. Simplified outline of rice crop processing (based on Thompson [69]).





Fig. 4. Simplified outline of millet crop processing (based on Reddy, unpublished PhD dissertation, 1994).

Millet processing (also from Harvey and Fuller 2005).

Hulled vs. Free-threshing Cereals

Crops	Hulled cereals (requiring dehusking)	Free-threshing cereals
Wheats	Triticum monococcum Triticum diococcum	Triticum aestivum (including T. sphaerococcum) Triticum durum
Barley	Hordeum vulgare var. vulgare	Hordeum vulgare var. nudum
Rice	Oryza sativa	
Millets	Sorghum bicolor race bicolor S. bicolor race kafir	Sorghum bicolor race caudatum, race guinea, race durra
	Panicum miliaceum, P. sumatrense	Eleusine coracana
	Setaria italica, S. pumila, S. verticillata, etc.	Pennisetum glaucum (syn. P. americanum, P. typhoides)
	Brachiaria ramosa	
	Paspalum scrobiculatum	
	Echinochloa frumentacea	
	Digitaria spp.	

Tabular summary of pulse (legume) processing variables (from Fuller and Harvey 2006)

Model/process	variant	pulse taxa	effects	remarks
Harvesting	uprooting	<i>Macrotyloma</i> (Watts 1908, 506); <i>Vigna radiata</i> (Watts 1908, 200)	incorporates weeds, especially climbers	
	cutting near base	<i>Lablab</i> (Watts 1908: 510); <i>Cajanus</i> (Westphal 1974; Van der Maesen 1989); <i>V. acutifolia</i> (Joan Ceas 1989a); <i>Macrotyloma</i> (Jansen 1989)	incorporates weeds, especially climbers	
	plucking pods	<i>Lablab</i> (Shrivashankar & Kulkarni 1989; Duke 1991); <i>Vigna radiata</i> (Watts 1908, 200; Weber 1991, 98)	selects against weeds	more likely in Neolithic due to uneven ripening; skip down to coarse sieving or pounding and rewinnowing
Threshing	free-threshing	<i>Lablab</i> , <i>Vigna</i>	frees pulses from pods and plants	some pods will not shatter; threshing of the by-product can be repeated one or more times to increase seed recovery
	pod-threshing	<i>Macrotyloma</i> , <i>Cajanus</i>	separates pods from plant	in <i>Cajanus</i> leaves are stripped or separated by simple shaking
Winnowing and Raking	free-threshing types	<i>Lablab</i> , <i>Vigna</i>	separates light material including pod fragments; product includes pulse seeds, large and small weeds, pod pedicels(?)	skip pounding and rewinnowing (pod-threshing) step. By-product may be used as fodder. If some pods are insufficiently broken, threshing may be repeated.
	pod-threshing types	<i>Macrotyloma</i> , <i>Cajanus</i>	separates light material; product includes pods, large heavy weeds, headed weeds, stem pieces. Pulse seeds from broken pods may enter by-product	By-product may be used as fodder. Mature seeds may enter dung. Possible stored as pods after this step. Possibly stored as pods after this step
Coarse sieving	free-threshing types	<i>Lablab</i> , <i>Vigna</i>	removes plants stalk parts, weed heads. Will lose some pulse seeds, especially unshattered pods.	By-product may be used as fodder. Mature seeds may enter dung.
	pod-threshing types	<i>Macrotyloma</i> , <i>Cajanus</i> , some <i>Lablab</i> (?)	removes small and large weed seeds, pulse pods and weed heads remain (could be hand-picked)	
Fine Sieving	free-threshing types	<i>Lablab</i> , <i>Vigna</i>	removes remaining small weeds, chaff fragments. Only weeds very similar in size and weight to pulse remain, possibly some pod pedicels (especially in <i>Vigna</i>). Will lose some small/immature pulse seeds.	store after this step as cleaned pulses; sieved again or hand-picked to remove remaining large weeds before cooking. Possible route to archaeological preservation.
	pod-threshing types	<i>Macrotyloma</i> , <i>Cajanus</i>		this step probably skipped
Pounding and rewinnowing	free-threshing types	this step unnecessary		
	pod-threshing types only	<i>Macrotyloma</i> , <i>Cajanus</i> , some <i>Lablab</i> (?)	removes pods, only some weed seeds or heads that are very close in size and weight to pulse remain	possibly a daily routine processing; most likely route to archaeological preservation
Parching		parching or dry-roasting reported for <i>Vigna</i> spp., <i>M. uniflorum</i> (Watts 1908). <i>Lablab</i> reported to be 'dried' before storage (Shrivashankar & Kulkarni 1989)	could lead to accidental charring and archaeological preservation	parched before grinding, or dry-roasting for consumption. Archaeological preservation route

Major publications in the development of archaeobotanical formation processes (for charred assemblages, especially from agricultural periods).

Körber-Grohne (1967, in German; 1981 in English [synoptic article]) discusses spatial distribution of archaeobotanical remains in relation structural features and potential processing and consumption activities.

Knörzer (1971, in German) observes that most charred assemblages includes cereals and herbaceous plant many of which are known field weeds. Thus most assemblages derive from arable ecosystems and weed species present can be used to infer aspects of agriculture. Also draws on Körber-Grohne's approach.

Dennell (1972) observes differences in composition of archaeobotanical assemblages between contexts, including the size of cereal grains. Carries out sieving experiments to confirm that past sieving of crops, as part of processing, may have contribute to the formation of distinct product/by-product assemblages. Subsequently, Dennell (1974; 1976) develops a general predictive model for comparison and interpretation of assemblages related to assumptions about the affecting of major crop processing stages on their composition. Hubbard (1976) disputes the statistical validity and assumptions of Dennell's model.

Hillman (1973) outlines need for ethnographic observation to produce generalisations that can be used to interpret archaeobotanical evidence

Hillman (1981; 1984) published crop-processing model on ethnographic work and turkey. Discusses its implication for interpreting archaeobotanical remains in terms of crop husbandry and producer/consumer sites.

G. Jones (1984; 1987) published similar processing sequence observed independently in Greece and discusses the potential of statistical analysis of weed seed types for distinguishing processing stages.

M. Jones (1985) applied elements of crop processing reasoning to question of distinguishing produce vs. consumer sites (but via method opposite Hillman 1981. See review in Van der Veen 1992). Also, a brief history of archaeobotanical interpretation in Europe.

Mikcesek (1987) general review article. Broad perspective but with less interest in crop processing, and a somewhat mistaken summary of early Dennell and Hillman work. Hastorf (1988) reviews importance of crop processing and highlights need for similar studies of New World crop species.

Boardman and Jones (1990) experimental charring demonstrating the much greater likelihood of chaff elements to be destroyed by charring, while grain are preserved, thus majorly biasing pre-charring ratios of chaff to grain.

Experimental work on processing, and charring maize reported by Goette, Williams, Johannessen and Hastorf (1994)

Reddy (1991; 1997; 2003) applies Hillman-Jones approach to Indian millets, with ethnographic and archaeological study.

Thompson (1996) applied Hillman-Jones approach to rice processing in Thailand

D'Andrea et al. (1999); Butler et al. (1999) preliminary report of ethnoarchaeological study of crop processing in Ethiopia, including tef and pulses.

Stevens (2003) develops the importance of routine processing (as the main source of waste) and considers how this allows contrasts to be interpreted in terms of labour mobilization. While recently Van Der Veen and Jones (2006) have critiqued this, they seem to have misunderstood Stevens and arrive at very nearly the same conclusion!

Application to phytolith analysis (for rice and millets) by Harvey and Fuller (2006)

A model of the variables in pulse processing, distinguishing pod-threshing and free-threshing pulse types (Fuller and Harvey 2006)

Selected Bibliography of Archaeobotanical Taphonomy

- Boardman, S. and G. E. M. Jones 1990. Experiments on the effects of charring on cereal plant components, *Journal of Archaeological Science* 17(1): 1-12
- Bottema, S. 1984. The composition of modern charred seed assemblages, in *Plants and Ancient Man* (W. van Zeist and W. A. Casparie eds.), pp. 207-212. Rotterdam: Balkema.
- Bruno M and Whitehead WT. 2003. Chenopodium cultivation and formative period agriculture at Ciripa, Bolivia. *Latin American Antiquity* 14: 339-355.
- Butler, A. 1992. Pulse agronomy: traditional systems and implications for early cultivation, in *Préhistoire L'Agriculture: Nouvelle Approches Expérimentales et Ethnographiques* (ed. P. C. Anderson), pp 67-78. Monographie du CRA 6, Editions du CNRS, Paris.
- Butler, A., Z. Tesfay, C. D'Andrea and D. Lyons 1999. The Ethnobotany of *Lathyrus sativus* L. in the Highlands of Ethiopia, in *The Exploitation of Plant Resources in Ancient Africa* (M. Van Der Veen ed.), pp. 123-136. New York: Kluwer Academic/ Plenum.
- Charles, Michael 1998. Fodder from Dung: the Recognition and Interpretation of Dung-Derived Plant Material from Archaeological Sites, *Environmental Archaeology* 1: 111-122
- D'Andrea, C., D. Lyons, M. Haile and A. Butler 1999. Ethnoarchaeological Approaches to the Study of Prehistoric Agriculture in the Highlands of Ethiopia, in *The Exploitation of Plant Resources in Ancient Africa* (M. Van Der Veen ed.), pp. 101-122. New York: Kluwer Academic/ Plenum.
- Dennell, R. W. 1972. The Interpretation of Plant Remains: Bulgaria, in *Papers in Economic Prehistory* (E. S. Higgs ed.), pp. 149-159. Cambridge: Cambridge University Press
- Dennell, R. W. 1974. Botanical evidence for prehistoric crop processing activities, *Journal of Archaeological Science* 1: 275-284
- Dennell, R. W. 1976. The economic importance of plant resources represented on archaeological sites, *Journal of Archaeological Science* 3: 229-247
- Fuller, D. Q. and Emma L. Harvey 2006. The archaeobotany of Indian Pulses: identification, processing and evidence for cultivation. *Environmental Archaeology* 11(2): 219-246
- Fuller, D.Q. and Chris J. Stevens. in press. Agriculture and the Development of Complex Societies: An Archaeobotanical Agenda. In *Ethnobotanist of Distant Pasts: Essays in Honour of Gordon Hillman*. Andrew Fairbairn and Ehud Weiss (eds.). Oxbow Books
- Fuller, D. Q., Chris J. Stevens and Meriel McClatchie. in press. Routine Activities, Tertiary Refuse and Labor Organization: Social Inference from Everyday Archaeobotany. In *Ancient Plants and People. Contemporary Trends in Archaeobotany*. Edited by Marco Madella and Manon Savard. University of Arizona Press, Tucson
- Goette, Williams, Johannessen and Hastorf (1994). Towards reconstructing ancient maize: experiments in processing and charring, *Journal of Ethnobiology* 14(1): 1-21
- Green, F. J. 1985. Plant remains in the modern soil and contamination from the surface, *East Anglian Archaeology* 27: 41-42
- Harvey, E. L. and D. Q. Fuller. 2005 Investigating crop processing through phytolith analysis: the case of rice and millets, *Journal of Archaeological Science* 32: 739-752
- Hillman, G. C. 1973. Crop husbandry and food production: modern models for the interpretation of plant remains, *Anatolian Studies* 23: 241-244
- Hillman, G. C. 1981. Reconstructing Crop Husbandry Practices from Charred Remains of Crops, in *Farming Practice in British Prehistory* (R. Mercer ed.), pp. 123-161. Edinburgh: University Press.

- Hillman, G. C. 1984. Interpretation of archaeological plant remains: The application of ethnographic models from Turkey, in *Plants and Ancient Man - Studies in Paleoethnobotany* (W. Van Zist and W. A. Casparie eds.), pp. 1-41. Rotterdam: A.A. Balkema.
- Hubbard, R. N. L. B. 1976. On the strength of the evidence for prehistoric crop processing activities, *Journal of Archaeological Science* 3: 257-265
- Hubbard, R. N. L. B. and A. Clapham 1992. Quantifying macroscopic plant remains, *Review of Palaeobotany and Palynology* 73: 117-132
- Jones, G. E. M. 1984. Interpretation of archaeological plant remains: Ethnographic models from Greece, Pp. 42-61 in W. Van Ziest and W. A. Casparie (eds.) *Plants and Ancient Man - Studies in Paleoethnobotany* . Rotterdam: A.A. Balkema.
- Jones, G. E. M. 1987. A statistical approach to the archaeological identification of crop processing, *Journal of Archaeological Science* 14: 311-323
- Jones, Martin K. 1985. Archaeobotany beyond subsistence reconstruction, in *Beyond Domestication in Prehistoric Europe* (G. W. Barker and C. Gamble eds.), pp. 107-128. Academic Press, New York and London.
- Keepax, C. 1977. Contamination of Archaeological Deposits by Seeds of Modern Origin with Particular Reference to the Use of Flotation Machines, *Journal of Archaeological Science* 4: 221-229
- Knörzer, K.-H. 1971. Urgeschichtliche Unkrauter im Rheinland. Ein Beitrag zur Entstehung der Segetalgesellschaften, *Vegetatio* 23: 89-111.
- Körber-Grohne, U. 1967. *Feddersen Wierde: die Ergebnisse der Ausgrabung der vorgeschichtlichen Wurt bei Bremerhaven in den Jahren 1955-1963* (Band 1). Wiesbaden.
- Körber-Grohne, U. 1981. Crop husbandry and environmental change in the Feddersen Wierde, near Bremerhaven, northwest Germany, in *The environment of man: the Iron Age to the Anglo-Saxon Period* (M. K. Jones and G. W. Dimbleby eds.), pp. 287-307. BAR British series 87. Oxford.
- Krueze, A 1990. Searching for "Single-activity refuse" in Linearbandkeramik Settlements. An Archaeobotanical Approach, in *Experimentation and Reconstruction in Environmental Archaeology* (D. E. Robinson ed.), pp. 63-75. Oxford: Oxbow Books
- Lundström-Baudais, K., A. M. Rachoud-Schneider, D. Baudais, and B. Poissonnier. 2002. Le broyage dans la chaîne de transformation du millet (*Panicum miliaceum*): outils, gestes et écofacts. In *Moudre et Broyer: I Méthodes*. Edited by H. Procopiou and R. Treuil, pp. 155-180. Paris: Comité des Travaux Historiques et Scientifiques.
- Miksicek, C. 1987. Formation Processes of the Archaeobotanical Record, *Advances in Archaeological Method and Theory* 10: 211-247
- Miller, N. F. 1984. The use of dung as fuel: an ethnographic example and an archaeological application, *Paleorient* 10(2): 71-79
- Minnis, Paul 1981. Seeds in Archaeological Sites: Sources and Some Interpretive Problems, *American Antiquity* 46: 143-151
- Murray, Mary Anne (2000) Cereal production and processing, in *Ancient Egyptian Materials and Technology* (ed. P. T. Nicholson and I. Shaw), pp. 505-536. Cambridge: Cambridge University Press
- Peña-Chocarro, Leonor (1999) *Prehistoric Agriculture in Southern Spain during the Neolithic and the Bronze Age. The Application of Ethnographic Models*. BAR International series 818, Oxford.

- Reddy, Seetha N. 1991. Complementary Approaches to Late Harappan Subsistence: An Example from Oriyo Timbo, Pp. 127-135 in R. H. Meadow (ed.) *Harappa Excavations 1986-1990 - A Multidisciplinary Approach to Third Millennium Urbanism*. Madison, Wisconsin: Prehistory Press
- Reddy, Seetha N. 1994. *Plant Usage and Subsistence Modeling: An Ethnoarchaeological Approach to the Late Harappan of Northwest India*, PhD Dissertation, University of Wisconsin. Ann Arbor Michigan: University Microfilms.
- Reddy, Seetha N. 1997. If the threshing floor could talk: integration of agriculture and pastoralism during the Late Harappan in Gujarat, India, *Journal of Anthropological Archaeology* 16: 162-187
- Reddy, Seetha N. 2003. *Discerning Palates of the Past: An Ethnoarchaeological Study of Crop Cultivation and Plant Usage in India*. Ann Arbor: Prehistory Press.
- Stevens, C. J. 2003. An investigation of agricultural consumption and production models for prehistoric and Roman Britain. *Environmental Archaeology* 8:61-76..
- Thompson, G. B. 1996. Ethnographic Models for Interpreting Rice Remains, in *The Excavation of Khok Phanom Di a prehistoric site in Central Thailand. Volume IV: Subsistence and Environment: The Botanical Evidence (The Biological Remains, Part II)* (C. Higham and R. Thosarat eds), pp.119-150. London: The Society of Antiquaries of London
- Van der Veen, M. 1985. Carbonised seeds, sample size and on-site sampling, in *Palaeoenvironmental Investigations. Research Design, Methods and Data Analysis* (N. R. J. Fieller, D. D. Gilbertson, and N. G. A. Ralph eds.), pp. 165-174. Oxford: BAR International Series 258.
- Van der Veen, M. 1992. *Crop Husbandry Regimes. An Archaeobotanical Study of Farming in northern England 1000 BC- AD 500*. Sheffield Archaeological monographs 3. J. R. Collis Publications, University of Sheffield.
- Van der Veen, M. 1999. The economic value of chaff and straw in arid and temperate zones, *Vegetation History and Archaeobotany* 8: 211-224
- Van der Veen, M. and N. Fieller 1982. Sampling seeds, *Journal of Archaeological Science* 9: 287-298
- Van der Veen, M. and G. Jones 2006. A re-analysis of agricultural production and consumption: implications for understanding the British Iron Age, *Vegetation History and Archaeobotany* 15(3): 217-228
- Viklund, K. 1998. *Cereals, Weeds and Crop Processing in Iron Age Sweden*, *Archaeology and Environment* 14. Umeaa: Department of Archaeology, University of Umeaa
- Wilson, D. G. 1984. The carbonisation of weed seeds and their representation in macrofossil assemblages, *Plants and Ancient Man - Studies in Paleoethnobotany* (W. Van Zist and W. A. Casparie eds.), pp. 201-210. Rotterdam: A.A. Balkema.
- Young, Ruth 1999. Ethnobotanical observations: Finger millet processing in east Africa, *Vegetation History and Archaeobotany* 8: 31-34