



MASARYKOVA UNIVERZITA
PŘÍRODOVĚDECKÁ FAKULTA



Obecná biologie – Fyziologie živočichů

Oddělení fyziologie a imunologie živočichů

Fyziologie: studium funkcí

Imunologie: studium obranyschopnosti

Srovnávací a ekologický přístup:

- živočichové včetně člověka
- širší mezioborový přístup než na LF


Výchova k vědecké práci:

- využívání informačních zdrojů
- experimentální laboratorní praxe
- hodnocení výsledků
- prezentace vlastních výsledků

Uplatnění:

- **Biomedicínský základní výzkum (AVČR, Univerzity, ústavy)**
- **Klinická pracoviště a laboratoře (humánní, veterinární)**
- **Zemědělsky orientovaný výzkum**
- **Soukromé firmy**

Skupina imunologie hmyzu



Pavel Hyršl



Oddělení fyziologie a imunologie živočichů

Ústav experimentální biologie

Přírodovědecká fakulta

Masarykova Univerzita v Brně

<http://www.sci.muni.cz/ofiz/>

IMUNITNÍ SYSTÉM HMYZU

BUNĚČNÁ IMUNITA

- pohyblivé krevní buňky – hemocyty
(počet a aktivita hemocytů je modulována humorálními faktory a neuroendokrinním systémem)

- koagulace
- fagocytóza (oxidativní vzplanutí?)
- nodulace
- enkapsulace

HUMORÁLNÍ IMUNITA

- molekuly v hemolymfě
- antigenem indukované antibakteriální a regulační faktory

- aglutininy
- koagulační kaskáda
- fenoloxidásová kaskáda
- baktericidní peptidy
- antimikrobiální faktory (lektiny, hemolisin)
- lysozym


stresové faktory

Hmyzí hemocyty:

- prohemocyty
- plasmatomy
- granulocyty
- spherulocyty
- oenocyty


Imunitní reakce hemocytů:

- fagocytóza
- enkapsulace
- nodulace
- koagulace



Obr. 1 : Navržené schema působení indukovaných antibakteriálních proteinů a peptidů na buněčnou stěnu bakterií (Hultmark 1993)

Figure 5: Diagram of an insect granular cell emphasizing its multifunctional role.



Elektroforéza SDS – PAGGE

(sodium dodecylsulfat polyacrylamide gradient gel electrophoresis)

- gely vznikají polymerací akrylamidu a N',N'-metylenbisakrylamidu
- SDS (sodium dodecylsulfát) se váže shodně na všechny proteiny v poměru 1,4 g / 1g proteinu a předává jim silný záporný náboj
- mercaptoethanol - rozštěpí disulfidické můstky v molekulách proteinů
- vertikální elektroforéza SE 600 (Hoefer Scientific Instruments)
- 5 % koncentrační gel a separační gel (14 x 10,5 cm) s gradientem akrylamidu 7,5 – 20,0 %
- dělení proteinů probíhalo ve dvou gelech, každý z nich pro 15 vzorků
- standard (SDS-PAGE Molecular Weight Standard Broad Range, Bio-Rad; 6,5 - 200 kDa)
- gely byly barveny stříbrem podle Kirkeby et al. (1993)

Vyhodnocování gelů:

- videodensitometr GS-670 a software Molecular Analyst (Bio-Rad)

Profilová analýza:

- **kvalitativní:**



MW jednotlivých proteinových frakcí hemolymfy

- **kvantitativní:**

poměr zastoupení jednotlivých proteinových frakcí
mezi samci a samicemi během vývoje

Profilová analýza:

- rozdělené vzorky na polyakrylamidovém gelu
- absorpční křivka (osa x – vzdálenost od startu, osa y – optická hustota)
- tabulka s určenou molekulovou hmotností frakcí (MW) a plochou píku




Peak No.	Area (OD*mm)	MW (Da)
1	0.1031	84013
2	0.1521	79290
3	0.4767	66371
4	0.5198	64548
5	0.3230	61972
6	0.8325	59245
7	0.1905	56035
8	0.2044	52885
9	0.3650	50993
10	0.3637	48126
11	0.2294	45910
12	0.1872	45130
13	0.4452	44363
14	0.1340	42593
15	0.2568	40719
16	0.4203	39515
17	0.1859	38265
18	0.1684	36739

**Proteinové spektrum hemolymfy
bource morušového (*Bombyx mori*)**

Bombyx mori (Lepidoptera, Bombycidae)

bulharský monovoltinní hybrid AS x KK






Životní cyklus bource morušového *B. mori*, upraveno podle Lim et al., 1990.

Bombyx mori – samice

S – standard; 1. - 9. den V. instaru; PR – prepupa 1. den;
Z – kukla začátek; S – kukla střed; K – kukla konec.



**Proteinové spektrum hemolymfy
zavíječe voskového (*Galleria mellonella*)**

Galleria mellonella (Lepidoptera, Pyralidae)



RESULTS

Protein Profiles







Fig. 1. Acrylamide gradient gel of proteins extracted from last instar larvae (samples 1-5) and pupae (samples 6-9) hemolymph of *G. mellonella*. Samples with increasing concentration of boric acid: 0.00 ppm (1, 6); 156 ppm (2, 7); 620 ppm (3, 8); 1250 ppm (4, 9) and 2500 ppm (5). Main protein groups of proteins are described on the left margin as well as 45 kDa protein fraction according to protein standards (S).

Aktivita lysozymu v hemolymfě



radiální difúze v agaróze

Silkworm (*Bombyx mori*) Hemocytes Do Not Produce Reactive Oxygen Metabolites As a Part of Defense Mechanisms

P. HYRŠL^a, M. ČÍŽ^b, L. KUBALÁ^b, A. LOJEK^{b*}

^aDepartment of Comparative Animal Physiology and General Zoology, Faculty of Science, Masaryk University, 611 37 Brno, Czechia
fax +420 541 211 293
e-mail alojek@ibp.cz

^bInstitute of Biophysics, Academy of Sciences of the Czech Republic, 612 65 Brno, Czechia

Received 6 October 2003
Revised version 22 December 2003

ABSTRACT. To investigate whether hemocytes of *Bombyx mori* (Lepidoptera) larvae produce reactive oxygen species (ROS) as part of the oxidative killing of invading pathogens, the production of ROS was measured as a luminol- and lucigenin-enhanced chemiluminescence of unstimulated or stimulated (zymosan particles, phorbol myristate acetate, calcium ionophore, rice starch or *Xenorhabdus nematophila*) hemolymph. No detectable ROS production was found. The spontaneous and activated ROS production measured with hemocytes, i.e. under the conditions when the antioxidative potential of hemolymph plasma was eliminated, was again undetectable. Likewise, ROS production by isolated hemocytes was observed by spectrophotometric (NBT test, cytochrome *c* assay) and fluorimetric (using dihydrorhodamine and hydroethidine probes) methods. Hence none of the experimental approaches used indicated the production of ROS by hemocytes of *B. mori* larvae as part of their immune response.

Hemocytes are basic to the invertebrate innate immune system that is divided into cellular and humoral defense responses. The most common types of hemocytes reported in the literature are prohemocytes, plasmatocytes, granulocytes, spherulocytes and encytes (Yamashita and Iwabuchi 2001; Lavine and Strand 2002). These have been identified by their morphology and histochemical and functional reactions (e.g., Gardiner and Strand 1999). Four basic types of hemocyte immune reactions have been described: phagocytosis, encapsulation, nodulation and coagulation. These activities are always connected with a particular type of hemocytes. During phagocytosis, plasmatocytes and granulocytes are mainly activated while other types of hemocytes have mostly no possibility to phagocytose. Mechanisms participating in the recognition of foreign material are still under study (Lavine and Strand 2002).

The production of reactive oxygen species (ROS) seems to be an important microbicidal factor in both invertebrate hemocytes and vertebrate phagocytes since an increase in ROS production by activated hemocytes of some invertebrates such as *Bivalvia*, *Clitellata*, *Malacostraca*, *Arachnida*, *Echinoidea* or *Ascidacea* has been reported (e.g., Nakamura *et al.* 1985; Ito *et al.* 1992; Bell and Smith 1993; Valembois and Lassegues 1995; Lambert and Nicolas 1998; Ordas *et al.* 2000; Pereira *et al.* 2001; Azumi *et al.* 2002). However, reports about similar mechanisms in insects are controversial. The aim of the present study was to investigate whether stimulated hemocytes of *Bombyx mori* can produce ROS as part of oxidative killing invading pathogens. Understanding of the defense mechanisms of *B. mori* has a high significance for providing a better insight into the evolution of animal immune systems; it also has a great impact on silk production sericulture.

MATERIALS AND METHODS

Sample preparation. The hemolymph of the Japan polyvoltine NO2 × CO2 hybrid of the silkworm, *Bombyx mori* LINNÉ 1758 (Lepidoptera, Bombycidae), was obtained from larvae of the 5th instar. Larvae were reared on mulberry leaves (*Morus alba*) *ad libitum*. The sex was not determined. After the collection of the hemolymph from the first-pair proleg (approximately 150 µL per larva), phenylthiourea was added to protect the hemolymph from melanization. (It was verified that this agent does not have a significant effect

*Corresponding author.

Reactive oxygen metabolites detection:

- chemiluminescence
- spectrophotometry
 - NBT test
 - cytochrome c reduction
- fluorometry
 - DHR 123 staining
 - HE staining
- total radical-trapping antioxidative potential (TRAP)

Entomopatogenní hlístovky (EPN)

třída: Nematoda (Nematoda)

řád: Hád'ata (Anguilata)

čeleď: *Steinernematidae*

Heterorhabditidae


- vyskytují se volně v půdě
- selektivita - výhradně entomopatogenní
- nízké teplotní optimum - do 20°C
- využívají se jako prostředek biologického boje



dospělci

invazní L3 larvy

Kultivace hlístovek na náhradním hostiteli:



Produktcode: 4-1273

nematop[®]

Dickmaulrüsslerbekämpfung



nema-green[®]

Biologische Rasenpflege



www.nema.de
Kontakt: E-Mail: info@nema.de, Tel.: 04322 44000, www.nema.de

Produktcode: 4-1272

nemaplus[®]

Trauermückenbekämpfung



METODY

1. Přirozená invaze EPN:

- larvy *G. mellonella* nebo larvy *Amphimallon* sp., *Hoplia* sp. + 25 EPN, 10% vlhkost
- inkubace 20 – 24 hodin za laboratorní teploty
- celkem bylo použito 70 ks *G. mellonella*, 23 ks *Amphimallon* sp. a 23 ks *Hoplia* sp.



Témata bakalářských a diplomových prací týkajících se entomopatogenních hlístovek jsou zaměřena na:

- testy patogenicity jednotlivých EPN
- srovnání invaze EPN do hostitele v různých prostředích
- obrana hostitele vůči EPN
- možnosti biologického boje s hmyzími škůdci

Výsledkem jsou:

- bakalářské a diplomové práce
- postery a přednášky na konferencích
- publikace

Folia Microbiol. 49 (3), 315–319 (2004) <http://www.bionetd.cas.cz/foia/E0139/>

Silkworm (*Bombyx mori*) Hemocytes Do Not Produce Reactive Oxygen Metabolites As a Part of Defense Mechanisms

P. HYŘŠL¹, M. ČÍŽ¹, L. KUBALA², A. LOJEK^{2*}

¹Department of Comparative Animal Physiology and General Zoology, Faculty of Science, Masaryk University, 611 37 Brno, Czechia
fax: +420 541 211 230
e-mail: alojek@biop.cz

²Institute of Biophysics, Academy of Sciences of the Czech Republic, 612 85 Brno, Czechia

Received 6 October 2003
Revised version 21 December 2003

ABSTRACT. To investigate whether hemocytes of *Bombyx mori* (*Lepidoptera*) larvae produce reactive oxygen species (ROS) as part of the oxidative killing of invading pathogens, the production of ROS was measured as a luminal- and lucigenin-enhanced chemiluminescence of unstimulated or stimulated (zymosan particles, phorbol myristate acetate, calcium ionophore, rice starch or *Acanthobolus nematophilus*) hemolymph. No detectable ROS production was found. The spontaneous and activated ROS production measured with hemocytes, i.e. under the conditions when the antioxidative potential of hemolymph plasma was eliminated, was again undetectable. Likewise, ROS production by isolated hemocytes was observed by spectrophotometric (NBT test, cytochrome c assay) and fluorimetric (using dihydroethidine and hydrochloride probes) methods. Hence none of the experimental approaches used indicated the production of ROS by hemocytes of *B. mori* larvae as part of their immune response.


Hemocytes are basic to the invertebrate innate immune system that is divided into cellular and humoral defense responses. The most common types of hemocytes reported in the literature are prohemocytes, plasmatocytes, granulocytes, spherulocytes and encystes (Yamashita and Iwabuchi 2001; Lavine and Strand 2002). These have been identified by their morphology and histochemical and functional reactions (e.g. Gardiner and Strand 1999). Four basic types of hemocyte immune reactions have been described: phagocytosis, encapsulation, nodulation and coagulation. These activities are always connected with a particular type of hemocytes. During phagocytosis, plasmatocytes and granulocytes are mainly activated while other types of hemocytes have mostly no possibility to phagocytose. Mechanisms participating in the recognition of foreign material are still under study (Lavine and Strand 2002).

The production of reactive oxygen species (ROS) seems to be an important microbicidal factor in both invertebrate hemocytes and vertebrate phagocytes since an increase in ROS production by activated hemocytes of some invertebrates such as *Bivalvia*, *Ciliolata*, *Malacostraca*, *Arachnida*, *Echinoidea* or *Ascidacea* has been reported (e.g. Nakamura et al. 1985; Ito et al. 1992; Bell and Smith 1993; Valenbois and Lassgues 1995; Lambert and Nicolas 1998; Ordás et al. 2000; Pereira et al. 2001; Azumi et al. 2002). However, reports about similar mechanisms in insects are controversial. The aim of the present study was to investigate whether stimulated hemocytes of *Bombyx mori* can produce ROS as part of oxidative killing invading pathogens. Understanding of the defense mechanisms of *B. mori* has a high significance for providing a better insight into the evolution of animal immune systems; it also has a great impact on silk production technology.

MATERIALS AND METHODS

Sample preparation. The hemolymph of the Japan polyvoltinism NO2 × CO2 hybrid of the silkworm, *Bombyx mori* LepS: 1758 (*Lepidoptera*, *Bombycidae*), was obtained from larvae of the 5th instar. Larvae were reared on mulberry leaves (*Morus alba* L.) *ad libitum*. The sex was not determined. After the collection of the hemolymph from the first pair proleg (approximately 150 µL per larva), phenylthiourea was added to protect the hemolymph from melanization. (It was verified that this agent does not have a significant effect

*Corresponding author.



Luminometer 2004, 10: 41–43
Published online in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/lum.755

ORIGINAL RESEARCH

Chemiluminescence of lucigenin is dependent on experimental conditions

Pavel Hyřšl¹, Antonín Lojek^{2*}, Milan Číž² and Lukáš Kubala²

¹Department of Comparative Animal Physiology and General Zoology, Faculty of Science, Masaryk University, Kotlářská 2, Brno 611 37, Czech Republic
²Institute of Biophysics, Academy of Sciences of the Czech Republic, Královopolská 135, Brno 612 85, Czech Republic

Received 15 December 2002; revised 8 December 2003; accepted 7 January 2004

ABSTRACT. The aim of the study was to test the effect of experimental conditions such as light irradiation and temperature on chemiluminescence (CL) of 10^{-6} – 10^{-7} mol/L lucigenin dissolved in various types of solvents. Irradiation by UV light (280, 297, 313 or 400 nm) induced a significant increase in CL of lucigenin dissolved in borate buffer. This effect was the most obvious for 10^{-6} – 10^{-7} mol/L lucigenin. All wavelengths used had a similar effect. UV irradiation did not induce changes in the CL activity of lucigenin dissolved in H_2O or in dimethyl sulphoxide (DMSO). Different results for various solvents were not dependent on pH. The CL activity of 10^{-6} mol/L and 10^{-7} mol/L lucigenin dissolved in borate buffer increased depending on the solution temperature (25°C, 30°C, 37°C or 40°C) already at the beginning of the analysis, with a further increase during 16 h incubation period. It can be summarized that temperatures higher than 25°C and intensive light irradiation are among those factors which significantly affect the result of analysis when lucigenin is used as a luminophore. Copyright © 2004 John Wiley & Sons, Ltd.

KEYWORDS: chemiluminescence; lucigenin; light irradiation; temperature

INTRODUCTION

Lucigenin has frequently been used in the luminescent detection of superoxide anion radical (O_2^-) production by various enzymatic and cellular systems (1–3). The three key steps for detecting O_2^- are: (a) one electron reduction of lucigenin (Luc^2) to cation radical ($\text{Luc}^{\bullet+}$); (b) reaction with O_2^- to yield lucigenin diacetate; and (c) the decomposition of lucigenin diacetate to electronically excited N-methylacridone which emits photons when related to ground state (1).

The auto-oxidation of $\text{Luc}^{\bullet+}$ with a simultaneous production of O_2^- can be a complicating factor (3, 4) and thus experimental conditions can significantly affect the amount of O_2^- detected. We observed an unexpected excitation of lucigenin in various types of previous experiments (see Results and Discussion). Therefore, the aim of this study was to test the effect of experimental conditions, such as temperature and light irradiation, on chemiluminescence (CL) of lucigenin dissolved in various types of solvent.

MATERIALS AND METHODS

Reagents

Solutions of 10^{-6} – 10^{-7} mol/L lucigenin (bis-N-methylacridinium nitrate, Sigma) were prepared in three types of solvents: sodium borate buffer, pH 9.0 (1.24 g H_2BO_3 and 763 g $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$ in 500 mL distilled water, both chemicals from Fluka), dimethyl sulphoxide (DMSO, Sigma) or distilled H_2O . All solutions were stored at -20°C before use. All chemicals used were of analytical grade.

Chemiluminescence assay

Lucigenin solutions or solvents alone were pre-incubated for 15 min to raise them to the required temperature (25°C, 30°C, 37°C or 40°C). CL emission of samples (reaction volume 200 µL) was measured in triplicates in 10 min intervals for 16 h using BioOrbit 1251 luminometer (Bio-Orbit, Finland). The data are presented in mV.

The effect of radiation

To test the effect of radiation, 1 mL 10^{-6} mol/L lucigenin was irradiated in plastic dishes at room temperature with UV light (280, 297, 313 and 400 nm) generated by the Oriol Solar Simulator for 10 min. The dose was

*Correspondence to: A. Lojek, Institute of Biophysics, Academy of Sciences of the Czech Republic, Královopolská 135, Brno 612 85, Czech Republic.
E-mail: alojek@biop.cz
Contract/grant sponsor: Grant Agency of the Czech Republic; Contract/grant number: 524/03/009.