

**Hraniční plazmolýza***Allium cepa*

molární koncentrace (mol/l)	0	0.1	0.2	0.3	0.4	0.5
	5	0	5	10	10	20
	0	0	10	30	30	50
	0	0	0	20	20	25
	0	0	10	20	20	40
	0	0	10	10	10	20
	0	0	10	20	20	40

**Hraniční plyzmolýza***Elodea canadensis*

molární koncentrace	0	0.1	0.2	0.3	0.4	0.5
	0	20	20	30	30	100
	0	0	20	50	50	50
	0	10	30	50	40	60
	0	10	20	40	30	50
	0	0	20	40	40	60
	0	10	30	50	50	60
	0	0	0	30	30	60

**Tlaková metoda**

Tlak

nezalitá rostlina		zalitá rostlina		nezalitá rostlina, list 10	
I. List	II. List	I. List	II. List	I. List	II. List
6	5	4	5	7	6
6	6	3.5	4	6	7
5.5	6.5	4.5	3.5	6	6.5

**Refraktometrie**

molární koncentrace (mol/l)	0.1	0.2	0.3	0.4	0.5	0.6
kalibrace	4	7.6	10.6	14	17	20.6
<i>Solanum tuberosum</i> (A)	5.6	6.9	10.2	11.2	15.3	16.9
<i>Solanum tuberosum</i> (B)	7	8	10.9	12.8	14.4	15.4

0.6	0.7	0.8
50	100	90
75	100	100
25	30	40
70	75	90
30	100	100
70	80	100

0.6	0.7	0.8
40	50	60
80	100	100
70	100	100
70	90	100
80	90	100
80	100	100
60	70	100

)min odříznutý

0.7	0.8
23.4	26
18.2	19.5
17.6	18.6