

The Histidine Kinases CYTOKININ-INDEPENDENT1 and ARABIDOPSIS HISTIDINE KINASE2 and 3 Regulate Vascular Tissue Development in *Arabidopsis* Shoots ^W

Jan Hejátko,^{a,1,2} Hojin Ryu,^{b,1} Gyung-Tae Kim,^{c,d} Romana Dobešová,^a Sunhwa Choi,^b Sang Mi Choi,^b Přemysl Souček,^{e,f} Jakub Horák,^a Blanka Pekárová,^a Klaus Palme,^g Břetislav Brzobohatý,^{e,f} and Ildoo Hwang^{b,2,3}

^aDepartment of Functional Genomics and Proteomics, Institute of Experimental Biology, Faculty of Science, Masaryk University, CZ-61137, Brno, Czech Republic

^bDepartment of Life Sciences and Functional Genomics Center, Pohang University of Science and Technology, Pohang 790-784, Korea

^cDepartment of Molecular Biotechnology, Dong-A University, Busan 604-714, Korea

^dEnvironmental Biotechnology National Core Research Center, Gyeongsang National University, Jinju 660-701, Korea

^eInstitute of Biophysics AS CR, CZ-612 65, Brno, Czech Republic

^fMendel University of Agriculture and Forestry in Brno, CZ-613 00, Brno, Czech Republic

^gInstitut für Biologie II/Botany, Freiburg Institute of Advances Studies, Centre of Biological Signaling Studies, Faculty of Biology, University of Freiburg, D-79104 Freiburg, Germany

The development and activity of the procambium and cambium, which ensure vascular tissue formation, is critical for overall plant architecture and growth. However, little is known about the molecular factors affecting the activity of vascular meristems and vascular tissue formation. Here, we show that the His kinase CYTOKININ-INDEPENDENT1 (CKI1) and the cytokinin receptors ARABIDOPSIS HISTIDINE KINASE2 (AHK2) and AHK3 are important regulators of vascular tissue development in *Arabidopsis thaliana* shoots. Genetic modifications of CKI1 activity in *Arabidopsis* cause dysfunction of the two-component signaling pathway and defects in procambial cell maintenance. *CKI1* overexpression in protoplasts leads to cytokinin-independent activation of the two-component phosphorelay, and intracellular domains are responsible for the cytokinin-independent activity of CKI1. *CKI1* expression is observed in vascular tissues of inflorescence stems, and CKI1 forms homodimers both in vitro and in planta. Loss-of-function *ahk2* and *ahk3* mutants and plants with reduced levels of endogenous cytokinins show defects in procambium proliferation and an absence of secondary growth. CKI1 overexpression partially rescues *ahk2 ahk3* phenotypes in vascular tissue, while the negative mutation CKI1^{H405Q} further accentuates mutant phenotypes. These results indicate that the cytokinin-independent activity of CKI1 and cytokinin-induced AHK2 and AHK3 are important for vascular bundle formation in *Arabidopsis*.

INTRODUCTION

Vascular tissue formation in plants is a process with broad developmental and physiological consequences. Factors regulating the proper formation of vascular tissue affect important developmental processes in plants, including the establishment of apical/basal symmetry during embryogenesis (Friml et al., 2003), organogenesis (Scheres et al., 1995; Mähönen et al., 2000), adaxial/abaxial cell fate determination (Emery et al., 2003; Prigge et al., 2005), and cell elongation and differentiation (Szekeres et al., 1996; Cano-Delgado et al., 2004).

Development of vascular tissue entails the differentiation of primary phloem and xylem from procambium, which contains vascular stem cells. Secondary vascular growth is characterized by vascular cambium originating from procambium and interfascicular cambium differentiating from phloem parenchyma and starch sheath cells (Altamura et al., 2001). The mitotic activity and differentiation of vascular and interfascicular cambial cells leads to the formation of secondary xylem and secondary phloem (Altamura et al., 2001; Ye et al., 2002).

Phytohormones appear to be regulatory factors of both primary and secondary vascular growth. Polar auxin transport is presumed to be necessary for the continuity of procambium (Jacobs, 1952; Sachs, 2000), and gibberellins are positive regulators of biomass production in hybrid aspen (*Populus tremula* × *Populus tremuloides*; Eriksson et al., 2000). Cytokinins have been suggested to be important regulators of primary vascular growth (Aloni, 1987; Medford et al., 1989), but their role in the regulation of procambium is just starting to emerge. Two-component signaling, wherein a His kinase receptor transfers a phosphate to downstream response regulators, is key for the

¹These authors contributed equally to this work.

²These authors contributed equally to this work.

³Address correspondence to ihwang@postech.ac.kr.

The authors responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantcell.org) are: Jan Hejátko (hejatk@sci.muni.cz) and Ildoo Hwang (ihwang@postech.ac.kr).

^WOnline version contains Web-only data.

www.plantcell.org/cgi/doi/10.1105/tpc.109.066696

cytokinin response (Hwang and Sheen, 2001; Kim et al., 2006). Cytokinin-induced signaling via its receptor ARABIDOPSIS HISTIDINE KINASE4 (AHK4) and the type-B response regulators ARR1, ARR10, and ARR12 is necessary for procambium formation in *Arabidopsis thaliana* roots (Scheres et al., 1995; Mähönen et al., 2000; Yokoyama et al., 2007). Reduction of endogenous cytokinins by ectopic overexpression of *CYTOKININ OXIDASE/DEHYDROGENASE1* (*CKX1*) or *CKX2* results in the exclusive formation of protoxylem in root vascular bundles (VBs) (Mähönen et al., 2000, 2006a). The role of cytokinin in vascular tissue formation is further suggested by the vascular tissue-specific expression of genes involved in cytokinin biosynthesis (Miyawaki et al., 2004; Zhao et al., 2005) and transport (Hirose et al., 2005, 2008). Factors involved in cytokinin signaling in poplar (*Populus* spp; Nieminen et al., 2008) and cytokinin biosynthesis in *Arabidopsis* (Matsumoto-Kitano et al., 2008) were shown to be principal regulators of the cambium activity and positive regulators of the radial growth via secondary thickening. Nonetheless, the nature of cytokinin action in the primary vascular meristems of shoots, which supply the majority of economically useful plant biomass, is still largely unknown. In addition to hormonal regulations, recent studies of dodeca-peptides, *CLV3/ESR-related41* (*CLE41*) and *CLE44*, and their receptor, *PHLOEM INTERCALATED WITH XYLEM*, showed that non-cell-autonomous communication between phloem and procambium is essential for procambium proliferation and polarity as well as xylem differentiation in the VB development (Fisher and Turner, 2007; Hidakawa et al., 2008). However, although few molecular factors regulating individual processes during vascular tissue formation and differentiation have been identified (Fukuda, 2004; Carlsbecker and Helariutta, 2005; Baucher et al., 2007), our knowledge of the molecular regulators of procambium and vascular cambium is still fragmentary.

Here, we report that the His kinase CYTOKININ-INDEPENDENT1 (CKI1) is important for vascular development via the regulation of procambium proliferation and/or the maintenance of its identity. Genetic manipulation of CKI1 activity leads to abnormal two-component signaling and defects in vascular tissue formation in *Arabidopsis* shoots. Cytokinin depletion and mutations in the cytokinin receptors *AHK2* and *AHK3* result in defects in vascular tissue formation in the inflorescence stem. Collectively, these results suggest that the two-component phosphorelay system is a key regulatory pathway for VB development in *Arabidopsis* shoots.

RESULTS

***CKI1* Is Expressed in Specific Cell Types of VBs in *Arabidopsis* Inflorescences**

To investigate the physiological function of the putative sensor His kinase CKI1 in *Arabidopsis* sporophyte development, we first determined the transcriptional activity of *CKI1* in *ProCKI1:uidA* and *ProCKI1:R12-uidA* transgenic lines that carry the *uidA* marker gene under the control of the *CKI1* promoter (Hejátko et al., 2003; Figure 1A; see Supplemental Figure 1 online). In *ProCKI1:uidA* and *ProCKI1:R12-uidA* plants (see Methods),

β -glucuronidase (GUS) activity was mostly detected in the vascular tissue of all floral organs, the top of the inflorescence stem and flower pedicels, and in the branching points adjacent to axillary meristems (Figures 1Aa to 1Ac and 1Ae). Weak but distinct GUS activity was also detectable in male sporogenous tissue (Figure 1Ad). In transverse sections of inflorescence stems, GUS activity was limited to specific cell types of VBs (Figure 1Ba).

To confirm the relevance of the GUS data with *CKI1* expression, the localization of *CKI1* mRNA and CKI1 protein was determined in situ on cross sections of inflorescence stems (Figure 1B). Similar to what was seen for GUS activity, *CKI1* mRNA was detected in differentiating xylem cells and in VB sheath cells (Figure 1Bb). Antibodies raised against the extracellular domain of CKI1 (α CKI1_{ED}) identified the protein in the procambium of VBs, with the most intense signals in the VB sheath cells located at the lateral procambium borders (Figures 1Bc and 1Bd). Weak CKI1 signals were also distinguishable in the xylem (Figure 1B; for the specificity of α CKI1_{ED}, see Supplemental Figure 2 online). The absence of *CKI1* promoter activity and *CKI1* mRNA in the procambium suggests the presence of a signal for procambial CKI1 localization. This was confirmed by immunolocalization of CKI1 in *CKI1*-overexpressing (*Pro35S:CKI1*) lines. Similar to wild-type plants, in *Pro35S:CKI1* lines, the CKI1 protein localized predominantly to procambial cells (Figures 1Bg and 1Bh). Collectively, these data suggest that CKI1 may be involved in growth and development of VB, particularly in procambium development. Weak expression of *CKI1* in the cortex (Figure 1B) might account for an additional role of CKI1 in other aspects of inflorescence stem growth.

***CKI1* Is Involved in Controlling Meristematic Activity and Vascular Tissue Formation**

To further assess the role of CKI1 in the sporophyte development of *Arabidopsis*, we employed a gain-of-function approach, as mutants completely lacking *CKI1* cannot be obtained due to the infertility of female gametes carrying *cki1* insertion alleles (Pischke et al., 2002; Hejátko et al., 2003). Ectopic overexpression of *CKI1* caused pleiotropic developmental changes (Figures 2A and 2B). *Pro35S:CKI1* transgenic lines were found to be partially or almost completely sterile and to have dramatically shorter siliques. Immunoblot analysis showed that sterility correlated well with *CKI1* expression levels (Figures 2A and 2B). *Pro35S:CKI1* lines also had unusually thick fasciated inflorescence stems, along with changes in overall VB architecture (Figure 2C). Ectopic formation of increased numbers of VBs was observed in transverse sections of inflorescence stems, which suggests higher mitotic activity and abnormal differentiation (Figure 2E). The overall number of cells in transgenic stems was dramatically increased compared with wild-type stems, as seen in transverse sections of inflorescence stems (Figures 2Ea and 2Ed). *CKI1*-overexpressing plants also developed additional inflorescence branches that were initiated from axillary meristems (Figures 2D and 2F). Longitudinal sections of axillary buds revealed additional meristematic tissues bearing many smaller cells (Figure 2Fb, arrow). These findings further suggest that CKI1 might be involved in the regulation of cell division in vascular and meristematic tissues and in their development.




Figure 1. Expression of *CKI1* in VBs.

(A) GUS activity in flowering transgenic plants harboring *ProCKI1:R12-uidA* [(a) and (c) to (e)] or *ProCKI1:uidA* (b). (a) Top of the *Arabidopsis* inflorescence. Note the intensity of the signal in the subapical region of the inflorescence stem, vascular tissues of floral organs, and floral pedicels (arrowheads). (b) and (c) Floral organs before (b) and at/just after anthesis (c). Note the predominant GUS staining in the pistil in the flowers before anthesis [(b); arrowhead]; conversely, the signal in the vascular tissue of stamens is stronger in flowers at/just after anthesis [(c); arrow]. (d) Male sporophytic tissue (arrowhead). (e) Axillary meristem. Bars = 500 μm in (a), (c), and (e) and 100 μm in (b) and (d).

(B) *CKI1* expression in VBs of the inflorescence stem of a *ProCKI1:R12-uidA* plant. GUS activity is seen in cells of the VB sheath located at the lateral (outer) borders of the VB (arrowhead) and xylem (arrows; see also [b]). (b) In situ

To examine *CKI1* action in vascular development and to avoid possible artifacts due to *CKI1* overexpression, we employed RNA interference (RNAi) to knock down the level of *CKI1*. The relative amounts of *CKI1* transcripts and proteins in RNAi transgenic plants were determined by immunostaining and quantitative real-time PCR (Figure 1B; see Supplemental Figure 3A online). Wild-type and transgenic plants were grown under long-day conditions to the stage at which the first silique is formed on the inflorescence. We found that in comparison to wild-type plants, the procambial cell file layers of RNAi lines were decreased (Figures 3Ac, 3Ad, and 3Ag). By contrast, the number of procambial cells in VBs of inflorescence stems in *Pro35S:CKI1* plants (122.8 ± 25 , $n = 6$; mean \pm SE) was increased compared with wild-type plants (77.6 ± 6.1 , $n = 8$) (Figures 3Ae and 3Af; see Supplemental Figure 4B online; for an example of quantification of procambial cells, see Supplemental Figure 4A online). *CKI1* expression in analyzed RNAi lines was not completely absent, as shown by both RNA and protein levels (Figure 1B; see Supplemental Figure 3A online), suggesting that even a partial reduction of *CKI1* expression might lead to phenotypic changes. We therefore analyzed two independent T-DNA insertion mutants in *CKI1*, *cki1-5/CKI1* and *cki1-6/CKI1* (Pischke et al., 2002). *CKI1* transcripts in heterozygous plants of both lines were reduced up to 50% of the wild-type level (see Supplemental Figure 3B online). Defects similar to, but not identical, those identified in the *CKI1* RNAi plants (i.e., reduction of procambium and abnormal cell shape) were observed in heterozygous *cki1-6* plants (Figure 3B; see Supplemental Figures 4A and 4B online). The number of procambial cells in VBs of inflorescence stems in *cki1* heterozygotes (51.1 ± 5 , $n = 14$) was significantly lower than in wild-type plants (84.9 ± 5.7 , $n = 12$). These results suggest that quantitative changes in the *CKI1* activity result in a mutant phenotype and, furthermore, indicate that *CKI1* is important for the maintenance of mitotic activity and/or the identity of procambial cells during VB development in *Arabidopsis*.

CKI1 Acts through the Two-Component Signaling Pathway

CKI1 shares similarity with members of the His kinase family, and *CKI1* His kinase activity has been reported in heterologous and *Arabidopsis* protoplast systems (Hwang and Sheen, 2001; Yamada et al., 2001; Mähönen et al., 2006a). To understand the mechanism by which *CKI1* affects vascular tissue development, we inspected the His kinase activity of *CKI1* in a two-component signaling network by measuring both the activity of the cytokinin-responsive *ARR6* promoter fused to a luciferase (*LUC*) reporter gene and measuring cytokinin-dependent *ARR2*

localization of *CKI1* mRNA. (c) to (h) In situ immunolocalization of *CKI1* using $\alpha\text{CKI1}_{\text{ED}}$ polyclonal antibodies in the cambium of VBs (deep-purple signal, arrows) on cross sections of inflorescence stems of wild-type [(c) and (d)], *CKI1* RNAi [(e) and (f)], and *Pro35S:CKI1* plants [(g) and (h)]. px, protoxylem; mx, metaxylem; arrowheads point to the strongest signal, located in cells on the outer border of the VB (cf. with [a] and [b]; arrowheads). Note the procambial localization of *CKI1* even in the *Pro35S:CKI1* line. Bars = 100 μm in (b), (c), (e), and (g) and 50 μm in (a), (d), (f), and (h).




Figure 2. Phenotype Analysis of *CKI1*-Overexpressing Plants.

(A) Expression analysis of *Pro35S:CKI1-HA* transgenic lines. Total protein and RNA from 2-week-old wild-type and transgenic plants were subjected to an immunoblot assay (top) and RT-PCR assay (bottom). RbcL and *actin* serve as input controls in the two assays.

(B) and **(C)** Ectopic expression of *CKI1* leads to sterility, many trichomes **(B)**, and thick fasciated inflorescence stems **(C)**.

(D) Ectopic expression of *CKI1* leads to additional vegetative tissues initiated from lateral meristems.

(E) The architecture of VBs in *Pro35S:CKI1* transgenic plants. Transverse sections **(a)**, **(b)**, **(d)**, and **(e)** and longitudinal sections **(c)** and **(f)** of the inflorescence stems of wild-type (top) and *Pro35S:CKI1* transgenic plants (bottom). The arrows indicate ectopically formed VBs.

(F) The node structures of wild-type and *Pro35S:CKI1* transgenic plants. Longitudinal sections of wild-type **(a)** and *Pro35S:CKI1* transgenic nodes **(b)**. The arrow indicates an ectopic axillary bud in a *Pro35S:CKI1* transgenic plant.

Bars = 100 μ m.

phosphorylation; both of these approaches have proved to be reliable indicators of two-component signaling outputs (Hwang and Sheen, 2001; Kim et al., 2006). *CKI1* induced *ARR6-LUC* activity in both the presence and absence of cytokinin, as previously shown (Hwang and Sheen, 2001; see Supplemental Figure 5A online). However, *CKI1* did not affect expression of the abscisic acid-responsive *RD29A* or auxin-responsive *GH3* promoters, suggesting a specificity of *CKI1*-mediated responses to the two-component phosphorelay (see Supplemental Figure 5A online). Then we tested whether *CKI1* could initiate a phosphorelay to *ARR2*, a type-B response regulator that is involved in cytokinin-mediated two-component responses (Hwang and Sheen, 2001). As previously demonstrated (Kim et al., 2006), the *ARR2* protein was phosphorylated in a cytokinin-dependent

manner, resulting in a gel band shift. By contrast, cytokinin-dependent phosphorylation of *ARR2* was abolished in protoplasts prepared from the loss-of-function *ahk2 ahk3* mutants (Figure 4A). When *CKI1* was expressed in *ahk2 ahk3* cells, *ARR2* phosphorylation was restored regardless of cytokinin treatment. However, overexpression of *CKI1^{H405Q}* carrying a mutation in the conserved functional His residue could not induce *ARR2* phosphorylation in the double mutant (Figure 4A). These results suggest that *CKI1* has cytokinin-independent His kinase activity in the two-component phosphorelay system.

The His residue at position 405 of *CKI1* amino acid sequence is reported to be a primary target of His kinase activity in the two-component phosphorelay (Hwang and Sheen, 2001). We previously showed that the *CKI1^{H405Q}* mutation diminishes the




Figure 3. CK11 Is Involved in the VB Development of Inflorescence Stems.

(A) Suppression of CK11 activity in *CK11* RNAi lines [(c), (d)], and [(g)] results in reduced and disorganized files of cambial cells. Conversely, the overexpression of *CK11* (f) results in an increase in the number of cambium layers. Note the presence of interfascicular cambium in toluidine blue staining of native tissue [(a) to (d)], suggesting the onset of secondary growth and, thus, cambium formation. Fixed material was subjected to phenotypic analysis before the onset of secondary growth, providing evidence for the role of CK11 in procambium development [(e) to (h)]. Overexpression of the negative allele *CK11^{H405Q}* leads to a dramatic reduction in procambium formation. Native staining of handmade sections [(a) to (d)] with toluidine blue and thin sections made from fixed and embedded material [(e) to (h)]. With native toluidine blue staining, the phloem appears as blue, the undifferentiated cambial zone as pink, metaxylem as blue-green, and protoxylem as purple. c, cambium; ic, interfascicular cambium; pc, procambium; mx, metaxylem; p, phloem. Bars = 100 μ m in (a), (c), and (e) to (h) and 50 μ m in (b) and (d).

(B) The phenotypes conferred by reducing *CK11* expression by T-DNA insertion resemble those of *CK11RNAi* plants. Transverse sections of the inflorescence stems of wild-type plants (Ws-2; [a]) and the heterozygous *CK11* T-DNA insertion lines *cki1-5/CK11* (b) and *cki1-6/CK11* (c). Bars = 100 μ m.

cytokinin-dependent activation of the *ARR6* promoter in wild-type protoplasts (Hwang and Sheen, 2001), suggesting that this mutation might act in a dominant-negative manner in AHK2-, AHK3-, and AHK4-mediated cytokinin signaling pathway.

To determine the mechanism for this negative regulation, wild-type protoplasts were transfected with *CK11^{H405Q}* and *ARR6-LUC* along with the His kinases *AHK2*, *AHK3*, or *AHK4* and treated with cytokinin. Interestingly, *CK11^{H405Q}* suppressed the AHK2-, AHK3-, and AHK4-mediated *ARR6-LUC* activation that was induced by exogenous cytokinins (Figure 4B; see Supplemental Figure 5B online). Accordingly, when wild-type protoplasts were transfected with *AHK2^{H597Q}*, *AHK3^{H460Q}*, or *AHK4^{H459Q}* carrying mutation in the conserved His residue along with wild-type *CK11* and the *ARR6-LUC* reporter gene, the CK11-mediated activation of the *ARR6* promoter was also blocked (Figure 4B; see Supplemental Figure 5B online). These data indicate that CK11 is connected to the two-component signal transduction pathway via its His kinase activity and that the negative effect of the *CK11^{H405Q}* protein is exerted via its inter-

ference with signaling mediated by the other His kinases AHK2, AHK3, or AHK4. Moreover, *Pro35S:CK11^{H405Q}* transgenic lines displayed defects in VBs (Figure 3Ah). Notably, these lines exhibited abnormal cell morphology with irregularly sized cells in both the xylem and phloem. These results provide additional experimental evidence for the functional importance of two-component mediated signaling in proper VB formation in *Arabidopsis*.

Dimerization of His kinases in plants and bacteria was previously demonstrated (Schaller et al., 1995; Surette et al., 1996; Tomomori et al., 1999; Gao et al., 2008; Grefen et al., 2008). Thus, we tested whether CK11 directly interacts with other His kinases in *Arabidopsis* two-component signaling. To do this, myc-tagged *CK11* was cotransfected with HA-tagged *CK11*, *AHK3*, or *AHK4* in *Arabidopsis* protoplasts. When whole protoplast lysates were immunoprecipitated with an anti-myc antibody, CK11-HA but not AHK3-HA or AHK4-HA was pulled down together with CK11-myc, either in the presence or absence of exogenous cytokinins (Figure 4C). Wild-type CK11 protein still interacted with the




Figure 4. CKI1-Mediated Signaling Is Connected to the Two-Component Signal Transduction Pathway.

(A) CKI1 induces cytokinin-independent ARR2 phosphorylation. Protoplasts from *ahk2 ahk3* plants were cotransfected with *ARR2-HA* along with *CKI1-HA* or *CKI1^{H405Q}-HA*, incubated for 6 h, and treated with 100 nM t-zeatin (cytokinin) in the presence of 100 μM cycloheximide for 1 h. Wild-type protoplasts transfected with *ARR2* served as a control. The mobility shift of ARR2 induced by phosphorylation was detected with an anti-HA antibody. Equal amounts of protein were loaded on each lane.

(B) A negative form of CKI1 protein represses the AHK4-mediated induction of *ARR6*. Protoplasts from wild-type plants were transfected with *ARR6-LUC* alone, wild-type *AHK4*, wild-type *AHK4* plus mutant *CKI1*, wild-type *CKI1* plus mutant *AHK4*. Error bars indicate SE (*n* = 2). *CKI1^{H405Q}* and *AHK4^{H459Q}* are negative versions of *CKI1* and *AHK4*, respectively. Rubisco large subunit (RbcL) stained by Coomassie blue was used as a protein loading control.

CKI1^{H405Q} mutant protein (Figure 4C), suggesting that the His kinase and phosphoryl transfer activities of CKI1 are not required for its dimerization. The self-interaction of CKI1 in planta was confirmed using a bimolecular fluorescence complementation system (Walter et al., 2004). Coexpressed CKI1-cYFP and CKI1-nYFP, but not CKI1-cYFP and AHK1-nYFP, produced strong yellow fluorescent protein (YFP) fluorescence at the plasma membrane in tobacco (*Nicotiana tabacum*) leaf cells (Figure 4D). To determine whether CKI1 forms a dimer or a higher-order multimer, detergent-solubilized proteins from protoplasts expressing *CKI1-HA* were treated with the cross-linker bis-sulfosuccinimidyl suberate (BS³) (Figure 4E). The intensity of immunoreactive bands corresponding to the approximate size of the CKI1 monomer was gradually reduced as the BS³ concentration increased, while the intensity of a higher band with the approximate predicted size of a CKI1 dimer was concomitantly increased. Taken together, these results indicate that CKI1 forms homodimers both in vitro and in planta. However, in contrast with sensor His kinases involved in ethylene signaling, which form heterodimers (Gao et al., 2008; Grefen et al., 2008), CKI1 does not form heterodimers with any of the tested His kinases.

To confirm the His kinase activity of CKI1 in planta, we examined the expression of type-A *ARR* genes, the cytokinin primary response genes (D'Agostino et al., 2000), in *Pro35S:CKI1*, *Pro35S:CKI1^{H405Q}*, and *CKI1* knockdown lines. The ectopic expression of *CKI1* induced the expression of a subset of type-A response regulators, including *ARR3*, 4, 7, 8, 9, 15, 16, and 17 (Figure 4F). By contrast, overexpression of *CKI1^{H405Q}* significantly reduced the expression of *ARR6*, 7, and 15 (Figure 4F), thus confirming the negative regulatory role of CKI1^{H405Q} in the two-component phosphorelay. Moreover, in heterozygous *cki1-5* and *cki1-6* lines, the expression of most of the inspected *ARR* genes was upregulated (Figure 4G), further demonstrating that CKI1 exerts its action through the two-component signal transduction pathway in planta. Furthermore, these results imply that changing CKI1 activity via site-directed mutagenesis and/or deregulation of endogenous *CKI1* expression leads to differential changes in expression of individual *ARRs*, suggesting a disturbance of the two-component phosphorelay.

The Cytoplasmic CKI1 Domain Is Necessary for Its His Kinase Activity

Our data suggested that CKI1 can activate the two-component phosphorelay via its His kinase activity, which is independent of exogenously added cytokinins. To unravel the potential importance of extracellular and intracellular CKI1 domains in CKI1-mediated signaling, we constructed chimeric receptors composed of CKI1 and AHK4 (CKI1-AHK4 and AHK4-CKI1) along with truncated forms of CKI1 (see Supplemental Figure 5C online). AHK4-CKI1, which consists of the extracellular and transmembrane domains of AHK4 fused to the kinase and receiver domains of CKI1, could activate the *ARR6* promoter as efficiently as wild-type CKI1, either in the presence or absence of cytokinins (see Supplemental Figure 5D online). However, CKI1-AHK4, which consists of the extracellular and transmembrane domains of CKI1 fused to the intracellular domain of AHK4, could not enhance the activity of the *ARR6* promoter, regardless of the presence or absence of cytokinins. Moreover, CKI1 Δ N, which lacks the extracellular domain of CKI1, still constitutively activated *ARR6-LUC*, unlike CKI1 Δ C, which consists of the extracellular and transmembrane domains of CKI1 (see Supplemental Figure 5D online). *Pro35S:AHK4-CKI1* transgenic lines displayed similar *CKI1*-overexpressing phenotypes with reduced fertility, shorter siliques, and additional inflorescence branches (see Supplemental Figures 6A and 6B online). They also had thick fasciated inflorescence stems with increased mitotic activity (see Supplemental Figure 6C online). Thus, the cytoplasmic kinase domain of CKI1 is sufficient for CKI1 cytokinin-independent His kinase activity in two-component signaling.

Cytokinins Regulate VB Development of *Arabidopsis* Inflorescence Stems via the AHK2 and AHK3 Signaling Pathway

Our data suggest that CKI1 regulates the development of vascular tissue in shoots via its His kinase activity. Proteins involved in the cytokinin-regulated two-component signaling pathway are known to regulate vascular tissue formation in *Arabidopsis* roots (Mähönen et al., 2000, 2006b; Hutchison et al., 2006). In addition,

Figure 4. (continued).

(C) CKI1-HA but none of the tested AHKs-HA proteins co-immunoprecipitate with myc-tagged CKI1. Mesophyll protoplasts from wild-type plants were transfected with *CKI1-HA*, *CKI1^{H405Q}-HA*, *AHK3-HA*, or *AHK4-HA*, with or without *CKI1-myc*, incubated for 6 h, and then immunoprecipitated with anti-myc antibodies. CKI1 proteins were detected with an anti-HA antibody.

(D) CKI1 forms homodimers in tobacco leaf cells. Confocal images of abaxial epidermal tobacco leaf cells expressing the indicated YFP-N and YFP-C fusion proteins demonstrate YFP fluorophore reconstitution due to protein-protein interaction of the tested proteins (top row). The bottom row shows the corresponding bright-field images of the transiently transformed cells. Bars = 50 μ m.

(E) CKI1 forms dimers. Protoplasts expressing *CKI1-HA* were solubilized with Triton X-100. Total protein was treated with increasing amounts of the cross-linker BS³ and subjected to SDS-PAGE. Two bands corresponding to the predicted sizes of the CKI1 monomer and dimer were detected with the anti-HA antibody.

(F) and **(G)** Genetic manipulation of CKI1 activity affects two-component signaling in planta. Transgenic plants overexpressing CKI1 or CKI1^{H405Q} **(F)** or CKI1 T-DNA insertion lines **(G)** show changes in the expression of specific type-A *ARRs*. Quantitative RT-PCR was performed with total RNA extracted from 3-week-old seedlings **(F)** or inflorescence stems **(G)** using gene-specific primers for type-A *ARRs* (see Supplemental Table 1 online for primer sequences). Error bars indicate SE ($n = 8$ **[F]** and 3 **[G]**). Asterisks indicate statistically significant differences from wild-type transgenic plants analyzed by Student's *t* test (* $P < 0.05$; ** $P < 0.01$).

the role of cytokinins in the cambium growth activity was recently identified (Matsumoto-Kitano et al., 2008; Nieminen et al., 2008). These results raised the possibility that two-component signaling cascades initiated by cytokinins as well as by CKI1 are also involved in the regulation of VB formation in shoots. Thus, we examined vascular tissue morphology in the inflorescence stems of plants with mutations in individual cytokinin receptors and in double mutants. In the *ahk2* mutant, the number of cell layers in the procambial region was decreased (Figures 5Ab, 5Ag, and 5Ai). A weaker phenotype was identifiable in the *ahk3* line (Figures 5Ac, 5Ah, and 5Am). In the *ahk2 ahk3* double mutant, reduction of the procambium and in the size of VBs was more pronounced than in either single mutant (Figures 5Ad, 5Ai, and 5An). A similar phenotype was also observed as a result of endogenous cytokinin depletion in *Pro35S:CKX3* and *Pro35S:CKX1* lines (Figures 5Ae, 5Aj, and 5Ao, respectively). In *ahk2 ahk3* plants and in lines with decreased endogenous cytokinin, we further observed that interfascicular cambium failed to form when compared with the wild type in Figure 5Aa, suggesting defects in the onset of secondary growth. Taken together, AHK2 and AHK3 together with CKI1 play important roles in proper VB development, especially in the maintenance of procambial cell identity and/or regulation of procambial cell proliferation.

To confirm that CKI1 can affect vascular tissue development via the AHK2/3 signaling pathway, we ectopically expressed *CKI1* or *CKI1^{H405Q}* in *ahk2 ahk3* mutants. Ectopic expression of *CKI1* partially rescued the growth defects of these mutants (Figures 5B and 5C). The rosette leaves and petioles of *ahk2 ahk3/Pro35S:CKI1* transgenic plants were similar to those of wild-type plants (Figure 5B). Overexpression of *CKI1* in the *ahk2 ahk3* background resulted in an increase of cambial layers, with a two- to threefold increase in the diameter of inflorescence stems compared with *ahk2 ahk3* (Figure 5D). In addition, a reduced number of cells with the irregular size of *ahk2 ahk3* in the xylem, phloem, and cambial layers were partially restored in these transgenic lines (Figure 5E). As a result, the radial growth was rescued, manifested by almost, but still partially, wild-type-like diameter of the inflorescence stem in *Pro35S:CKI1/ahk2 ahk3* (Figure 5E). By contrast, ectopic expression of the dominant-negative mutation *CKI1^{H405Q}* further accentuated the mutant phenotypes of *ahk2 ahk3* plants. In comparison to the *ahk2 ahk3* mother line, the aerial parts and diameters of inflorescence stems of *ahk2 ahk3/Pro35S:CKI1^{H405Q}* plants were much smaller (Figures 5B and 5D). The cambial cell layers were unidentifiable, and vascular tissue differentiation was nearly abolished (Figure 5E). Collectively, these results suggest that CKI1 is functionally conserved with AHK2 and AHK3 in VB development but that it still has its own specificity in the regulation of vascular tissue development.

DISCUSSION

Cytokinin-Independent CKI1 Regulates Two-Component Phosphorelay in *Arabidopsis*

CKI1 was the first His kinase implicated in the perception of cytokinins (Kakimoto, 1996). However, CKI1 does not contain the

cytokinin binding CHASE domain and does not bind cytokinins in vitro (Yamada et al., 2001). Defects in megagametogenesis conferred by a *cki1* loss-of-function allele, together with observations of *CKI1* expression in the ovule and endosperm, show that CKI1 is critical in female gametophyte development (Pischke et al., 2002; Hejátko et al., 2003). However, the mechanisms underlying the involvement and action of CKI1 signaling in specific biological processes during *Arabidopsis* gametophyte and/or sporophyte development remain largely unknown. Furthermore, when overexpressed in plants, calli, or protoplasts, *CKI1* induces typical cytokinin responses, including shoot regeneration, delay of leaf senescence, and activation of the cytokinin-responsive *ARR6* promoter in the absence of exogenously applied cytokinins (Kakimoto, 1996; Hwang and Sheen, 2001). It was suggested that ectopic expression of *CKI1* allows the expressing cells to sense low concentrations of endogenous cytokinins that are otherwise unable to trigger shoot formation (Kakimoto, 1996). Here, we have shown that CKI1 can mediate cytokinin-independent regulation of the two-component signaling pathway. Thus, rather than recognition of endogenous cytokinin levels as suggested previously (Hwang and Sheen, 2001), the cytokinin-independent His kinase activity of CKI1 probably leads to the cytokinin-like phenotype in calli overexpressing *CKI1* (Kakimoto, 1996). However, the possibility that the extracellular domain of CKI1 allows another mode of cytokinin-independent regulation of its His kinase activity cannot be excluded.

We found here that CKI1 shares at least some of the signaling proteins with the two-component phosphorelay system in the cytokinin signaling pathway. Based on our results, and of studies showing dephosphorylation of AHP1 and AHP2 by CKI1 in vitro (Nakamura et al., 1999), functional complementation of bacterial and yeast His kinase mutants by CKI1 (Yamada et al., 2001), and CKI1 interaction with AHP proteins in a yeast two-hybrid system (Dortay et al., 2006), we conclude that CKI1 can activate the two-component phosphorelay in *Arabidopsis* via proteins involved in the cytokinin signaling pathway. Whether CKI1 activity directly affects cytokinin signaling and/or other adaptive responses mediated by the two-component phosphorelay (e.g., osmosensing or abscisic acid responses), however, remains to be determined. *CKI1* overexpression activated *ProARR6:LUC*, a marker for two-component signaling in protoplasts, but we could not observe similar activation of *ARR6* in a late developmental stage of *CKI1*-overexpressing plants. This result implies that a single cell system may not always reflect different developmental stages at which multiple cells incorporate diverse external and/or internal signals to properly execute growth and development programs (Figure 4; see Supplemental Figure 5 online).

The output of the two-component phosphorelay was proposed to be a result of interactions of multiple His kinases and their kinase and phosphatase activities (Mähönen et al., 2006a). In this model, the final output of the two-component phosphorelay also depends on the expression levels of cytokinin binding and cytokinin nonbinding His kinases, including CKI1. The phosphatase activity of the receiver domain of CKI1 has been demonstrated (Nakamura et al., 1999), suggesting that CKI1 might contribute to two-component phosphorelay regulation via both kinase and phosphatase activities. Here, we have shown that both overexpression and downregulation of CKI1 affects the




Figure 5. Cytokinin Regulates VB Formation via AHK2 and AHK3 Phosphorelay.

(A) Transverse sections of the inflorescence stems of wild-type (Columbia-0 [Col-0]) (**[a]**, **[f]**, and **[k]**), *ahk2* (**[b]**, **[g]**, and **[l]**), *ahk3* (**[c]**, **[h]**, and **[m]**), *ahk2 ahk3* (**[d]**, **[i]**, and **[n]**), and *Pro35S:CKX* (**[e]**, **[j]**, and **[o]**) lines. Note the reduction of procambial layers in *ahk2* and *ahk3* plants and particularly in *ahk2 ahk3* double mutants. The overall reduction in VB size is apparent in *ahk2 ahk3* and in *Pro35S:CKX* lines, suggesting positive regulation of cytokinin signaling via the *AHK2/AHK3* pathway in VB development in *Arabidopsis* inflorescence stems. Staining of handmade sections (**[a]** to **[j]**) with toluidine blue and thin sections made from fixed and embedded material (**[k]** to **[o]**). c, cambium; ic, interfascicular cambium; pc, procambium; mx, metaxylem; p, phloem. Bars = 100 μ m in (**a**) to (**e**) and (**k**) to (**o**) and 50 μ m in (**f**) to (**j**).

(B) The dwarfism resulting from deletion of *ahk2 ahk3* is rescued in the presence of CK11. Three-week-old wild-type (Col-0) and transgenic plants expressing *Pro35S:CK11-HA* or *Pro35S:CK11^{H405Q}-HA* in the *ahk2 ahk3* background were used for phenotypic analysis.

(C) Expression analysis of HA-tagged CK11 and CK11^{H405Q} proteins under the control of the *Pro35S* promoter in transgenic lines. Total proteins from 3-week-old plants of each designated line were subjected to 7.5% SDS-PAGE. Actin proteins detected by immunoblotting serve as input controls.

(D) and **(E)** Ectopic expression of *CK11* rescues the abnormal vasculature of the *ahk2 ahk3* mutant. Microscopy images of transverse sections of the inflorescence stems of wild-type (**a**), *ahk2 ahk3* (**b**), *Pro35S:CK11-HA/ahk2 ahk3* (**c**), and *Pro35S:CK11^{H405Q}-HA/ahk2 ahk3* (**d**) plants. The arrow indicates extensively reduced VB. Bars = 100 μ m.

output of the two-component signaling pathway, as measured by regulation of the expression of *ARRs*. This observation accords well with the above-described model proposed by Mähönen et al. (2006a) and suggests that an equilibrium of individual inputs into the two-component pathway is critical for proper vascular tissue development in *Arabidopsis* shoots. Type-A *ARR* genes have been identified as negative regulators of cytokinin signaling (To et al., 2004). Auxin-induced regulation of *ARR7* and *ARR15* was recently identified as a mechanism of auxin-dependent spatial-specific attenuation of cytokinin signaling during stem cell niche formation in *Arabidopsis* roots (Müller and Sheen, 2008). Thus, upregulation of negative type-A *ARRs* in knock-down *CK11* lines might disrupt the proper regulation of two-component signaling in procambial development; therefore, these lines may partially phenocopy plants deficient in cytokinin signaling. However, whether CKI1-regulated expression of *ARR* genes represents another cytokinin-independent mechanism for regulation of the cytokinin two-component pathway remains to be determined. Furthermore, it is still uncertain if cytokinin-responsive *ARR* genes are direct regulators required for cambial development, although it is evident that perturbation of cytokinin homeostasis affects cambial activity in shoots and roots (Matsumoto-Kitano et al., 2008; Nieminen et al., 2008).

CKI1 Together with AHK2/AHK3 Is Involved in the Maintenance of Procambial Activity during VB Development in *Arabidopsis* Shoots

Vascular development in *Arabidopsis* can be divided into three major steps: (1) initiation and maintenance of (pro)cambium, (2) asymmetric cell patterning and differentiation into xylem and phloem precursor cells, and (3) their final specification into distinct xylem and phloem cell types. While auxin initiates and maintains continuous vascular pattern formation of procambial cells via polar auxin transport (Fukuda, 2004; Friml et al., 2004), cytokinin signaling mediated by AHK2/3 is unlikely to be involved in the initiation of procambial cell files as knockout lines still contain functional VBs (Figure 5). Rather, our results suggest that CKI1 and AHK2/3 are required for the proliferation and maintenance of procambial cells and vascular stem cells, which give rise to primary vascular tissues and vascular cambium.

Besides the procambium activity, the activity of the shoot apical meristem (SAM) seems to be genetically linked with the regulation of vascular tissue development (Baucher et al., 2007). Cytokinins were shown to be positive regulators of the shoot meristem size (Higuchi et al., 2004; Nishimura et al., 2004; Kurakawa et al., 2007). Thus, the downregulation of the diameter of the inflorescence stem and the size of VBs in *ahk2 ahk3* mutants might be at least partially due to the defects in the SAM activity during procambium initiation. Accordingly, formation of enlarged and fasciated inflorescence stems in *Pro35S:CK11* lines could be affected by the increased mitotic activity in the SAM upon *CK11* overexpression. However, we could not observe any quantitatively significant change of the SAM size in the transgenic lines overexpressing *CK11* (see Supplemental Figure 8A online). In addition, we have analyzed the VB phenotype at the very base of the first internodium and, thus, in a position spatially and developmentally well dissected from the shoot apical mer-

istem. The analysis was performed at the stage when the first silique was formed on the inflorescence. This stage corresponds to the end of the primary growth, which is primarily governed by the procambial activity (Altamura et al., 2001). Taken together, although we cannot completely exclude the possibility that some of the observed phenotype changes originate in the SAM, the defects in the procambium activity due to impaired cytokinin signaling seem to be at least one of the substantial contributions to the observed defects in the primary radial growth of *ahk2 ahk3* and *Pro35S:CKX1(2)* lines.

Interestingly, AHK2/3-mediated cytokinin signaling seems to be also involved in secondary VB development. Formation of interfascicular cambium is one of the anatomically well distinguishable markers of the secondary growth initiation in *Arabidopsis* (Altamura et al., 2001). We observed that the formation of interfascicular cambium was absent and/or substantially reduced in the *ahk2 ahk3* double mutant and in *Pro35S:CKX1(2)* lines, which suggests possible defects in the onset of the secondary thickening. It is possible that CKI1 maintains the basal meristematic activity of procambial cells and that AHK2/3 fine-tunes (pro)cambial activity following environmental and/or developmental cues that regulate endogenous cytokinin levels (Samuelson et al., 1992; Yang et al., 2001; Takei et al., 2004; Werner et al., 2006; Matsumoto-Kitano et al., 2008). Recently, results showing the involvement of cytokinin in the regulation of cambium in *Arabidopsis* and poplar were published (Matsumoto-Kitano et al., 2008; Nieminen et al., 2008). A reduction of cytokinins in null mutants of the *Arabidopsis* cytokinin biosynthetic genes *ipt1,3,5,7* and in transgenic poplar trees overexpressing *Arabidopsis* *CYTOKININ OXIDASE/DEHYDROGENASE2* resulted in impaired cambial formation, indicating that cytokinins are important regulators of vascular cambium. These results are consistent and complementary with our findings of a role for His kinase-mediated two-component signaling in vascular tissue formation of *Arabidopsis* shoots.

However, it should be emphasized here that in addition to hormonal regulations, other signals (e.g., weight of the produced biomass of the plant body) (Ko et al., 2004) are also integrated in the regulation of the secondary thickening. Thus, this type of signal might contribute to the observed defects in the onset of secondary thickening in *ahk2 ahk3* and cytokinin-deficient plants, both of which are deficient in radial growth, thus revealing lowered production of the shoot biomass.

It has long been known that roots and shoots respond differently to cytokinins in *Arabidopsis* (Werner et al., 2003). In *Arabidopsis* roots, cytokinins have been shown to be necessary for the periclinal procambial cell divisions required for the proliferation of vascular cell files (Scheres et al., 1995; Mähönen et al., 2000, 2006b). Similar to what we have found, this suggests a positive role of cytokinins for procambium proliferation and/or maintenance. However, CKX1(2)-mediated depletion of cytokinins in *Arabidopsis* roots leads to the formation of abnormal vascular tissue that is devoid of phloem but which exhibits abundant protoxylem formation (Mähönen et al., 2006b). This is apparently not the case in the inflorescence stem, where CKX1(3)-mediated cytokinin depletion led to the formation of VBs of reduced size; however, all cell types (i.e., protoxylem, metaxylem, and phloem) still could be detected (Figure 5C). Accordingly,

we did not observe specific phenotypic changes in root and hypocotyl vascular development in *CK11* knockdown lines (see Supplemental Figures 8B and 8C online). This could be explained by a lower sensitivity of the inflorescence stem to cytokinin depletion and by the specificity of CK11 and AHK2/3 signaling. CK11-driven, cytokinin-independent regulation of VB development could contribute to the lower sensitivity and resulting phenotype in *Pro35S:CKX1(3)* inflorescence stems. Alternatively, modified developmental pathways might operate in root and shoot vascular tissue development.

Here, we have shown that His kinases in *Arabidopsis* regulate vascular tissue formation in shoots via the regulation of procambium activity (see Supplemental Figure 9 online). This is of great economic importance as procambium and vascular cambium activities regulate biomass production in plants. Thus, regulation of the activity of individual His kinases by means of genetic engineering might be used to regulate biomass production in plants and might help us to lower our dependence on other, mostly nonrenewable, energy resources.

METHODS

Plant Materials

The *Arabidopsis thaliana* Col-0 ecotype and the mutant carrying the *ahk2-1* and *ahk3-1* mutant alleles (Higuchi et al., 2004), both in the Col-0 background, were used. Wild-type and mutant plants were grown in an environmentally controlled room at 23°C under white light with 14-h-light/10-h-dark cycles.

Transient Expression in *Arabidopsis* Protoplasts

Transient expression in protoplasts was performed as previously described (Hwang and Sheen, 2001). Typically, 2×10^4 protoplasts were transfected with 20 μ g total plasmid DNA consisting of different combinations of the reporter, effectors, and internal control. Transfected protoplasts were incubated at 10^4 cells per mL with or without 100 nM t-zeatin (Sigma-Aldrich) for 6 h. As an internal control, the *GUS* reporter gene fused to the *Arabidopsis* ubiquitin promoter (*UBI10-GUS*) was used. The results shown are the means and error bars of relative LUC activities obtained from duplicate samples. All assays were performed at least three times, and similar results were obtained in all experiments.

Plasmid Constructs and Generation of Transgenic Plants

Full-length and truncated *CK11* were amplified by PCR from genomic *Arabidopsis* DNA. Full-length *AHK4* was obtained by PCR from an *Arabidopsis* cDNA library. Chimeric *AHK4-CK11* and *CK11-AHK4* constructs were generated by overlap extension PCR using the overlapping primers 5'-CATCTCTCTCCTTGTGCTTGAGCTGCACCATACAGTATATA-3' and 5'-CTTCGACTTTTACTATGTGCATCATAAACACACAAC-CATAC-3', respectively. The coding regions of all proteins were tagged with either two copies of the hemagglutinin epitope (HA), the myc epitope, or green fluorescent protein and inserted into a plant expression vector containing the *35S4PPDK* promoter and the *NOS* terminator (Hwang and Sheen, 2001). Transgenic *Arabidopsis* plants expressing *CK11-HA* under the control of the *35S4PPDK* promoter were generated by the floral dip method and BASTA selection as described (Clough and Bent, 1998). *Pro35S:CKX2* and *Pro35S:CKX3* lines were generated as described (Pernisova et al., 2009). The ectopic expression of *CK11* was tested by RT-PCR and immunoblot analysis. Phenotypic analyses of

transgenic lines were performed with homozygous T3 plants. All mutants were generated by site-directed mutagenesis using the QuikChange site-directed mutagenesis kit (Stratagene) according to the manufacturer's instructions. All constructs were confirmed by sequencing. To analyze the specificity of the *CK11* promoter, two different constructs were made and introduced into *Arabidopsis* ecotype Col-0. The first construct, *ProCK11:uidA*, contains a 2.7-kb fragment of upstream genomic DNA that includes the putative translational start site of the short open reading frame (MKRAF) in the 5' untranslated region of the *CK11* mRNA. The primers SII-ckpr (5'-GTAACCGCGGGAGGAGGCACAAAATGACGAA-3') and B-ckpr (5'-GCTGGGATCCTCATATTATCTTCTTCTCGGAGC-3') were used for PCR amplification of the putative promoter region of CK11; this fragment is translationally fused with the *uidA* coding sequence (Hejátko et al., 2003). The second construct, *ProCK11:R12-uidA*, also contains a translational fusion of *uidA* with the same genomic fragment described above; however, the 3' end of this fragment was extended to include the CGT codon that encodes the R12 residue of CK11 (see Supplemental Figure 1 online). Multiple independent transgenic lines were inspected in both cases, and no apparent differences in the resulting distribution of GUS activity were detectable.

Expression Analysis

In situ mRNA and GUS staining were performed as previously described (Hartmann et al., 2000; Hejátko et al., 2003; Brewer et al., 2006). Polyclonal rabbit anti-CK11 antibody was prepared against the peptide from the CK11 extracellular domain (GATRIKHQAEEKAKYQC, α CK1_{ED}; Sigma-Genosys) and used for indirect immunolocalization on Steedman's wax sections as described (Vitha et al., 2000). Two batches of polyclonal sera (anti-CK11_{ED120} and anti-CK11_{ED121}) isolated from two independently immunized rabbits were tested (see Supplemental Figure 2 online). If not otherwise mentioned, anti-CK11_{ED121} was used. The alkaline phosphatase-conjugated secondary antibody was visualized by 5-bromo-4-chloro-3-indol phosphate (BCIP)/*p*-Nitro-Blue tetrazolium chloride (NBT) staining. Antibody specificity was characterized on immunoblots using recombinant proteins expressed in *Escherichia coli* and on immunoblot using plant protein extracts (see Supplemental Figure 2 online). Preimmune serum was used as a negative control in immunolocalizations of CK11 on sections, and no signal was obtained.

RT-PCR and Quantitative Real-Time PCR Analysis

Total RNA was isolated using Trizol reagent (Invitrogen) according to the manufacturer's instructions. For RT-PCR, first-strand cDNA was synthesized from 1 μ g RNA with oligo(dT) primers and ImProm-II reverse transcriptase (Promega). The expression of *CK11* was verified with 30 cycles using a gene-specific primer set, CK11fwd (5'-AACAGCTCAAG-GACACCAAG-3') and CK11rev (5'-GCGTTCCTTCATTTTCAATA-3'), and *actin* gene as a control using ACTfwd (5'-GTACAACATGTTCTCAGGT-3') and ACTrev (5'-GAAGCATTTTCTGTGGACAA-3') primers. For quantitative real-time RT-PCR, first-strand cDNA was prepared with SuperScript II reverse transcriptase (Invitrogen) and the ACT-L and rCK11rt primers. The subsequent quantitative PCR was performed in a Light Cycler 2.0 (Roche) with SYBR Premix ExTaq system (Takara) as a fluorescent dye that monitors DNA content. To amplify gene-specific products, the following primers were used: fACTrt (5'-CAGTGTCTG-GATCGGAGGAT-3'), rACTrt (5'-TGAACAATCGATGGACCTGA-3'), fCK11rt (5'-CTATTGGGAACCCAGAGGACG-3'), rCK11rt (5'-AAGCT-TCTTCCCCTGTCGC-3'), and type-A *ARRs* (see Supplemental Table 1 online). The steady state levels of the transcripts were determined by standard curve quantitation. All quantitative RT-PCR experiments were performed with biologically independent samples at least three times.

Coimmunoprecipitation and Immunoblot Analysis

Protoplasts were transfected with either HA- or myc-tagged *CKI1*, *AHK3*, *AHK4*, *CKI1^{H405Q}*, or *AHK4* and then incubated for 6 h to allow protein expression. Total protein was extracted from the transfected protoplasts in IP buffer (50 mM Tris, pH 7.5, 150 mM NaCl, 5 mM EDTA, 1% Triton X-100, protease inhibitor cocktail [Roche], and 1 mM DTT) and incubated with a monoclonal anti-HA antibody (Roche) or a monoclonal anti-c-myc antibody (Cell Signaling). The protein-antibody complex was precipitated with protein A/G plus-agarose beads (Calbiochem). For cross-linking experiments, protoplasts transfected with *CKI1-HA* were incubated for 6 h and lysed with protein extraction buffer (50 mM sodium phosphate, pH 7.4, 5 mM EDTA, 1% Triton X-100, 1 mM DTT, and protease inhibitor cocktail). Total protein extracts were incubated with different concentrations of the cross-linker BS³ (Pierce) for 1 h at 4°C before being quenched with 25 mM Tris-HCl, pH 7.5, for 30 min. Immunoprecipitated proteins and total proteins were subjected to 7.5 or 10% SDS-PAGE and blotted onto Immobilon-P membranes (Millipore). The blots were probed with a peroxidase-conjugated anti-HA antibody (Roche) or a monoclonal anti-myc antibody. Extracellular domains of *AHK4* (*AHK4_{ED}*, D127-P395) and *CKI1* (*CKI1_{ED}*, E28-Q345) were cloned into *E. coli* expression vector pDEST17 and expressed as a recombinant protein in a translational fusion with His-Tag. One hundred micrograms of the total protein (bacterial lysate) was separated using 15% SDS-PAGE, blotted on polyvinylidene fluoride (PVDF) membrane, and immunodetected using monoclonal anti-polyHistidine (Sigma-Aldrich) or polyclonal anti-*CKI1_{ED}* 1:10,000 in blocking buffer (5% milk in 10 mM Tris-HCl, pH 7.5, 150 mM NaCl, and 0.1% Tween-20). The detection was performed using alkaline phosphatase-conjugated goat anti-rabbit secondary antibodies (Sigma-Aldrich) diluted 1:30,000 in blocking buffer and anti-mouse-AP antibodies (Sigma-Aldrich) diluted 1:20,000 in a blocking buffer with BCIP/NBT substrate for 10 min, within the linear range of signal development. All experiments were performed at least three times.

Analysis of CKI1 Dimerization Using Bimolecular Fluorescence Complementation

Entry clones containing *CKI1* and *AHK1* cDNA were prepared according to the manual for Gateway technology in pDONR207 (Invitrogen), verified by sequencing, and subsequently recombined via the LR reaction into pSPYNE-35S and pSPYCE-35S (Walter et al., 2004). Transient transformation of tobacco (*Nicotiana tabacum*) leaves and immunodetection of fusion proteins were performed as previously described (Horak et al., 2008). Confocal laser scanning microscopy was performed using an Olympus IX81 microscope equipped with a Fluoview 500 confocal unit at a setup recommended by the manufacturer for YFP fluorescence detection.

Histological Analysis

Tissue samples were fixed for 24 h in 3% glutaraldehyde and 4% paraformaldehyde in 0.1 M phosphate buffer, pH 7.2, or in FAA containing 5% acetic acid, 45% ethanol, and 5% formaldehyde. The fixed samples were then rinsed with 0.1 M phosphate buffer, pH 7.2, and dehydrated through a graded ethanol series. The specimens were infiltrated and embedded in Spurr's resin (Ted Pella) or Technovit resin (Kulzer and Co.) for 48 h at 65°C. Sections (0.5 or 4 μm) were made using an MT-X ultramicrotome (RMC), stained in 0.1% toluidine blue, and photographed with a Zeiss Axioplan2 microscope. For native staining, handmade sections were prepared with a razor blade from the base of the inflorescence stems when the first silique appeared. Sections were stained with toluidine blue (0.05% [w/v] solution in water) for 1 min, destained in distilled water for 30 seconds, mounted in 50% glycerol, and observed with a microscope (Olympus BX 61) using differential interference contrast microscopy.

Accession Numbers

Sequence data from this article can be found in the Arabidopsis Genome Initiative or GenBank/EMBL databases under the following accession numbers: *CKI1* (AT2G47430), *AHK1* (AT2G17820), *AHK2* (AT5G35750), *AHK3* (AT1G27320), *AHK4* (AT2G01830), *ARR2* (AT4G16110), *CKX1* (AT2G41510), *CKX2* (AT2G19500), and *CKX3* (AT5G56970). Germplasm identification numbers from this article are as follows: *cki1-5* (CS6360) and *cki1-6* (CS6361).

Supplemental Data

The following materials are available in the online version of this article.

Supplemental Figure 1. Schematic Representation of the *ProCKI1:uidA* (Up) and *ProCKI1:R12-uidA* (Down) Constructs Used in Analysis of the Transcriptional Specificity of the *CKI1* Promoter.

Supplemental Figure 2. Anti-*CKI1_{ED}* Antibody (α *CKI1_{ED}*) Specifically Recognizes the Extracellular Domain of *CKI1*.

Supplemental Figure 3. Transcript Levels of *CKI1* in RNAi Lines and T-DNA Insertion Heterozygous Lines.

Supplemental Figure 4. *CKI1* Regulates the Number of Procambial Cells.

Supplemental Figure 5. *CKI1* Specifically Enhances the Activity of the Two-Component, Cytokinin-Responsive *ARR6* Promoter in a Cytokinin-Independent Manner.

Supplemental Figure 6. The Ectopic Expression of *CKI1* or *AHK4-CKI1* Leads to Sterility and Formation of Short Siliques and to Additional Vegetative Tissues Initiated from Lateral Meristems.

Supplemental Figure 7. Expression of Individual Constructs Used in the BIFC Assay in Figure 5B Was Determined by Immunostaining and Ponceau S Staining to Prove Equal Protein Loading (Red Bands).

Supplemental Figure 8. *CKI1* Activity Does Not Affect Either the SAM Activity or the Vascular Bundle Development in Hypocotyl and Root.

Supplemental Figure 9. A Proposed Model for *CKI1* and Cytokinin Action Mechanism in the Vascular Bundle Development of Inflorescence Stems.

Supplemental Table 1. Gene-Specific Primers for Type-A *ARRs*.

ACKNOWLEDGMENTS

We thank Filip Rolland and Jiří Friml for critically reading the manuscript, Thomas Schumüller for *Pro35S:CKX1* and *Pro35S:CKX2* seeds, and Chiharu Ueguchi and Yka Helariutta for *ahk* mutant seeds. This work was supported by grants to I.H. from the Plant Diversity Research Center of MOST, the Plant Signaling Network Research Center, and Technology Development Program for Agriculture and Forestry (309017-5) and, in part, by a grant to G.-T.K. from the Environmental Biotechnology National Core Research Center of MOST. The work was also supported by the Ministry of Education of the Czech Republic (LN00A081, LC06034, and MSM0021622415), the Academy of Sciences of the Czech Republic (IAA600380507), the Bundesministerium für Bildung und Forschung, and Fonds of Chemical Industry. J.H. was supported by the Deutscher Akademischer Austausch Dienst. S.M.C. was a recipient of a Brain Korea 21 fellowship.

Received March 3, 2009; revised June 16, 2009; accepted June 30, 2009; published July 21, 2009.

REFERENCES

- Aloni, R. (1987). Differentiation of vascular tissues. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **38**: 179–204.
- Altamura, M.M., Possenti, M., Matteucci, A., Baima, S., Ruberti, I., and Morelli, G. (2001). Development of the vascular system in the inflorescence stem of *Arabidopsis*. *New Phytol.* **151**: 381–389.
- Baucher, M., El Jaziri, M., and Vandeputte, O. (2007). From primary to secondary growth: Origin and development of the vascular system. *J. Exp. Bot.* **58**: 3485–3501.
- Brewer, P.B., Heisler, M.G., Hejatko, J., Friml, J., and Benkova, E. (2006). In situ hybridization for mRNA detection in *Arabidopsis* tissue sections. *Nat. Protocols* **1**: 1462–1467.
- Cano-Delgado, A., Yin, Y., Yu, C., Vafeados, D., Mora-Garcia, S., Cheng, J.C., Nam, K.H., Li, J., and Chory, J. (2004). BRL1 and BRL3 are novel brassinosteroid receptors that function in vascular differentiation in *Arabidopsis*. *Development* **131**: 5341–5351.
- Carlsbecker, A., and Helariutta, Y. (2005). Phloem and xylem specification: pieces of the puzzle emerge. *Curr. Opin. Plant Biol.* **8**: 512–517.
- Clough, S.J., and Bent, A.F. (1998). Floral dip: A simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* **16**: 735–743.
- D'Agostino, I.B., Deruere, J., and Kieber, J.J. (2000). Characterization of the response of the *Arabidopsis* response regulator gene family to cytokinin. *Plant Physiol.* **124**: 1706–1717.
- Dortay, H., Mehnert, N., Burkle, L., Schmulling, T., and Heyl, A. (2006). Analysis of protein interactions within the cytokinin-signaling pathway of *Arabidopsis thaliana*. *FEBS J.* **273**: 4631–4644.
- Emery, J.F., Floyd, S.K., Alvarez, J., Eshed, Y., Hawker, N.P., Izhaki, A., Baum, S.F., and Bowman, J.L. (2003). Radial patterning of *Arabidopsis* shoots by class III HD-ZIP and KANADI genes. *Curr. Biol.* **13**: 1768–1774.
- Eriksson, M.E., Israelsson, M., Olsson, O., and Moritz, T. (2000). Increased gibberellin biosynthesis in transgenic trees promotes growth, biomass production and xylem fiber length. *Nat. Biotechnol.* **18**: 784–788.
- Fisher, K., and Turner, S. (2007). PXY, a receptor-like kinase essential for maintaining polarity during plant vascular-tissue development. *Curr. Biol.* **17**: 1061–1066.
- Friml, J., Vieten, A., Sauer, M., Weijers, D., Schwarz, H., Hamann, T., Offringa, R., and Jurgens, G. (2003). Efflux-dependent auxin gradients establish the apical-basal axis of *Arabidopsis*. *Nature* **426**: 147–153.
- Friml, J., et al. (2004). A PINOID-dependent binary switch in apical-basal PIN polar targeting directs auxin efflux. *Science* **306**: 862–865.
- Fukuda, H. (2004). Signals that control plant vascular cell differentiation. *Nat. Rev. Mol. Cell Biol.* **5**: 379–391.
- Gao, Z., Wen, C.K., Binder, B.M., Chen, Y.F., Chang, J., Chiang, Y.H., Kerris III, R.J., Chang, C., and Schaller, G.E. (2008). Heteromeric interactions among ethylene receptors mediate signaling in *Arabidopsis*. *J. Biol. Chem.* **283**: 23801–23810.
- Grefen, C., Stadele, K., Ruzicka, K., Obrdliik, P., Harter, K., and Horak, J. (2008). Subcellular localization and in vivo interactions of the *Arabidopsis thaliana* ethylene receptor family members. *Mol. Plant* **1**: 308–320.
- Hartmann, U., Hohmann, S., Nettesheim, K., Wisman, E., Saedler, H., and Huijser, P. (2000). Molecular cloning of SVP: A negative regulator of the floral transition in *Arabidopsis*. *Plant J.* **21**: 351–360.
- Hejatko, J., Pernisova, M., Eneva, T., Palme, K., and Brzobohaty, B. (2003). The putative sensor histidine kinase CK11 is involved in female gametophyte development in *Arabidopsis*. *Mol. Genet. Genomics* **269**: 443–453.
- Higuchi, M., et al. (2004). In planta functions of the *Arabidopsis* cytokinin receptor family. *Proc. Natl. Acad. Sci. USA* **101**: 8821–8826.
- Hirakawa, Y., Shinohara, H., Kondo, Y., Inoue, A., Nakanomyo, I., Ogawa, M., Sawa, S., Ohashi-Ito, K., Matsubayashi, Y., and Fukuda, H. (2008). Non-cell-autonomous control of vascular stem cell fate by a CLE peptide/receptor system. *Proc. Natl. Acad. Sci. USA* **105**: 15208–15213.
- Hirose, N., Makita, N., Yamaya, T., and Sakakibara, H. (2005). Functional characterization and expression analysis of a gene, OsENT2, encoding an equilibrative nucleoside transporter in rice suggest a function in cytokinin transport. *Plant Physiol.* **138**: 196–206.
- Hirose, N., Takei, K., Kuroha, T., Kamada-Nobusada, T., Hayashi, H., and Sakakibara, H. (2008). Regulation of cytokinin biosynthesis, compartmentalization and translocation. *J. Exp. Bot.* **59**: 75–83.
- Horak, J., Grefen, C., Berendzen, K.W., Hahn, A., Stierhof, Y.D., Stadelhofer, B., Stahl, M., Koncz, C., and Harter, K. (2008). The *Arabidopsis thaliana* response regulator ARR22 is a putative AHP phospho-histidine phosphatase expressed in the chalaza of developing seeds. *BMC Plant Biol.* **8**: 77.
- Hutchison, C.E., Li, J., Argueso, C., Gonzalez, M., Lee, E., Lewis, M. W., Maxwell, B.B., Perdue, T.D., Schaller, G.E., Alonso, J.M., Ecker, J.R., and Kieber, J.J. (2006). The *Arabidopsis* histidine phosphotransfer proteins are redundant positive regulators of cytokinin signaling. *Plant Cell* **18**: 3073–3087.
- Hwang, I., and Sheen, J. (2001). Two-component circuitry in *Arabidopsis* cytokinin signal transduction. *Nature* **413**: 383–389.
- Jacobs, W.P. (1952). The role of auxin in differentiation of xylem around a wound. *Am. J. Bot.* **39**: 301–309.
- Kakimoto, T. (1996). CK11, a histidine kinase homolog implicated in cytokinin signal transduction. *Science* **274**: 982–985.
- Kim, H.J., Ryu, H., Hong, S.H., Woo, H.R., Lim, P.O., Lee, I.C., Sheen, J., Nam, H.G., and Hwang, I. (2006). Cytokinin-mediated control of leaf longevity by AHK3 through phosphorylation of ARR2 in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **103**: 814–819.
- Ko, J.H., Han, K.H., Park, S., and Yang, J. (2004). Plant body weight-induced secondary growth in *Arabidopsis* and its transcription phenotype revealed by whole-transcriptome profiling. *Plant Physiol.* **135**: 1069–1083.
- Kurakawa, T., Ueda, N., Maekawa, M., Kobayashi, K., Kojima, M., Nagato, Y., Sakakibara, H., and Kyozuka, J. (2007). Direct control of shoot meristem activity by a cytokinin-activating enzyme. *Nature* **445**: 652–655.
- Mahonen, A.P., Bishopp, A., Higuchi, M., Nieminen, K.M., Kinoshita, K., Tormakangas, K., Ikeda, Y., Oka, A., Kakimoto, T., and Helariutta, Y. (2006b). Cytokinin signaling and its inhibitor AHP6 regulate cell fate during vascular development. *Science* **311**: 94–98.
- Mahonen, A.P., Bonke, M., Kauppinen, L., Riikonen, M., Benfey, P. N., and Helariutta, Y. (2000). A novel two-component hybrid molecule regulates vascular morphogenesis of the *Arabidopsis* root. *Genes Dev.* **14**: 2938–2943.
- Mahonen, A.P., Higuchi, M., Tormakangas, K., Miyawaki, K., Pischke, M.S., Sussman, M.R., Helariutta, Y., and Kakimoto, T. (2006a). Cytokinins regulate a bidirectional phosphorelay network in *Arabidopsis*. *Curr. Biol.* **16**: 1116–1122.
- Matsumoto-Kitano, M., Kusumoto, T., Tarkowski, P., Kinoshita-Tsujimura, K., Vaclavikova, K., Miyawaki, K., and Kakimoto, T. (2008). Cytokinins are central regulators of cambial activity. *Proc. Natl. Acad. Sci. USA* **105**: 20027–20031.
- Medford, J.I., Horgan, R., El-Sawi, Z., and Klee, H.J. (1989). Alterations of endogenous cytokinins in transgenic plants using a chimeric isopentenyl transferase gene. *Plant Cell* **1**: 403–413.
- Miyawaki, K., Matsumoto-Kitano, M., and Kakimoto, T. (2004). Expression of cytokinin biosynthetic isopentenyltransferase genes in

- Arabidopsis: Tissue specificity and regulation by auxin, cytokinin, and nitrate. *Plant J.* **37**: 128–138.
- Müller, B., and Sheen, J.** (2008). Cytokinin and auxin interaction in root stem-cell specification during early embryogenesis. *Nature* **453**: 1094–1097.
- Nakamura, A., Kakimoto, T., Imamura, A., Suzuki, T., Ueguchi, C., and Mizuno, T.** (1999). Biochemical characterization of a putative cytokinin-responsive His-kinase, CKI1, from *Arabidopsis thaliana*. *Biosci. Biotechnol. Biochem.* **63**: 1627–1630.
- Nieminen, K., et al.** (2008). Cytokinin signaling regulates cambial development in poplar. *Proc. Natl. Acad. Sci. USA* **105**: 20032–20037.
- Nishimura, C., Ohashi, Y., Sato, S., Kato, T., Tabata, S., and Ueguchi, C.** (2004). Histidine kinase homologs that act as cytokinin receptors possess overlapping functions in the regulation of shoot and root growth in *Arabidopsis*. *Plant Cell* **16**: 1365–1377.
- Pernisova, M., Klima, P., Horak, J., Valkova, M., Malbeck, J., Soucek, P., Reichman, P., Hoyerova, K., Dubova, J., Friml, J., Zazimalova, E., and Hejatkó, J.** (2009). Cytokinins modulate auxin-induced organogenesis in plants via regulation of the auxin efflux. *Proc. Natl. Acad. Sci. USA* **106**: 3609–3614.
- Pischke, M.S., Jones, L.G., Otsuga, D., Fernandez, D.E., Drews, G. N., and Sussman, M.R.** (2002). An *Arabidopsis* histidine kinase is essential for megagametogenesis. *Proc. Natl. Acad. Sci. USA* **99**: 15800–15805.
- Prige, M.J., Otsuga, D., Alonso, J.M., Ecker, J.R., Drews, G.N., and Clark, S.E.** (2005). Class III homeodomain-leucine zipper gene family members have overlapping, antagonistic, and distinct roles in *Arabidopsis* development. *Plant Cell* **17**: 61–76.
- Sachs, T.** (2000). Integrating cellular and organismic aspects of vascular differentiation. *Plant Cell Physiol.* **41**: 649–656.
- Samuelson, M.E., Eliasson, L., and Larsson, C.M.** (1992). Nitrate-regulated growth and cytokinin responses in seminal roots of barley. *Plant Physiol.* **98**: 309–315.
- Schaller, G.E., Ladd, A.N., Lanahan, M.B., Spanbauer, J.M., and Bleecker, A.B.** (1995). The ethylene response mediator ETR1 from *Arabidopsis* forms a disulfide-linked dimer. *J. Biol. Chem.* **270**: 12526–12530.
- Scheres, B., Dilaurenzio, L., Willemsen, V., Hauser, M.T., Janmaat, K., Weisbeek, P., and Benfey, P.N.** (1995). Mutations affecting the radial organization of the *Arabidopsis* root display specific defects throughout the embryonic axis. *Development* **121**: 53–62.
- Surette, M.G., Levit, M., Liu, Y., Lukat, G., Ninfa, E.G., Ninfa, A., and Stock, J.B.** (1996). Dimerization is required for the activity of the protein histidine kinase CheA that mediates signal transduction in bacterial chemotaxis. *J. Biol. Chem.* **271**: 939–945.
- Szekeres, M., Nemeth, K., Koncz-Kalman, Z., Mathur, J., Kauschmann, A., Altmann, T., Redei, G.P., Nagy, F., Schell, J., and Koncz, C.** (1996). Brassinosteroids rescue the deficiency of CYP90, a cytochrome P450, controlling cell elongation and de-etiolation in *Arabidopsis*. *Cell* **85**: 171–182.
- Takei, K., Ueda, N., Aoki, K., Kuromori, T., Hirayama, T., Shinozaki, K., Yamaya, T., and Sakakibara, H.** (2004). AtIPT3 is a key determinant of nitrate-dependent cytokinin biosynthesis in *Arabidopsis*. *Plant Cell Physiol.* **45**: 1053–1062.
- To, J.P., Haberer, G., Ferreira, F.J., Deruere, J., Mason, M.G., Schaller, G.E., Alonso, J.M., Ecker, J.R., and Kieber, J.J.** (2004). Type-A *Arabidopsis* response regulators are partially redundant negative regulators of cytokinin signaling. *Plant Cell* **16**: 658–671.
- Tomomori, C., et al.** (1999). Solution structure of the homodimeric core domain of *Escherichia coli* histidine kinase EnvZ. *Nat. Struct. Biol.* **6**: 729–734.
- Vitha, S., Baluska, F., Braun, M., Samaj, J., Volkmann, D., and Barlow, P.W.** (2000). Comparison of cryofixation and aldehyde fixation for plant actin immunocytochemistry: Aldehydes do not destroy F-actin. *Histochem. J.* **32**: 457–466.
- Walter, M., Chaban, C., Schütze, K., Batistic, O., Weckermann, K., Nake, C., Blazevic, D., Grefen, C., Schumacher, K., Oecking, C., Harter, K., and Kudla, J.** (2004). Visualization of protein interactions in living plant cells using bimolecular fluorescence complementation. *Plant J.* **40**: 428–438.
- Werner, T., Kollmer, I., Bartrina, I., Holst, K., and Schumling, T.** (2006). New insights into the biology of cytokinin degradation. *Plant Biol.* **8**: 371–381.
- Werner, T., Motyka, V., Laucou, V., Smets, R., Van Onckelen, H., and Schumling, T.** (2003). Cytokinin-deficient transgenic *Arabidopsis* plants show multiple developmental alterations indicating opposite functions of cytokinins in the regulation of shoot and root meristem activity. *Plant Cell* **15**: 2532–2550.
- Yamada, H., Suzuki, T., Terada, K., Takei, K., Ishikawa, K., Miwa, K., Yamashino, T., and Mizuno, T.** (2001). The *Arabidopsis* AHK4 histidine kinase is a cytokinin-binding receptor that transduces cytokinin signals across the membrane. *Plant Cell Physiol.* **42**: 1017–1023.
- Yang, J., Zhang, J., Wang, Z., Zhu, Q., and Wang, W.** (2001). Hormonal changes in the grains of rice subjected to water stress during grain filling. *Plant Physiol.* **127**: 315–323.
- Ye, Z.H., Freshour, G., Hahn, M.G., Burk, D.H., and Zhong, R.** (2002). Vascular development in *Arabidopsis*. *Int. Rev. Cytol.* **220**: 225–256.
- Yokoyama, A., Yamashino, T., Amano, Y., Tajima, Y., Imamura, A., Sakakibara, H., and Mizuno, T.** (2007). Type-B ARR transcription factors, ARR10 and ARR12, are implicated in cytokinin-mediated regulation of protoxylem differentiation in roots of *Arabidopsis thaliana*. *Plant Cell Physiol.* **48**: 84–96.
- Zhao, C., Craig, J.C., Petzold, H.E., Dickerman, A.W., and Beers, E.P.** (2005). The xylem and phloem transcriptomes from secondary tissues of the *Arabidopsis* root-hypocotyl. *Plant Physiol.* **138**: 803–818.

The Histidine Kinases CYTOKININ-INDEPENDENT1 and ARABIDOPSIS HISTIDINE KINASE2 and 3 Regulate Vascular Tissue Development in Arabidopsis Shoots

Jan Hejatko, Hojin Ryu, Gyung-Tae Kim, Romana Dobesova, Sunhwa Choi, Sang Mi Choi, Premysl Soucek, Jakub Horak, Blanka Pekarova, Klaus Palme, Bretislav Brzobohaty and Ildoo Hwang
PLANT CELL 2009;21;2008-2021; originally published online Jul 21, 2009;
DOI: 10.1105/tpc.109.066696

This information is current as of September 11, 2009

Supplemental Data	http://www.plantcell.org/cgi/content/full/tpc.109.066696/DC1
References	This article cites 62 articles, 35 of which you can access for free at: http://www.plantcell.org/cgi/content/full/21/7/2008#BIBL
Permissions	https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&issn=1532298X&WT.mc_id=pd_hw1532298X
eTOCs	Sign up for eTOCs for <i>THE PLANT CELL</i> at: http://www.plantcell.org/subscriptions/etoc.shtml
CiteTrack Alerts	Sign up for CiteTrack Alerts for <i>Plant Cell</i> at: http://www.plantcell.org/cgi/alerts/ctmain
Subscription Information	Subscription information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at: http://www.aspb.org/publications/subscriptions.cfm