



## A spatial model of tree $\alpha$ -diversity and tree density for the Amazon

HANS TER STEEGE<sup>1,2,\*</sup>, NIGEL PITMAN<sup>3</sup>, DANIEL SABATIER<sup>4</sup>,  
HERNAN CASTELLANOS<sup>5</sup>, PETER VAN DER HOUT<sup>6</sup>, DOUGLAS C.  
DALY<sup>7</sup>, MARCOS SILVEIRA<sup>8</sup>, OLIVER PHILLIPS<sup>9</sup>, RODOLFO  
VASQUEZ<sup>10</sup>, TINDE VAN ANDEL<sup>1</sup>, JOOST DUIVENVOORDEN<sup>11</sup>,  
ALEXANDRE ADALARDO DE OLIVEIRA<sup>12</sup>, RENSKÉ EK<sup>1</sup>, RAMESH  
LILWAH<sup>13</sup>, RAQUEL THOMAS<sup>14</sup>, JESSICA VAN ESSEN<sup>15</sup>, CLAUDIA  
BAIDER<sup>16</sup>, PAUL MAAS<sup>1</sup>, SCOTT MORI<sup>6</sup>, JOHN TERBORGH<sup>3</sup>, PERCY  
NÚÑEZ VARGAS<sup>17</sup>, HUGO MOGOLLÓN<sup>18</sup> and WILFRIED MORAWETZ<sup>19</sup>

<sup>1</sup>National Herbarium of the Netherlands (NHN), Utrecht University Branch, Heidelberglaan 2, 3584 CS Utrecht, The Netherlands; <sup>2</sup>Department of Natural Resources, International Institute for Geo-information and Earth Observation (ITC), P.O. Box 6, 7500 AA Enschede, The Netherlands; <sup>3</sup>Center for Tropical Conservation, Duke University, Box 90381, Durham, NC 27708-0381, USA; <sup>4</sup>Institut de Recherche pour le Développement (IRD), 911 Avenue Agropolis, BP 5045, F-34032 Montpellier Cedex 1, France; <sup>5</sup>Universidad Nacional Experimental de Guayana, Calle Chile, Urbaniz Chilemex, Puerto Ordaz, Edo. Bolívar, Venezuela; <sup>6</sup>Department of Plant Ecology, Utrecht University, P.O. Box 80084, 3504 TB Utrecht, The Netherlands; <sup>7</sup>Institute of Systematic Botany, The New York Botanical Garden, Bronx, NY 10458-5126, USA; <sup>8</sup>Departamento de Ciências da Natureza/Parque Zoobotânico, Universidade Federal do Acre, BR 354, km 05, Rio Branco, Acre 69915-900, Brazil; <sup>9</sup>Centre for Biodiversity and Conservation, School of Geography, University of Leeds, Leeds LS2 9JT, UK; <sup>10</sup>Proyecto Flora del Perú, Jardín Botánico de Missouri, Jaen, Cajamarca, Peru; <sup>11</sup>Institute for Biodiversity and Ecosystem Dynamics, Universiteit van Amsterdam, Kruislaan 318, 1098 SM Amsterdam, The Netherlands; <sup>12</sup>Lab. Botânica, Universidade Paulista, Av. Paulista 900 1º andar, 01310-100 São Paulo-SP, Brazil; <sup>13</sup>Environmental Protection Agency, IAST Building, Turkyen Campus, Greater-Georgetown, Guyana; <sup>14</sup>Iwokrama International Centre for Rain Forest Conservation and Development, 67 Bel Air, P.O. Box 10630, Georgetown, Guyana; <sup>15</sup>Department of Comparative Physiology, Utrecht University, Padualaan 14, 3584 CH Utrecht, The Netherlands; <sup>16</sup>Universidade de São Paulo, IB/Ecologia, Rua do Matão/Trav. 14, 321, 05508-900 São Paulo-SP, Brazil; <sup>17</sup>Herbario Vargas, Universidad Nacional San Antonio de Abad de Cusco, Cusco, Peru; <sup>18</sup>NUMASHIR, Fundación para la Conservación de Ecosistemas Amenazados, Casilla postal 17 01 9149, Quito, Ecuador; <sup>19</sup>Institut für Botanik – Spezielle Botanik, Universität Leipzig, Johannisallee 21–23, D-04103 Leipzig, Germany; \*Author for correspondence (e-mail: h.tersteegen@bio.uu.nl; fax: +31-30-251-8061)

Received 11 July 2002; accepted in revised form 9 December 2002

**Key words:** Amazon, Guayana Shield, Kriging, Planning inventories, Quantile regression, Tree  $\alpha$ -diversity, Tree density

**Abstract.** Large-scale patterns of Amazonian biodiversity have until now been obscured by a sparse and scattered inventory record. Here we present the first comprehensive spatial model of tree  $\alpha$ -diversity and tree density in Amazonian rainforests, based on the largest-yet compilation of forest inventories and bolstered by a spatial interpolation technique that allows us to estimate diversity and density in areas that have never been inventoried. These data were then compared to continent-wide patterns of rainfall seasonality. We find that dry season length, while only weakly correlated with average tree  $\alpha$ -diversity, is a strong predictor of tree density and of maximum tree  $\alpha$ -diversity. The most diverse forests for any given DSL are concentrated in a narrow latitudinal band just south of the equator, while the least diverse

forests for any given DSL are found in the Guayana Shield and Amazonian Bolivia. Denser forests are more diverse than sparser forests, even when we used a measure of diversity that corrects for sample size. We propose that rainfall seasonality regulates tree  $\alpha$ -diversity and tree density by affecting shade tolerance and subsequently the number of different functional types of trees that can persist in an area.

## Introduction

Biodiversity mapping and modelling is an increasingly useful tool for ecology and conservation, but it has not yet produced a reliable picture of diversity patterns in the world's most species-rich forest (Gentry 1988a, 1988b; Phillips et al. 1994; Gaston 2000; Myers et al. 2000; Condit et al. 2002).

The Amazon, which harbours a very significant portion of the world's diversity (Myers et al. 2000), is a perfect natural laboratory for the study of tree diversity because of its size and its bio-physical gradients. The number of plots has risen steadily over the years and is probably much greater than in any other tropical region. Most studies examining the patterns of plant diversity in the Neotropics, however, have relied heavily on plots in western Amazonia (Gentry 1988a, 1988b; Phillips et al. 1994; Clinebell et al. 1995; Terborgh and Andresen 1998; Givnish 1999; ter Steege et al. 2000). In western Amazonia species richness is found to be strongly correlated with total annual rainfall (Gentry 1988a, 1988b), often considered a proximate estimate of ecosystem productivity (Gentry 1988a, 1988b; Phillips et al. 1994; Clinebell et al. 1995). The rainfall–diversity relationship has been confirmed by other studies outside the Amazon (Hall and Swaine 1976; Huston 1980, 1994). In each of these studies fertility was negatively correlated with rainfall, which has complicated the search for single causal relationships. Ter Steege et al. (2000) showed that, when the whole of the Amazon was taken into account, rainfall was a poor estimator of tree  $\alpha$ -diversity in 1-ha plots.

A closer inspection of the data of ter Steege et al. (2000) suggests that while rainfall and rainfall seasonality may not be good estimators of average tree  $\alpha$ -diversity, they may be good estimators of maximum diversity. As many more factors may have an effect on tree  $\alpha$ -diversity, these may obscure this relationship. If such factors interact rather than act in an additive manner, standard regression methods are not capable of detecting proper relationships. Recently, a more advanced method has become available for ecology (Cade et al. 1999) that has been pioneered in the field of econometrics (Koenker and Bassett 1978). With this technique, quantile regression, interacting factors may be studied even if not all such factors have been measured. Further details can be found in the Methods section.

Since our previous publication (ter Steege et al. 2000) much more plot data have become available and in this paper we have two main objectives:

1. Use a statistical model to generate an improved, most accurate spatial model to date, of tree  $\alpha$ -diversity of the Amazon, including an error map that may assist to plan future plot establishment. The results of this model will be available on our website (see below) and regularly updated.
2. Describe the rainfall–diversity model as a model of limiting factors, whereby

rainfall seasonality is one of the main factors limiting tree  $\alpha$ -diversity at a regional level, other factors being interactive with it.

We have no *a priori* assumptions what the most important other interacting factors might be, but by eliminating plots on ‘special’ soil types, such as white sands, floodplains and swamps, that generally have lower diversity than *terra firme* forest plots (ter Steege et al. 2000), we already eliminate some sources of variation. If the unknown factor(s) act at a regional level, mapping them should reveal a geographical pattern. Hence, we also map the other total of unknown limiting factors to discern this.

## Methods

The total number of plots with data, presently known to us in the Amazon, which here includes the forest region of the Guayana Shield, is 425.<sup>1</sup> For the analyses we used forest plots on *terra firme* only, leaving out plots on white sands, floodplains and swamps, plots outside the Amazon, and plots with incomplete data. The final selection comprised 275 plots. This set mainly included 1-ha plots, but in areas of low plot density also smaller (plots that had more than 150 individuals of trees with a dbh  $\geq 10$  cm) or larger ones. Most (228) plots were 1 ha in size, 18 plots were between 0.25 and 1 ha, 20 plots were between 1 and 4 ha, and 9 were plotless line samples with 200–800 individuals (equivalent to ca. 0.4–1.6 ha). To standardize  $\alpha$ -diversity we used Fishers’s  $\alpha$  (Fisher et al. 1943), which corrects well for variation in sample size (Taylor et al. 1976; Leigh 1995; Rosenzweig 1995; Condit et al. 1998).

Tree density was calculated on a hectare basis. For the plotless samples the tree density was extracted from the publications concerned, if this data was presented there. Otherwise the samples were omitted for the density estimates.

### *Mapping tree $\alpha$ -diversity*

Because the plots over-sample some areas of Amazonia and under-sample others, we used a spatial model based on universal kriging (Burrough and McDonnell 1998; Kalunzy et al. 1998) to estimate diversity in  $1^\circ$  cells across the region. The theory and process is explained briefly below:

The regionalized variable theory assumes that the spatial variation of any variable  $Z$  at location  $x$  can be expressed as the sum of a structural component ( $m$ : a constant mean or a trend), a random but spatially correlated component ( $\varepsilon'$ : the regionalized variable), and a residual error term ( $\varepsilon''$ ), that is not spatially correlated (Burrough and McDonnell 1998):

---

<sup>1</sup> This was the number when the analyses were carried out. During manuscript review a few more plots came to our knowledge. They are listed in the Appendix and will be used in the updated map on our website.

$$Z = m(x) + \varepsilon'(x) + \varepsilon''$$

We used a loess regression to detect the structural component ( $m$ ). The loess regression explained 64% of the variation in Fisher's  $\alpha$ . The residuals from the loess regression are equal to  $\varepsilon'(x) + \varepsilon''$  and were analysed with variogram analysis (Burrough and McDonnell 1998; Kalunzy et al. 1998). In the residuals, a clear spatial structure was present, and this improved very much after the removal of two outlier plots. An exponential model fitted the variogram data well and showed autocorrelation up to ca.  $2.2^\circ$ . As a consequence, interpolation was carried out with the exponential variogram results and with inclusion of a spatial trend model. This is known as universal kriging.

The statistical model used by kriging also produces error estimates. These errors were also mapped for inspection and may serve as a useful tool for planning further exploration.

Estimates are only presented for the forest area of the region based on an interpretation of NOAA-AVHRR NDVI satellite data (<http://terra.whrc.org/>). This area includes 567  $1^\circ$  grid cells.

To examine the effect of spatial and ecological variables on diversity we used the kriging model predictions for each  $1^\circ$  grid cell. We took this course because several sites had many points close together, violating a main assumption of regression analysis – independence of data (this assumption is opposite to the requirement of kriging, which assumes and then estimates spatial dependence in the data). We tested the accuracy of the model's estimates by removing each single plot from the dataset, allowing the model to re-estimate the diversity of its cell with the remaining dataset, and then comparing the estimated diversity (for the grid cell) with the actual diversity of the removed plot. In this way the plot's own value has no influence on its estimated value. Thus, the complete interpolation process was carried out 275 times. We will call this the jackknifed estimation (a more correct term might be the studentized estimation – Venables and Ripley 1998).

#### *Rainfall and tree $\alpha$ -diversity*

To study the effect of rainfall on tree  $\alpha$ -diversity we used dry season length (DSL), which was defined as the number of months with  $<100$  mm of rain. DSL was extracted for every  $1^\circ$  grid cell in the region from a digitised map (data from Sombroek 2001; Figure 1). Sombroek (2001) did not distinguish between 0 and 1 month(s) of dry period and for calculations we classified both as zero.

Limiting factor theory assumes that factors influencing the dependent variable act heterogeneously (read: independently) over the landscape. Near the upper bound of the distribution, the factor under consideration is considered to be actively limiting the dependent variable. Variation below these maxima is caused by the interacting influence of other (often unmeasured) factors. We used quantile regression (Koenker and Bassett 1978; Cade et al. 1999) to examine whether DSL was correlated with maximum diversity and tree density (Cade et al. 1999). We used the module *regq*, written for S-plus (Cade et al. 1999). Two important assumptions of quantile




Figure 1. Map of DSL (data after Sombroek 2001), expressed as the number of months with <100 mm of rain.

regression are the existence of other interacting limiting factors and the fact that the dependent variable will attain its true maximum for the limiting factor (the measured independent variable) under investigation. While there are several factors that can be thought to influence diversity (disturbance, soil type (which was already eliminated)), we have no absolute proof of maximum tree  $\alpha$ -diversity anywhere in the Amazon. However, it is difficult to imagine plots richer than some in western or central Amazonia where plots of ca. 600 individuals may contain ca. 300 species (e.g.: Gentry 1988a; de Oliveira and Mori 1999; Vasques-Martinez and Phillips 2000).

We used an interference interaction model (Cade et al. 1999), in linear form, where tree  $\alpha$ -diversity ( $Z$ ) is a linear (negative) function of DSL and its interaction with OIF:

$$Z = \alpha + \beta(1 - \text{OIF})\text{DSL} + \varepsilon$$

If  $\text{OIF} = 0$  (the upper bound of the distribution for DSL) this reduces to

$$Z_{\max} = \alpha + \beta(\text{DSL}) + \varepsilon$$

This is equal to the maximum  $Z$  for a given DSL (in other words, it is the real relationship between DSL and average tree  $\alpha$ -diversity for the grid cells). The parameters  $\alpha$  and  $\beta$  were estimated by calculating the regression equation for the

90th percentile, where OIF approaches zero. To study whether OIF is either related to climatic factors or has an otherwise geographic pattern its estimated value was calculated as

$$\text{OIF} = (Z_{\text{max}} - Z)\beta\text{DSL} + \varepsilon'$$

## Results

### *Mapping diversity and density*

Both the raw and the interpolated plot data show the region's peak of average tree  $\alpha$ -diversity as a large swath of mostly a-seasonal forest stretching along the equator from Ecuador, northern Peru, and southern Colombia into central Amazonian Brazil (Figure 2). *Terra firme* tree  $\alpha$ -diversity is highest in the vicinity of Iquitos, Peru, and Manaus, Brazil and lowest in Bolivia and the Guayana Shield forests, with the exception of central-south French Guiana.

Average tree density for 1° grid cells in Amazonia varies from ca. 400 to 750 trees ha<sup>-1</sup> and shows a distribution similar to that of tree  $\alpha$ -diversity, but with a more pronounced peak in western Amazonia (Figure 3). Forests in western Amazonia are on average 150 stems ha<sup>-1</sup> denser than those in eastern Amazonia.

The accuracy of the spatial model is high and the jackknifed estimation explains 80% of variance in the diversity of the removed plots ( $n = 275$ , data not shown).

The errors in the spatial model are largest where there are few plots in the 'neighbourhood' (Figure 4). These areas include the region in Pará below the Guianas, the Tapajós River basin, central Colombian Amazon, and the border region of northwestern Peru and Ecuador.

### *Rainfall, tree $\alpha$ -diversity and density*

We find DSL to be negatively correlated with diversity and density. However, DSL is a much stronger predictor of estimated maximum diversity (Figure 5;  $R^2 = 0.91$ ,  $P \ll 0.001$  for the 90th quantile) than of estimated (average) diversity ( $R^2 = 0.35$ ,  $P \ll 0.001$ ,  $n = 567$  grid cells). This same pattern is also present in the raw plot data (Figure 1, inset), the only difference being that very high local values have been smoothed out by the spatial interpolation.

The effect of the OIF is in fact of such magnitude that minimum tree  $\alpha$ -diversity is almost equal for each DSL. The regression line for the 10th quantile is nearly horizontal, its slope still significantly different from zero, but more so very significantly different from the slope of the 90th percentile. This indeed supports the existence of interaction factors acting on tree  $\alpha$ -diversity.

Average tree density is more strongly under control of seasonality than is tree  $\alpha$ -diversity (Figure 6;  $R^2 = 0.52$ ,  $P \ll 0.001$ ). The higher and lower quantile regression lines are nearly parallel (Figure 6), suggesting that seasonality is the main effect, with other factors being mostly additive, rather than interacting.




Figure 2. Map of average tree  $\alpha$ -diversity (Fisher's  $\alpha$ ) based on 275 plots in *terra firme* forest in the Amazon and Guayana Shield rain forest area mapped on a  $1^\circ$  grid cell scale. The extent of tropical wet/moist and semi-deciduous forest area is based on NOAA-AVHRR NDVI satellite data (<http://terra.whrc.org/>). Diversity is highest in a narrow band from western Amazonia towards central Amazonia with peaks around Iquitos and Manaus. Dots indicate the maximum Fisher's  $\alpha$  found at one location ( $n = 275$ ). Because many plots may be in close proximity (and overlapping), much less than 275 dots are visible.

Even though Fisher's  $\alpha$  values are standardized for density (at the plot level), we still find a relatively strong positive relationship between interpolated tree density and interpolated Fisher's  $\alpha$  for the 567 grid cells (Figure 7;  $R^2 = 0.46$ ,  $P \ll 0.001$ ).

A spatial null-hypothesis is that grid cells with average Fisher's  $\alpha$  above the 90th quantile and those below the 10th (Figure 5) are randomly distributed over the spatial gradient of DSL (cf. Figure 1). When we map these grid cells, those with the highest diversity for any DSL regime (OIF = 0) are found within a single band in central Amazonia, just south of the equator. The least diverse grid cells, where seasonality is only of limited influence on tree  $\alpha$ -diversity (OIF = maximum) are concentrated in two geographically distinct areas: the northwestern Guayana Shield and Bolivia. In Figure 8 this is shown in a slightly different manner as the strength of the OIF between 0 and 100%. The band where the OIF are of little influence is clearly visible below the Amazon River.

The OIF have a strong unimodal relationship with latitude, peaking at  $5^\circ$  S (Figure 9). This unimodal pattern is obvious with and without the data related to the




Figure 3. Map of average tree density (trees  $\geq 10$  cm dbh  $\text{ha}^{-1}$ ) across the Amazon. Forests in western Amazonia have on average 150 more stems per ha than those in eastern Amazonia.




Figure 4. Map of standard errors in tree  $\alpha$ -diversity (Fisher's  $\alpha$ ) as produced by the spatial model obtained by universal kriging (Figure 2).






Figure 5. Modelled Fisher's  $\alpha$  as a function of DSL. An increase in dry months strongly limits the average maximum tree  $\alpha$ -diversity found at a site. Dark green: grid cells with Fisher's  $\alpha$  over the 90th quantile (rainfall seasonality is the main limiting factor to tree  $\alpha$ -diversity); light green: grid cells with Fisher's  $\alpha$  below the 10th quantile (rainfall seasonality has no marked effect on tree  $\alpha$ -diversity in these grid cells). The slopes of the two regression lines differ significantly. The inset shows the same relationship for actual plot data with least squares regression to show a weak overall relation between DSL and tree  $\alpha$ -diversity.

10th quantile (Figure 9) and also if data is plotted for each DSL value separately (data not shown). In other words, even if rainfall has been corrected for (correlation between OIF and DSL is non-existent –  $R^2 = 0.003$ ), there is still a very strong regional spatial pattern in tree  $\alpha$ -diversity, peaking at 5° S.

## Discussion

### *Mapping tree $\alpha$ -diversity and tree density in the Amazon*

As the accuracy of the spatial model, tested with the jackknifed estimation was high, this suggests that the tree  $\alpha$ -diversity of unmeasured areas can be estimated with a fairly high level of accuracy. The same pattern as shown in Figure 2 was also apparent in an earlier diversity map (based on inverse distance weighting) drawn with data from >100 fewer plots (ter Steege et al. 2000), also suggesting that the pattern may be robust despite the immense areas still unexplored by botanists and ecologists in Amazonia. New data not yet included in the map (Henkel (in press); D. Sabatier et al., unpublished data; J. Schulz and J. Lindeman, unpublished data; A. Duque, unpublished data) also correspond well to the modelled pattern.




Figure 6. Average tree density (trees  $\text{ha}^{-1}$ ) of Amazonian  $1^\circ$  grid cells as a function of DSL. Dark brown: grid cells with density higher than the 90th quantile; light brown: grid cells with density below the 10th quantile. Although the slopes of the 90th and 10th quantile differ significantly, they are practically similar in absolute value, suggesting that seasonality is a main factor influencing tree density in the Amazon.






Figure 7. Fisher's  $\alpha$  is still influenced by average tree density in  $1^\circ$  grid cells, suggesting that density has a regional effect of diversity. This effect is present in the high, medium and low quantile data (see Figure 4).



*Figure 8.* Spatial distribution of the OIF. OIF is displayed as a percentage of its maximum: 100 indicates interacting factors at maximum and rainfall not important in determining tree  $\alpha$ -diversity; 0 indicates that rainfall is the major limiting factor for tree  $\alpha$ -diversity.



*Figure 9.* Latitude vs. OIF. OIF are least limiting (most positive) around 5° S, the central band of the Amazon forest region.

The pattern is in stark contrast, however, with previous maps of diversity in the Amazon (Prance 1982; Conservation International 1990) and suggests that the main diversity pattern in the Amazon may be much more simple in its spatial configuration, with consequences for inferred forest history (see also Bush 1994).

*Errors are useful for planning*

Although the general prediction of tree  $\alpha$ -diversity of excluded plots by the model was quite accurate, errors are not equally distributed over the map. Errors in the spatial model are largest where there are few plots in the 'neighbourhood' (Figure 4). As mentioned above, these areas include the region in Pará below the Guianas, the Tapajós River basin, central Colombian Amazon, and the border region of northwestern Peru and Ecuador. Mapping of diversity in the Amazon would benefit most from new plots in these areas. In the southern part of Colombia (Leticia) and adjacent Brazil, errors are slightly smaller and, in addition to that, new plots have already been established there (A. Duque, personal communication).

*The main pattern: drought is limiting diversity but what is cause and what is correlation?*

DSL was a much stronger predictor of estimated maximum diversity than of estimated (average) diversity on the modelled and raw data (Figure 5), as predicted by limiting factor theory. Based on the results of the quantile regression we suggest that DSL has a strong (linear) effect on diversity, an effect that is masked by OIF in the full dataset. The effect of the OIF is in fact of such magnitude that minimum tree  $\alpha$ -diversity is almost equal for each DSL.

While our results are in general agreement with previous studies (Hall and Swaine 1976; Huston 1980; Phillips et al. 1994; Clinebell et al. 1995), the strong relationship between maximum tree  $\alpha$ -diversity and DSL and potential interacting limiting factors demonstrates why researchers see such contrasting results when they compare rainfall regime and tree diversity – they look at mean diversity, not maximum potential diversity. Secondly, our results support hypotheses that consider the effect of the length of the growing season and number of potential co-existing functional types, which can be considered a proxy measure for diversity. Recently, two such mechanistic models have shown that, at a global level, the length of the growing season may indeed determine the number of functional types, and hence species, that can exist (Kikuzawa 1996; Kleidon and Mooney 2000).

We suggest that, in the Amazon, density and diversity may be driven by the same ecological factor (see also Pitman 2000): DSL. High constant moisture is thought to increase shade tolerance, higher densities, and subsequently a higher number of functional guilds in the understorey (Huston 1994). Indeed, in western Amazonia the higher density (of individuals  $\geq 10$  cm dbh) is a result of higher density of smaller understorey trees (Pitman 2000) and this effect may even be stronger for yet smaller size classes (Givnish 1999 – who did *not* find the effect in the size classes  $\geq 10$  cm dbh in the 'Gentry dataset', however! If our thoughts are correct, these

'extra' individuals should, at least for a significant part, consist of other species and of other functional guilds, increasing diversity more than if they would consist of seedlings and saplings of the overstorey trees alone. This would also explain that even when we correct for density by using Fisher's  $\alpha$  there is still an effect of average tree density on tree  $\alpha$ -diversity (Figure 7).

We do not have complete data for all plots (individual trees and their diameters) but propose a simple test for this hypothesis. If indeed an increased diversity in western Amazonia is caused by the higher diversity of small individuals, calculating Fisher's  $\alpha$  separately for individuals  $<20$  cm and for trees  $\geq 20$  cm should reveal a difference when carried out for eastern, central and western Amazonia.

One other causal factor that is in accordance with the results, are insect and fungal pressure, which are both thought to increase strongly with (more predictable, constant) moisture and mediate tree diversity through species-specific density-dependent mortality (Givnish 1999; E.G. Leigh, personal communication; but see Hille Ris-Lambers et al. 2000; ter Steege and Zagt 2002). This prediction should also lead to testable hypotheses in dry and wet areas with maximum diversity.

#### *Other spatial causes*

The OIF have a strong unimodal relationship with latitude (Figures 8 and 9), peaking at  $5^\circ$  S, and this unimodal relationship is not related to rainfall. We suggest that at present this maximum diversity (for any given seasonality) in greater Amazonia is best explained by a stochastic null-model (Colwell and Lees 2000) which predicts that even in the absence of an environmental gradient, the highest number of species will be found in the central portion of a geographical domain, which for the Amazon forest coincides largely with a band around  $5^\circ$  S. This higher 'regional species pool' results in higher potential tree  $\alpha$ -diversity in plots as well, which is our measure used.

In addition to that, we hypothesise that the consistently low diversity in the Guianas may also be the result of the small size of this area, compared to the major Amazon forest block (see Terborgh 1973; Rosenzweig 1995; ter Steege et al. 2000). The Bolivian area includes many wet savannahs and may have become forested only very recently in geological time (Mayle et al. 2000), which may also contribute to its low diversity and high abundance of well-dispersed families with several fast growing species, such as Moraceae (ter Steege et al. 2000).

#### *A caution for conservation planning*

While the data presented increase our insight into potential factors influencing tree  $\alpha$ -diversity, we caution the use of diversity maps as a single source of data for conservation planning. Although the Guayana Shield and the Bolivian area have much lower diversity than central and western Amazonian areas, their species are not just a sub-sample of the latter area. In addition to that, within the areas of equal diversity, gradients in species composition ( $\beta$ -diversity) are continuously present (western Amazonia and Panama: see Condit et al. 2002; Guayana Shield: D.

Sabatier, unpublished data; H. ter Steege, unpublished data). Such information is also crucial if a representative portion of the area's diversity is to be protected.

#### *A bit about the future*

The map and some associated data will be published on the website of our Amazonian Plot Network, accessible through: <http://www.bio.uu.nl/~herba/research.html>.

It is our intention to update this web-based map regularly as newly published plot data become available. Some of us and colleagues have already started with the establishment of new plots in areas where coverage is low (Pitman et al., Peru, Ecuador) or are planning to do so (C. Peres, Tapajós and other Brazilian parts; B. Hoffman, South Suriname). We would also welcome and acknowledge plot data sent to us by others.

As there are much fewer plots on floodplains (64) and even fewer in swamps (20) and on white sand podzols (14), maps for these plots are not well feasible (for some information see ter Steege et al. 2000). On our website a preliminary map of floodplain diversity will be present. We especially encourage the establishment of plots in these less diverse ecosystems.

Further research plans include the role of neutral models and ecology in patterns of  $\beta$ -diversity, especially in western Amazonia and the Guayana Shield, two areas for which the comparative taxonomy for a great number of plots can be resolved with the least difficulties. To further resolve the role of ecology in shaping tree composition of forest communities (and plots therein) it is also necessary to establish plots in the various forest systems found.

For more information please visit: <http://www.bio.uu.nl/~herba/research.html>.

#### **Acknowledgements**

The start of this project was supported by the Tropenbos Guyana Programme, through grant 'Conservation and Sustainable Use of Botanical Diversity in Guyana' (B7-6201/98-13/FOR) of the European Community. We thank all those who worked with us in the field for their support and Peter Burrough, Alfred de Gier, Wim Dijkman, Karl Eichhorn, Gerrit Heil, Mark van Nieuwstadt, Marinus Werger, René Verburg and Roderick Zagt and two anonymous reviewers for comments on previous versions of the manuscript.

#### **Appendix. References to plot data in the Amazon.**

Absy M.L., Prance G.T. and Barbosa E.M. 1987. Inventário florístico de floresta natural na área da estrada Cuiabá-Porto Velho (BR - 364). *Acta Amazonica* 16/17 (Supl.): 85–121.

Almeida S.S., Lisboa P.L.B. and Silva A.S. 1993. Diversidade florística de uma comunidade arbórea na estação científica 'Ferreira Pena', em Caxiuanã (Pará). Boletim do Museu Paraense Emílio Goeldi. Serie Botânica 9: 93–128.

Alonso A., Dalmeijer F., Mistry S., Ros C., Udvardy S., Comiskey J., Núñez P., Chuspe M.E., Baldeón S., Betrán H., Santistebán J. and Valencia G. 1998. Flora I. Biodiversity assessment and long term monitoring, lower Urubamba Region. In: Alonso A. and Dalmeijer F. (eds.), Biodiversity Assessment in the Lower Urubamba Region, Perú. SI/MAB Series #2. Smithsonian Institution, Washington, DC.

Amaral I.L. 1996. Diversidade florística em floresta de terra firme, na região de Urucu – AM. Dissertação de Mestrado, Instituto Nacional de Pesquisas da Amazônia (INPA) /Fundação Universidade do Amazonas (FUA). Manaus, Brazil.

Amaral I.L., Matos F.D.A. and Lima J. 2000. Composição florística e parâmetros estruturais de um hectare de floresta densa de terra firme no Rio Uatumã, Amazonia, Brasil. Acta Amazonica 30: 377–392.

Ayres J.M. 1981. Observações sobre a ecologia e o comportamento do Cuciús (*Chiropotes satanes* e *C. satanus*). M.Sc. dissertation, INPA, Manaus, Brazil.

Ayres J.M. 1993. As matas da Várzea do Mamirauá. MCT-CNPq/Sociedade Civil Mamirauá, Brasília, Brazil.

Balee W. 1986. Análise preliminar de inventário florestal e a etnobotânica Ka'apor (Maranhão). Boletim do Museu Paraense Emílio Goeldi. Serie Botânica 2: 141–167.

Balee W. 1987. A etnobotânica quantitativa dos índios Tembé (Rio Gurupi, Pará). Boletim do Museu Paraense Emílio Goeldi. Serie Botânica 3: 29–50.

Balee W. and Campbell D.G. 1990. Evidence for the successional status of liana forest (Xingu river basin, Amazonian Brazil). Biotropica 22: 36–47.

Balslev H., Luteyn J., Ollgaard B. and Holm-Nielsen L.B. 1987. Composition and structure of adjacent unflooded and floodplain forest in Amazonian Ecuador. Opera Botanica 92: 37–57.

Balslev H., Valencia R., Paz Y Mino G., Christensen H. and Nielsen I. 1998. Species count of vascular plants in one hectare of humid lowland forest in Amazonian Ecuador. In: Dallmeier F. and Comiskey J.A. (eds), Forest Biodiversity in North, Central and South America, and the Caribbean. Research and Monitoring. MAB Series Vol. 21. UNESCO, Paris, pp. 585–594.

Barbosa E.M. 1988. Análise estrutural de uma floresta natural na reserva de pesquisa ecológica do INPA em Ouro Preto do Oeste – Rondônia. M.Sc. Thesis, INPA, Manaus, Brazil.

Bergmans F. and Vroomans V. 1995. Availability and sustainability of forest resources: a research to the structure and species composition of a north Bolivian Amazon forest and its actual and future resource availability. Thesis, Larenstein Agricultural College, Velp, The Netherlands.

Black G.A., Dobzhansky T.H. and Pavan C. 1950. Some attempts to estimate species diversity and population density of trees in Amazonian forests. Botanical Gazette 111: 413–425.

Boom B.M. 1986. A forest inventory in Amazonian Bolivia. Biotropica 18: 287–294.

Boubli J.P. 1999. Feeding ecology of black-headed Uacaris (*Cacajó melanocephalus melanocephalus*) in Pico da Neblina National Park, Brazil. *International Journal of Primatology* 20: 719–749.

Cain S.A., De Oliveira Castro M., Murça Pires J. and Da Silva N.T. 1956. Application of some phytosociological techniques to Brazilian rain forest. *American Journal of Botany* 43: 911–941.

Campbell D.G., Daly D.C., Prance G.T. and Ubirajara N.M. 1986. Quantitative ecological inventory of terra firme and varzea tropical forest on the Rio Xingu, Brazilian Amazon. *Brittonia* 38: 369–393.

Campbell D.G., Stone J.L. and Rosas A. 1992. A comparison of the phytosociology and dynamics of three floodplain (varzea) forests of known ages, Rio Jurua, western Brazilian Amazon. *Botanical Journal of the Linnean Society* 108: 213–237.

Castellanos H.G. 1998. Floristic composition and structure, tree diversity, and the relationship between floristic distribution and soil factors in El Caura Forest Reserve, southern Venezuela. In: Dallmeier F. and Comiskey J.A. (eds), *Forest Biodiversity in North, Central and South America, and the Caribbean. Research and Monitoring. MAB Series Vol. 21.* UNESCO, Paris, pp. 507–533.

Cerón C.E. and Montalvo C.A. 1997. Composición y estructura de una hectárea de bosque en la Amazonía Ecuatoriana – con información etnobotánica de los Huaorani. In: Valencia R. and Balslev H. (eds), *Estudios sobre diversidad de plantas. Memorias de II congreso Ecuatoriana de Botánica.* Pontificia Universidad Católica del Ecuador, Quito, Ecuador, pp. 153–172.

Cerón C.E., Fernández D.M., Jiménez E.D. and Pillojo I. 2000. Composición y estructura de una igapo Ecuatoriano. *Cinchona* 1: 41–70.

Comiskey J.A., Aymard G. and Dallmeier F. 1994. Structure and composition of lowland mixed forest in the Kwakwani region of Guyana. *Biollonia* 10: 13–28.

Dallmeier F., Kabel M. and Foster R.B. 1996. Floristic composition, diversity, mortality and recruitment of different substrates: lowland tropical forest, Pakitza, Rio Manu, Peru. In: Wilson D.E. and Sandoval A. (eds), *Manu: The Diversity of Southeastern Peru.* Smithsonian Institution, Washington, DC, pp. 61–88.

Davis T.A.W. and Richards P.W. 1934. The vegetation of Moraballi creek, British Guiana. An ecological study of a limited area of tropical rain forest. II. *Journal of Ecology* 22: 106–155.

De Lima Filho D. de A., Matos F.D. de A., do Amaral I.L., Revilla J, Coêlho L. de S., Ramos J.F. and dos Santos J.L. 2001. Inventário florístico de floresta ombrófila densa de terra firme, na região do Rio Uruca – Amazonas, Brasil. *Acta Amazonica* 31: 565–579.

De Walt S.J., Bourdy G., Chavez de Michel L.R. and Quenevo C. 1999. Ethnobotany of the Tacana. Quantitative inventories of two permanent plots of northwestern Bolivia. *Economic Botany* 53: 237–260.

Duivenvoorden J.F. and Lips J.M. 1995. A Land-Ecological Study of Soils, Vegetation and Plant Diversity in Colombian Amazonia. *Tropenbos Series 12.* Tropenbos Foundation, Wageningen, The Netherlands.

Duivenvoorden J.F. and Lips J.M. 1998. Mesoscale patterns of tree species



diversity in Colombian Amazonia. In: Dallmeier F. and Comiskey J.A. (eds), Forest biodiversity in North, Central and South America, and the Caribbean. Research and Monitoring. MAB Series Vol. 21. UNESCO, Paris, pp. 535–549.

Dezzeo N., Maquirino P., Berry P.E. and Aymard G. 2000. Principales bosques en el area de San Carlos de Río Negro, Venezuela. *Scientia Guaianæ* 11: 15–36.

Ek R.C. 1997. Botanical Diversity in the Tropical Rain Forest of Guyana. Tropenbos-Guyana Programme, Georgetown, Guyana.

Fanshawe D.B. 1954. Forest types of British Guiana. *Caribbean Forester* 15: 73–111.

Ferreira L.V. 1997. Effects of the duration of flooding on species richness and floristic composition in three hectares in the Jaú National park in floodplain forest in Central Amazonia. *Biodiversity and Conservation* 6: 1353–1363.

Ferreira, L.V. and Prance, G.T. 1998a. Structure and species richness of low-diversity floodplain forest on the Rio Tapajós, Eastern Amazonia, Brazil. *Biodiversity and Conservation* 7: 585–596.

Ferreira L.V. and Prance G.T. 1998b. Species richness and floristic composition in four hectares in the Jaú National Park in upland forests in Central Amazonia. *Biodiversity and Conservation* 7: 1349–1364.

Ferreira L.V. and Rankin-de Merona J.M. 1998. Floristic composition and structure of a one-hectare forest in Central Amazonia. In: Dallmeier F. and Comiskey J.A. (eds), Forest Biodiversity in North, Central and South America, and the Caribbean. Research and Monitoring. MAB Series Vol. 21. UNESCO, Paris, pp. 649–662.

Finol Urdaneta H. undated. Silvicultura de la Mora de Guiana (*Mora gonggrijpii*). Facultad de Ciencias Forestales. Instituto de Silvicultura, Mérida, Venezuela. 73 pp.

Gazel M. 1983. Croissance des arbres et productivité des peuplement en forêt dense équatoriale de Guyane. Office National de Forêts, Cayenne, France. 129 pp.

Gentry A.H. 1988a. Tree species richness of upper Amazonian forests. *Proceedings of the National Academy of Sciences, USA* 85: 156–159.

Gentry A.H. 1988b. Changes in plant community diversity and floristic composition on environmental and geographical gradients. *Annals of the Missouri Botanical Garden* 75: 1–34.

Gentry A.H. 1990. Floristic similarities and differences between southern Central America and upper and central Amazonia. In: Gentry A.H. (ed.), *Four Neotropical Rain Forests*. Yale University Press, New Haven, Connecticut, pp. 141–157.

Gentry A.H. and Terborgh J. 1990. Composition and dynamics of the Cocha Cashu 'mature' floodplain forest. In Gentry A.H. (ed.), *Four Neotropical Rain Forests*. Yale University Press, New Haven, Connecticut, pp. 542–564.

Henkel T.W. in press. Monodominance in the ectomycorrhizal *Dicymbe corymbosa* (Caesalpinaceae) from Guyana. *Journal of Tropical Ecology*.

Johnston M. and Gillman M. 1995. Tree population studies in low-diversity forests, Guyana. I. Floristic composition and stand structure. *Biodiversity and Conservation* 4: 339–362.

Jonkers W.B.J. 1987. Vegetation Structure Logging Damage and Silviculture in a Tropical Rain Forest in Suriname. Wageningen Agricultural University, Wageningen, The Netherlands. 172 pp.

Klinge H., Adis J. and Worbes M. 1995. The vegetation of a seasonal varzea forest in the lower Solimoes River, Brazilian Amazonia. *Acta Amazonica* 25: 201–220.

Korning J. and Balslev H. 1994. Growth and mortality of trees in Amazonian tropical rain forest in Ecuador. *Journal of Vegetation Science* 4: 77–86.

Laurance W.F., Ferreira L.V., Rankin-de Merona J.M. and Hutchings R.W. 1998. Influence of plot shape on estimates of tree diversity and community composition in central Amazonia. *Biotropica* 30: 662–665.

Lleras E. 1997. Upper Rio Negro region. Brazil, Colombia, Venezuela. In: Davis S.D., Heywood V.H., Herrera-MacBryde O., Vila-Lobos J. and Hamilton A.C. (eds), *Centres of Plant Diversity. A Guide and Strategy for Their Conservation*. Vol. 3: The Americas. World-Wide Fund for Nature and World Conservation Union, Gland, Switzerland, pp. 333–337.

Londoña Vega A.C. 1993. Análisis estructural de dos bosques asociados a unidades fisiográficas contrastantes, en la región de Aracua (Amazonia Colombiana). Universidad de Colombia, Medellín, Colombia.

Maas P.J.M. 1971. Floristic observations on forest types in western Suriname I and II. *Proceedings of the Koninklijke Nederlandse Academie van Wetenschappen Ser. C* 74: 269–302.

Maciel U.N. and Lisboa P.L.B. 1989. Estudo florístico de 1 hectarea de mata de terra firme Km 15 da rodovia Presidente Médici-Costa Marques (RO-429), Rondônia. *Boletim do Museu Paraense Emílio Goeldi. Serie Botânica* 5: 25–37.

Mackensen J., Tillery-Stevens M., Klinge R. and Folster H. 2000. Site parameters, species composition, phytomass structure and element stores of a terra-firme forest in east-Amazonia, Brazil. *Plant Ecology* 151: 101–119.

Marin E. and Chaviel A. 1996. La vegetación: bosques de tierra firme. In: Rosales J. and Huber O. (eds), *Ecología de la Cuenca del Río Cauca, Venezuela I. Caracterización general*. *Scientia Guianae* 6, pp. 60–65.

Matos F.D.A. and Amaral I.L. 1999. Análise ecológica de um hectare em floresta ombrófila densa de terra-firme, Estrada da Várzea, Amazonas, Brasil. *Acta Amazonica* 29: 365–379.

Milliken W. 1998. Structure and composition of one hectare of central Amazonian terra firme forest. *Biotropica* 30: 530–537.

Milliken W. and Ratter J.A. 1998. The vegetation of the Ilha de Maraca. In: Milliken W. and Ratter J.A. (eds), *Maraca: The Biodiversity and Environment of an Amazonian Rain Forest*. John Wiley and Sons, New York, pp. 71–112.

Miranda I.S. 2000. Análise florística e estrutural da vegetação lenhosa do Rio Comemoração, Pimenta Bueno, Rondônia, Brasil. *Acta Amazonica* 30: 393–422.

Mori S.A. and Boom B.M. 1987. The forest. *Memoirs of the New York Botanical Garden* 44: 9–29.

Mori S.A., Rabelo, B.V., Tsou, C.-H. and Daly, D. 1989. Composition and

structure of an eastern Amazonian forest at Camaipa, Amapá, Brazil. *Boletim do Museu Paraense Emílio Goeldi. Serie Botânica* 5: 3–18.

Muniz F.H., Cesar O. and Monteiro R. 1994. Aspectos florísticos quantitativos e comparativos da vegetação arbórea da reserva florestal do Savecém, São Luís, Maranhão (Brasil). *Acta Amazonica* 24: 189–218.

Nascimento M.T., Proctor, J. and Villela, D.M. 1997. Forest structure, floristic composition and soils of an Amazonian monodominant forest on Maraca Island, Brazil. *Edinburgh Journal of Botany* 54: 1–38.

Nebel G., Kvist L.P., Vanclay J.K., Christensen H., Freitas L. and Ruiz J. 2001. Structure and floristic composition of flood plain forests in the Peruvian Amazon. I. Overstorey. *Forest Ecology and Management* 150: 27–57.

de Oliveira A.A. and Mori S.A. 1999. A central Amazonian terra firme forest I. High tree species richness on poor soils. *Biodiversity and Conservation* 8: 1219–1244.

Peres C.A. 1994. Primate response to phenological changes in an Amazonian terra firme forest. *Biotropica* 26: 98–112.

Phillips O.L., Hall P., Gentry A.H., Sawyer A.S. and Vasquez R. 1994. Dynamics and species richness of tropical rain forests. *Proceedings of the National Academy of Sciences, USA* 91: 2805–2809.

Pires J.M., Dobzhansky T.H. and Black G.A. 1953. An estimate of the number of species of trees in an Amazonian forest community. *Botanical Gazette* 114: 467–477.

Pires J.M. and Koury H.M. 1958. Estudo de um trecho de mata várzea próximo Belém. *Boletim Técnico de Instituto Agrônomo do Norte* 36: 3–44.

Pitman N.C.A. 2000. A large-scale inventory of two Amazonian tree communities. Ph.D. Thesis. Duke University, Durham, North Carolina.

Pitman N.C.A., Terborgh J., Silman M.R. and Nunez-V P. 1999. Tree species distributions in an upper Amazonian forest. *Ecology*: 80: 2651–2661.

Ramdass I., Singh J., Tamessar M. and Gopaul D. 1997. National Activity on the Use of Biodiversity, Guyana. UNDP/Office of the President/University of Guyana, Georgetown, Guyana.

Rankin-de Merona J.M., Hutchings R.W. and Lovejoy T.E. 1990. Tree mortality and recruitment over a five-year period in undisturbed upland rain forest of the central Amazon. In: Gentry A.H. (ed.), *Four Neotropical Rain Forests*. Yale University Press, New Haven, Connecticut, pp. 573–584.

Rankin-de Merona J.M., Prance G.T., Hutchings R.W., Da Silva M.F., Rodrigues W.A. and Uehling M.E. 1992. Preliminary results of a large-scale tree inventory of upland rain forest in the Central Amazon. *Acta Amazonica* 22: 493–534.

Rodrigues W.A. 1963. Estudo de 2,6 hectares de mata terra firme da Serra do Navio, Território do Amapá. *Boletim do Museu Paraense Emílio Goeldi Nova Série* 19: 1–44.

Sabatier D. 1990. Diversité des arbres et du peuplement forestier en Guyane. In: *Gestion de l'écosystème forestier et aménagement de l'espace régional*. SEPANGUI Cayenne, pp. 41–47.

Sabatier D. and Prevost M.F. 1987. Une forêt à Cacaoyers sauvages sur le Haut-Camopi, en Guyane Française. ORSTOM, Cayenne, France. 29 pp.

Sabatier D., Grimaldi M., Prevost M.-F., Guillaume J., Godron M., Dosso M. and Curmi P. 1997. The influence of soil cover organization on the floristic and structural heterogeneity of a Guianan rain forest. *Plant Ecology* 131: 81–108.

Salomão R.P. 1991. Uso de parcelas permanentes para estudos da vegetação da floresta tropical úmida. I. Município de Marabá, Pará. *Boletim do Museu Paraense Emílio Goeldi. Serie Botânica* 7: 543–604.

Salomão R.P. and Lisboa P.L. 1988. Análise ecológica da vegetação de uma floresta pluvial tropical de terra firme, Rondônia. *Boletim do Museu Paraense Emílio Goeldi. Serie Botânica* 4: 195–233.

Salomão R.P., Silva M.F.F. and Lisboa P.L. 1988. Inventário ecológico em floresta pluvial tropical de terra firme, Serra Norte, Carajás, Pará. *Boletim do Museu Paraense Emílio Goeldi. Serie Botânica* 4: 1–46.

Schulz J.P. 1960. *Ecological Studies on Rain Forest in Northern Surinam*. Van Eedenfonds, Amsterdam, The Netherlands.

Seidel R.S. 1995. Inventário de los árboles en tres parcelas de bosque primario en la Serranía de Marimón, Alto Beni. *Ecología en Bolivia* 25: 1–35.

Silva A.S.L., Lisboa P.L.B. and Maciel U.N. 1992. Diversidade florística e estrutura em floresta densa da bacia do Rio Juruá-Am. *Boletim do Museu Paraense Emílio Goeldi. Serie Botânica* 8: 203–253.

Silva M.F.F. and Rosa N.A. 1989. Análise do estrato arbóreo da vegetação sobre jazidas de cobre na Serra do Carajás-PA. *Boletim do Museu Paraense Emílio Goeldi. Serie Botânica* 5: 175–206.

Silva M.F.F., Rosa N.A. Oliveira J. 1987. Estudos botânicos na área do projecto Ferro Carajás. 5. Aspectos florísticos da mata do Rio Gelado, Pará. *Boletim do Museu Paraense Emílio Goeldi. Serie Botânica* 3: 1–20.

Silva M.F.F., Rosa N.A. and Salomão R.P. 1986. Estudos botânicos na área do projecto Ferro Carajás. 3. Aspectos florísticos da mata do aeroporto de Serra Norte – PA. *Boletim do Museu Paraense Emílio Goeldi. Serie Botânica* 2: 169–187.

Smith D.N. and Killeen T.J. 1998. A comparison of the structure and composition of montane and lowland tropical forest in the Serranía Pilón, Beni, Bolivia. In: Dallmeier F. and Comiskey J.A. (eds), *Forest Biodiversity in North, Central and South America, and the Caribbean*. Research and Monitoring. MAB Series Vol. 21. UNESCO, Paris, pp. 681–700.

Spichiger R., Loizeau P.-A., Latour C. and Barriera G. 1996. Tree species richness of a South-Western Amazonian forest (Jenaro Herrera, Peru, 73°40' / 4°54' S). *Candollea* 51: 559–577.

Stergios B., Comiskey J.A., Dallmeier F. and Licata A Nino M. 1998. Species diversity, spatial distribution and structural aspects of semi-deciduous lowland gallery forests in the western Llanos of Venezuela. In: Dallmeier F. and Comiskey J.A. (eds), *Forest Biodiversity in North, Central and South America, and the Caribbean*. Research and Monitoring. MAB Series Vol. 21. UNESCO, Paris, pp. 449–479.

Stern M.J. 1998. Field comparisons of two rapid vegetation assessment tech-

niques with permanent plot inventory data in Amazonian Peru. In: Dallmeier F. and Comiskey J.A. (eds), *Forest Biodiversity Research, Monitoring and Modelling. Conceptual Background and Old World Case Studies*. MAB Series Vol. 20. UNESCO, Paris, pp. 269–283.

Stevenson P.R., Castellanos M.C. and del Pilar Medina A. 1999. Elementos arboreos de los bosques de uno plano inundable en el parque nacional natural Tinigua, Colombia. *Caldasia* 21: 38–49.

Thomas R.S. 2001. Forest productivity and resource availability in lowland tropical forests in Guyana. Ph.D. Thesis, Imperial College London. Tropenbos-Guyana Series 7. Tropenbos-Guyana Programme, Georgetown, Guyana.

Uhl C. and Murphy P.G. 1981. Composition, structure, and regeneration of tierra firme forest in the Amazon basin of Venezuela. *Tropical Biology* 22: 219–237.

Urrego Giraldo L.E. 1997. Los bosques inundables del Medio Caqueta. Caracterización y sucesión Estudios en la Amazonia Colombiana XIV. Tropenbos Colombia, Bogota, Colombia.

Valencia R., Balsev H., Palacios W., Neill D., Josse C., Tirado M. and Skov F. 1998. Diversity and family composition of trees in different regions of Ecuador: a sample of 18 one-hectare plots. In: Dallmeier F. and Comiskey J.A. (eds), *Forest Biodiversity in North, Central and South America, and the Caribbean*. Research and Monitoring. MAB Series Vol. 21. UNESCO, Paris, France, pp. 569–584.

Valencia R., Balslev H. and Paz Y Mino G. 1994. High tree alpha-diversity in Amazonian Ecuador. *Biodiversity and Conservation* 3: 21–28.

van Andel T. 2001. Non-timber forest products of the North-West District of Guyana. Part I. Ph.D. Thesis, Utrecht University. Tropenbos-Guyana Series 8A. Tropenbos-Guyana Programme, Georgetown, Guyana.

Vasques-Martinez R. and Phillips O.L. 2000. Allpahuayo: floristics, structure and dynamics of a high-diversity forest in Amazonian Peru. *Annals of the Missouri Botanical Garden* 87: 499–527.

Worbes M., Klinge H., Revilla J.D. and Martius C. 1992. On the dynamics, floristic subdivision and geographical distribution of várzea forests in Central Amazonia. *Journal of Vegetation Science* 3: 553–564.

*Other used plot data from unpublished sources*

C. Baider (Brazil),

H. Castellanos (Venezuela),

D. Daly and M. Silveira (Brazil),

R. Ek and P. van der Hout (Guyana), D. Sabatier et al. (French Guiana),

W. Morawetz et al. (Venezuela).

*New unused plot data (to be used in our first web update)*

Henkel T.W. in press. Monodominance in the ectomycorrhizal *Dicymbe corymbosa* (Caesalpinaceae) from Guyana. *Journal of Tropical Ecology*.

Jardim M.A.G. and Vieira I.C.G. 2001. Composição florística e estrutura de uma floresta de várzea do estuário Amazônico, Ilha do Combu, Estado do Pará, Brasil. *Boletim do Museu Paraense Emílio Goeldi. Serie Botânica* 17: 333–354.

Wittman F., Anhof D. and Junk W.J. 2002. Tree species distribution and community structure of central Amazonian várzea forests by remote-sensing techniques. *Journal of Tropical Ecology* 18: 805–820.

J. Lindeman and J. Schulz (Suriname, unpublished data).

## References

- Burrough P.A. and McDonnell R.A. 1998. *Principles of Geographic Information Systems*. Oxford University Press, Oxford, UK.
- Bush M.B. 1994. Amazonian speciation: a necessarily complex model. *Journal of Biogeography* 21: 5–17.
- Cade B.S., Terrel J.W. and Schroeder R.L. 1999. Estimating effects of limiting factors with regression quantiles. *Ecology* 80: 311–323.
- Condit R., Foster R.B., Hubbell S.P., Sukumar R., Leigh E.G., Manokaran N. et al. 1998. Assessing forest diversity on small plots: calibration using species–individual curves from 50-ha plots. In: Dallmeier F. and Comiskey J.A. (eds), *Forest Biodiversity Research, Monitoring and Modelling. Conceptual Background and Old World Case Studies*. MAB Series Vol. 20. UNESCO, Paris, pp. 247–268.
- Condit R., Pitman N., Leigh E.G. Jr., Chave J., Terborgh J., Foster R.B. et al. 2002. Beta-Diversity in Tropical Forest Trees. *Science* 295: 666–669.
- Colwell R.K. and Lees D.C. 2000. The mid-domain effect: geometric constraints on the geography of species richness. *Trends in Ecology and Evolution* 15: 70–76.
- Clinebell R.R., Phillips O.L., Gentry A.H., Stark N. and Zuuring H. 1995. Prediction of neotropical tree and liana species richness from soil and climatic data. *Biodiversity and Conservation* 4: 56–90.
- Conservation International 1990. Workshop 90. *Biological Priorities for Conservation in Amazonia*. Conservation International, Washington, DC.
- Fisher A.A., Corbet A.S. and Williams C.B. 1943. The relation between the number of species and the number of individuals in a random sample of an animal population. *Journal of Animal Ecology* 12: 42–58.
- Gaston K.J. 2000. Global patterns in biodiversity. *Nature* 504: 220–227.
- Gentry A.H. 1988a. Tree species richness of upper Amazonian forests. *Proceedings of the National Academy of Sciences, USA* 85: 156–159.
- Gentry A.H. 1988b. Changes in plant community diversity and floristic composition on environmental and geographical gradients. *Annals of the Missouri Botanical Garden* 75: 1–34.
- Givnish T.J. 1999. On the causes of gradients in tropical tree diversity. *Journal of Ecology* 87: 193–210.
- Hall J.B. and Swaine M.D. 1976. Classification and ecology of closed-canopy forest in Ghana. *Journal of Ecology* 64: 913–951.
- Henkel T.W. in press. Monodominance in the ectomycorrhizal *Dicymbe corymbosa* (Caesalpinaceae) from Guyana. *Journal of Tropical Ecology*.
- Hille Ris-Lambers J., Clark J.S. and Beckage B. 2000. Density-dependent mortality and the latitudinal gradient in species diversity. *Nature* 417: 732–735.
- Huston M.A. 1980. Soil nutrients and tree species richness in Costa Rican forests. *Journal of Biogeography* 7: 147–157.
- Huston M.A. 1994. *Biological Diversity. The Coexistence of Species on Changing Landscapes*. Cambridge University Press, Cambridge, UK, 681 pp.
- Kalunzy S.P., Vega S.C., Cardoso T.P. and Shelly A.A. 1998. *S+ Spatial Stats. User's Manual for Windows and UNIX*. Mathsoft Inc., Seattle, Washington.
- Kikuzawa K. 1996. Geographical distribution of leaf life span and species diversity of trees simulated by a leaf-longevity model. *Vegetatio* 122: 61–67.
- Kleidon A. and Mooney H.A. 2000. A global distribution of biodiversity inferred from climatic constraints; results from a process-based modelling study. *Global Change Biology* 6: 507–523.
- Koenker R.W. and Bassett G.W. 1978. Regression quantiles. *Econometrica* 46: 33–50.

- Leigh E.G. 1995. Epilogue: research on Barro Colorado Island 1980–1994. In: Leigh E.G., Rand A.S and Windsor D.M. (eds), *The Ecology of a Tropical Rain Forest, Seasonal Rhythms and Long Term Changes*. Smithsonian Institution Press, Washington, DC, pp. 496–503.
- Mayle F.E., Burbridge R. and Killeen T.J. 2000. Millennial-scale dynamics of southern Amazonian rain forests. *Science* 290: 2291–2294.
- Myers N., Mittermeier R.A., Mittermeier C.G., da Fonseca G.A.B. and Kent J. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403: 853–858.
- de Oliveira A.A. and Mori S.A. 1999. A central Amazonian terra firme forest I. High tree species richness on poor soils. *Biodiversity and Conservation* 8: 1219–1244.
- Phillips O.L., Hall P., Gentry A.H., Sawyer A.S. and Vasquez R. 1994. Dynamics and species richness of tropical rain forests. *Proceedings of the National Academy of Sciences, USA* 2805–2809.
- Pitman N.C.A. 2000. A large-scale inventory of two Amazonian tree communities, Ph.D. Thesis, Duke University, Durham, North Carolina.
- Prance G.T. 1982. Forest refuges: evidence from woody angiosperms. In: Prance G.T. (ed.), *Biological Diversification in the Tropics*. Columbia University Press, New York, pp. 137–138.
- Rosenzweig M.L. 1995. *Species Diversity in Space and Time*. Cambridge University Press, Cambridge, UK, 436 pp.
- Sombroek W. 2001. Spatial and temporal patterns of Amazon rainfall. Consequences for the planning of agricultural occupation and protection of primary forests. *Ambio* 30: 388–396.
- Taylor L.R., Kempton R.A. and Woiwod I.P. 1976. Diversity statistics and the log-series model. *Journal of Animal Ecology* 45: 255–272.
- Terborgh J. 1973. On the notion of favorableness in plant ecology. *American Naturalist* 107: 481–501.
- Terborgh J. and Andresen E. 1998. The composition of Amazonian forests: patterns at local and regional scale. *Journal of Tropical Ecology* 14: 645–664.
- ter Steege H. and Zagt R. 2002. Density and diversity. *Nature* 417: 698–699.
- ter Steege H., Sabatier S., Castellanos H., van Andel T., Duivenvoorden J., de Oliveira A.A. et al. 2000. An analysis of Amazonian floristic composition, including those of the Guiana Shield. *Journal of Tropical Ecology* 16: 801–828.
- Vasques-Martinez R. and Phillips O.L. 2000. Allpahuayo: floristics, structure and dynamics of a high-diversity forest in Amazonian Peru. *Annals of the Missouri Botanical Garden* 87: 499–527.
- Venables W.N. and Ripley B.D. 1998. *Modern Applied Statistics with S-plus*. Springer, New York, 548 pp.

