

## Acidophilous mixed forests in the Ojcow National Park: thirty years pressure of air pollution

Anna MEDWECKA-KORNAŚ and Stefan GAWROŃSKI

### CONTENTS

1. Introduction
  2. Environmental conditions in the ONP
  3. Methods of investigations
  4. Some features of *Pino-Quercetum* and its soil conditions in the ONP
  5. Changes in structure and floristic composition in the stands of *Pino-Quercetum*
  6. Shifts in the share of major syntaxonomic groups of species in *Pino-Quercetum*
  7. Possible causes of vegetation changes in the *Pino-Quercetum* association
  8. Syntaxonomical conclusions
  9. Outlook
- Summary  
References

### 1. INTRODUCTION

Areas for which early phytosociological documentation exists, are particularly suitable for estimation of vegetation changes. This is also true for the Ojcow National Park (ONP), located in southern Poland, about 22 km NNW of Cracow (Fig. 1).

The plant communities of this area have been mapped about 30 years ago (MEDWECKA-KORNAŚ and KORNAŚ 1963). Apart from the published data also some unpublished materials from the 1950-ties have been preserved, which were now taken into consideration.

Our paper deals with the acidophilous mixed forest association *Pino-Querce-*




Fig. 1. Distribution of phytosociological records of acidophilous mixed forests *Pino-Quercetum* in the Ojcow National Park. Record numbers as in Table 3-5. Solid symbols: 1958-59 records; repeated in 1986-88, open symbols: additional records from 1986-88; g - records in strict reserves; a - *Fagus sylvatica* variant; b - *Abies alba* variant; c - *Pinus sylvestris* variant; d - forests; e - Park borders; f - roads.

tum Kozl. 1925 in which more distinct transformations occurred than in any other forest community in the study area. They resulted from the mass-dying of coniferous trees, mainly the silver fir, *Abies alba*, and the Scots pine, *Pinus sylvestris*. Short reports on these events were already published (ZABECKI 1984, MEDWECKA-KORNAS and GAWRONSKI 1990). Now we present the phytosociological tables with complete field records from the years 1958-59 and 1986-88 and a discussion based on pedological data and numerical indicator values, especially those of ELLENBERG (1974).

Nomenclature of taxa follows TUTIN et al. (1964-1980) for vascular plants and OCHYRA and SZMAJDA (1978) for mosses.

#### Acknowledgements

The authors wish to express their gratitude to Mrs. Barbara Szczepanowicz, M.Sci., who completed the analyses.

The project was supported by the grant of the Polish Academy of Sciences CPBP 04.04.

## 2. ENVIRONMENTAL CONDITIONS IN THE ONP

The Ojcow National Park includes the valley of rivulet Pradnik, deeply incised in jurassic limestones and surrounded by a gently rolling plateau. The relief of the Park is highly varied. The local flora consists of about 950 species of vascular plants, and the vegetation is composed of epilithic xerothermic grasslands, meadows, shrub thickets and various types of forests, mainly neutrophilous ones. Mixed forests of coniferous and deciduous trees (*Pino-Quercetum*) form the sole acidophilous forest association in the ONP (MEDWECKA-KORNAS and KORNAS 1963).

In 1960-70 a rapid decrease of vitality of coniferous trees became evident in the forests of the ONP, followed by their mass-dying and an accumulation of dead organic matter on the forest ground (KAMIENIECKI and SZCZESNY 1972). The main factor responsible for these processes seems to be the increase of air pollution originating in the near-by urban and industrial centres of Silesia, the coal power station in Siersza, the lead-zinc smelting works near Olkusz and the city of Cracow. The distant sources of pollution in the eastern part of Germany and in Czechoslovakia also contributed to the worsening of the situation (RYSZKOWSKI and BALAZY 1988). The pollutants were transported to the Park mainly by the dominant western winds.

The average yearly  $\text{SO}_2$  concentration in 1977-85 ranged in the Park from 55.0  $\mu\text{g}/\text{m}^3/24$  hours to 91.9  $\mu\text{g}/\text{m}^3/24$  hours (JUSZKIEWICZ and PARTYKA

1987), the permissible concentration in protected areas according to the Polish norms - being 75  $\mu\text{g}/\text{m}^3/24$  hours. The mean monthly values of  $\text{SO}_2$  content were higher in winter than in summer and reached 200% or more of the permissible concentration. The short term concentrations were sometimes very high (JUSZKIEWICZ and PARTYKA 1987). As demonstrated by the forest monitoring system, organized by the Institute of Forest Research in Warsaw, the mean atmospheric  $\text{SO}_2$  concentrations in the Ojcow National Park (estimated as 31.4  $\text{mg}/\text{m}^3/24$  hours in 1987/88) were about twice as high as in the Bialowieza National Park located in a relatively clean area of northeastern Poland. The concentrations of fluorine and  $\text{NO}_2$  (0.12 and 0.72  $\text{mg}/\text{m}^3/24$  hours, respectively) were about 10 times higher than in Bialowieza (data for 1987/88). Dust fall in some parts of the ONP exceeded now and then 100  $\text{t}/\text{km}^2/\text{year}$  (the admissible norm for protected area being 40  $\text{t}/\text{km}^2/\text{year}$ ). In 1982 when industrial activity was markedly lowered, the dust fall in the ONP was estimated as 51.3  $\text{t}/\text{km}^2/\text{year}$  (MANECKI et al. 1988). The slightly soluble in water fraction amounted to 78.1%; as major components were identified quartz and silicate glass, as minor component e.g. gypsum, and as trace components e.g. dolomite and calcite (silicate glass and gypsum being of industrial origin). The amount of heavy metal deposition (in  $\text{kg}/\text{km}^2/\text{year}$ ) was: Fe 1202, Zn 116.8, Mn 29.1, Pb 22.4, Cu 9.2, Ni 8.9. Zinc was found in the form highly soluble in water. The precipitations were not extremely acid, because of their neutralization by dust. Their pH was on an average 4.7 in winter and 5.3 in summer; the mean monthly values ranged from pH 3.9 (January 1989) to pH 5.9 (April 1988) (ZAJAC et al. in print). It is worthy of note that in non-polluted areas the pH of 5.6-6.0 in atmospheric precipitations is considered to be natural (SMITH 1981).

The degree of air pollution in the ONP was characterized also by means of bioindication (GRODZINSKA 1980, KAZMIERCZAKOWA et al. 1984) and by soil analyses in an exemplary stand of mature mixed forest (*Pino-Quercetum*) on the plateau (SWIEBODA 1980). The results of these analyses for the 4-14 cm soil horizon were (in p.p.m.): S: 80, Zn: 340, Pb: 85, Cr: 44, V: 21, Ni: 10, Co: 9, Cu: 6. Concentrations of sulphur, zinc and lead were many times higher than in a control plot, and those of zinc approached the values toxic for plants. These estimations were made around 1970, when the injuries to pines were not yet visible.

The local distribution of pollutants in the ONP is uneven because of the highly varied relief. This was confirmed by direct measurements and the "bark test" of HÄRTEL and GRILL (1972), which consists in measuring the electrolyt-

ic conductivity and sulphate concentration in water extracts. It was applied to the bark of *Pinus sylvestris* in the ONP by MEDWECKA-KORNAS et al. (1989).

### 3. METHODS OF INVESTIGATIONS

**Vegetation.** In the phytosociological investigations both thirty years ago and at present, the method of BRAUN-BLANQUET (1964) was used. The number of records available was limited by the relatively small area of the ONP (about 1500 ha) and by the scarcity of *Pino-Quercetum* stands. The sites of the former records have been located with the help of the map of their distribution 1:10'000 and the field notes from 1958-59. Only in few cases the localization of the former study plots was uncertain. Due to disturbances some areas of the repeated record were smaller than the original ones. For complementary information new records were made in forest stands not studied in 1958-59, but still containing a number of species typical of *Pino-Quercetum*. These records are placed in the last columns in the Tables 3-5. The individual syntaxonomic groups are numbered in the tables with Roman numerals, which are used also in the lists of sporadic species. The following abbreviations are used for the vegetation layers: a - tree layer, b - shrub layer, c - ground (herb and moss) layer.

**Soil.** The soil characteristic of the *Pino-Quercetum* stands was based on examination of twelve soil profiles distributed by four in each variant of this association. The pits were 70-80 cm deep and were dug in October 1989. In the soil samples the following values were measured: pH (potentiometrically), organic carbon (by the method of Turin), total nitrogen (by the method of Kjeldahl), assimilable  $K_2O$  and  $P_2O_5$  (by the method of Egner-Riehm) and  $CaO$  (with flame photometer). The granulometric composition of the soil was determined by the method of Casagrande modified by Proszynski.

### 4. SOME FEATURES OF *PINO-QUERCETUM* AND ITS SOIL CONDITIONS IN THE ONP

The delimitation and syntaxonomical position of *Pino-Quercetum* (= *Quercetum medioeuropaeum* MEDWECKA-KORNAS 1952) accepted in the present paper is that of the earlier papers on the vegetation of the ONP (MEDWECKA-KORNAS and KORNAS 1963) as well as of numerous other Polish publications (cfr. MEDWECKA-KORNAS et al. 1966). The stands of *Pino-Quercetum* cover about 37% of the Park area, more than any other plant community. They be-

long to the subassociation *P.-Q. luzuletosum* and are differentiated into three variants. The *Fagus sylvatica* variant occurs mainly in the transition belt between valley slopes and the plateau and covers about 4% of the Park area. The *Abies alba* variant and the *Pinus sylvestris* variant are developed mainly on the plateau. They occupy about 5% and 28% of the Park area, respectively. High share of Scots pine in the forests of the ONP results mainly from it being planted on abandoned fields and forest clearings, resulting from the intensive forest exploitation at the end of the 19th and the beginning of the 20th centuries, before the establishment of the National Park in 1957 (MICHALIK 1974).

The soils in *Pino-Quercetum* of the ONP developed in Quaternary loess deposits. They are deep in the *Abies* and *Pinus* variants, but relatively shallow in the *Fagus* variant. In two of the four profiles studied in the *Fagus* variant calcareous boulders and the underlying calcareous rocks were found already in the depth of 40 cm and 75 cm, respectively. The litter layer measured in most cases 2-5 cm, and the zone of litter humification ( $A_{0f}$ ) was distinct, but shallow (1-2 cm). The mineral humus horizon ( $A_1$ ), dark grey or black, measured 2-6-(8) cm. In some profiles a thin  $A_{1+3}$  horizon, with slightly lightened patches, was observed. The deeper horizon ( $A_3/B_1$ ) showed traces of leaching and formed a transgression zone to the deeper, textural and more or less compact  $B_1/C$  (or  $B_1/D$ ) horizon, usually extending down below the pit depth (GRESZTA and BITKA 1977, MEDWECKA-KORNAS 1952). The colour of soil (excepted the humic layers) was yellow, and the granulometric composition in near all samples was that of silty loam (Table 1). The results of chemical analyses are presented as mean values calculated separately for groups of four profiles in each variant of the association (Table 2). A similar range of variability was found for the C/N ratio in the  $A_1$  horizon in the *Fagus* variant (from 1:11-1:23), the *Abies* variant (1:10-1:23) and the *Pinus* variant (1:10-1:24). In half of the stands it did not exceed 1:16. The soil reaction was always acid, with the exception of the  $B_1/D$  horizon in the *Fagus sylvatica* variant, where it was directly influenced by the calcareous bed-rock. The pH values (in  $H_2O$ ) of the litter horizon ( $A_{0f}$ ) were similar to those of the humic-mineral horizon ( $A_1$ ) and ranged from 3.8 in some samples of the *Abies* and *Pinus* variants to 5.2 in the *Fagus* variant. The pH of the upper soil layers was slightly lower than that of the deeper layers. The contents of assimilable potassium and phosphorus of the soil ranged from 6.4-66.4 mg/100 g and 1.7-18.0 mg/100 g, respectively, with marked differences between individual profiles and their particular horizons. In several cases they attained their

**Table 1.** Granulometric composition of soil (mean values of four profiles in each of the three variants of *Pino-Quercetum*).

Horizon	Depth (cm)	Fraction content (%)					
		mm					
		1-0.1	0.1-0.05	0.05-0.02	0.02-0.005	0.005-0.002	<0.002
<i>Fagus sylvatica</i> variant							
A <sub>1</sub>	2-4(8)	15.7	16.7	35.0	19.3	5.0	8.3
A <sub>1+3</sub>	4-10	10.3	17.0	30.3	23.0	9.0	10.3
A <sub>3</sub> /B <sub>t</sub>	10-40	10.8	10.9	40.0	19.7	10.1	8.5
B <sub>t</sub> /C/D	40	11.7	13.2	37.0	23.2	4.5	10.2
<i>Abies alba</i> variant							
A <sub>1</sub>	2-5(8)	24.3	19.8	29.2	14.8	2.7	9.2
A <sub>1+3</sub>	5-9	7.5	9.5	38.0	23.5	12.5	9.0
A <sub>3</sub> /B <sub>t</sub>	9-40	5.7	12.2	45.8	21.0	7.8	7.5
B <sub>t</sub> /C	40-80	6.0	10.5	41.0	22.5	4.3	15.7
<i>Pinus sylvestris</i> variant							
A <sub>1</sub>	2-6	17.3	21.0	32.7	15.0	6.0	8.0
A <sub>3</sub> /B <sub>t</sub>	6-40	5.8	10.3	41.6	24.5	8.2	9.6
B <sub>t</sub> /C	40-75	5.0	9.5	37.0	25.7	8.2	14.5

minimum in upper leached soil horizons (A<sub>1+3</sub> or A<sub>3</sub>/B<sub>t</sub>). The distribution of assimilable calcium was similar. The hydrogen ions in the absorbing complex exceeded (with few exceptions of the B<sub>t</sub>/D horizons) the sum of the exchangeable cations. The degree of V-saturation was generally rather low (in the A<sub>1</sub> horizon: 14.4-42.1 in the *Fagus* variant, 12.5-13.8% in the *Abies* variant, 12.5-58.2% in the *Pinus* variant). The decrease of the V-values in the A<sub>1+3</sub> or A<sub>3</sub>/B horizons, in relation to C (or D), was a rule. As proved by these characters, the soils of the *Pino-Quercetum* association in the ONP represent the type of Haplic Lurisol (FAO-UNESCO 1988) or of "soil lessivé" (leached soil) developed in loess (KROLIKOWSKI et al. 1986, GRESZTA and BITKA 1977). According to this classification used in Poland they have to be regarded as intermediate between the subtypes of proper leached soils and slightly podzolized leached soils, in some stands being closer to the first of those subtypes, and in others to the second one. The differences between individual profiles, however, were of a minor rank, and no correlation has been discovered between them and the differentiation of the *Pino-Quercetum* vegetation.

**Table 2.** Chemical soil properties in the stands of *Pino-Quercetum* in the Ojców National Park (mean values for four profiles in each variant). Analyses of the A<sub>1+3</sub> horizon concern three profiles in the *Fagus sylvatica* variant and two profiles in the *Abies alba* variant.

Genetic horizons	Depth cm	Organic C %	Organic matter in %	Total N in %	C/N	pH in H <sub>2</sub> O	pH in KCl	Assimilable K <sub>2</sub> O mg/100 g of soil	Assimilable P <sub>2</sub> O <sub>5</sub> mg/100 g of soil	CaO mg/100 g of soil	Absorbing complex H S T miliequiv. per 100 g of soil	V in % (S/Tx100)
<i>Fagus sylvatica</i> variant												
A <sub>1</sub>	2-4(8)	5.29	9.13	0.396	15.9	4.7	3.9	11.8	3.0	197.0	17.9	25.3
A <sub>1+3</sub>	4-10	2.95	5.08	0.208	16.3	4.8	3.7	8.8	5.3	164.3	13.4	28.3
A <sub>3</sub> /B <sub>t</sub>	10-40	0.96	1.66	0.053	18.4	4.7	3.8	6.9	2.2	67.2	9.3	26.6
B <sub>t</sub> /C(D)	>40	0.61	1.06	0.041	15.6	5.6	4.5	9.5	2.9	286.3	5.3	56.3
<i>Abies alba</i> variant												
A <sub>1</sub>	2-5(8)	9.33	16.09	0.567	17.7	4.0	3.4	32.0	10.6	70.0	38.8	13.0
A <sub>1+3</sub>	5-9	1.78	3.08	0.113	15.6	3.6	3.3	7.4	3.5	13.7	12.0	24.4
A <sub>3</sub> /B <sub>t</sub>	9-40	0.41	0.71	0.036	11.6	4.3	3.9	12.8	5.8	6.6	5.9	27.7
B <sub>t</sub> /C	40-80	0.23	0.40	0.012	13.2	4.5	3.9	11.8	4.4	32.5	7.8	36.3
<i>Pinus sylvestris</i> variant												
A <sub>1</sub>	2-6	7.86	13.54	0.447	16.5	4.2	3.6	14.2	6.3	84.3	23.6	28.1
A <sub>3</sub> /B <sub>t</sub>	6-40	0.46	0.80	0.048	10.3	4.5	4.0	9.1	4.0	11.6	7.4	32.8
B <sub>t</sub> /C	40-75	0.20	0.35	traces	--	4.7	3.9	12.3	4.3	66.3	6.8	43.1





Table 3 (continued)

Successive number	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<b>Shrubs</b>														
III <i>Sorbus aucuparia</i>	b	1.1	1.1	2.2	+	1.1	+	1.1	+	1.1	+	1.1	+	V
IV <i>Corylus avellana</i>	c	1.1	1.1	2.1	+	1.1	+	1.1	+	1.1	+	1.1	+	V
V <i>Fraxinus alnus</i>	c	+	+	+	+	2.1	+	+	+	+	+	+	+	IV
IV <i>Lonicera xylosteum</i>	c	+	+	+	+	+	+	+	+	+	+	+	+	IV
IV <i>Evonymus verrucosus</i>	c	+	+	+	+	+	+	+	+	+	+	+	+	IV
IV <i>Viburnum opulus</i>	c	+	+	+	+	+	+	+	+	+	+	+	+	IV
IV <i>Daphne mezereum</i>	c	+	+	+	+	+	+	+	+	+	+	+	+	IV
IV <i>Rhamnus catharticus</i>	c	+	+	+	+	+	+	+	+	+	+	+	+	IV
IV <i>Ribes uva-crispa</i>	c	+	+	+	+	+	+	+	+	+	+	+	+	IV
V <i>Rubus idaeus</i>	c	+	+	+	+	+	+	+	+	+	+	+	+	IV
V <i>Sambucus nigra</i>	c	+	+	+	+	+	+	+	+	+	+	+	+	IV
<b>Ground layer</b>														
I <i>Ch. Pino-Quercetum</i> (8 loc.), <i>Pino-Quercetum</i>	1.1	2.2	1.1	2.2	1.2	1.2	3.3	2.2	1.1	2.2	1.2	1.1	1.2	2.2
* <i>Luzula luzuloides</i>	1.1	2.2	+	+	+	+	+	+	+	+	+	+	+	V
<i>Veronica officinalis</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Genista germanica</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	I
* <i>Hieracium vulgatum</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	I
* <i>Hieracium sabaudum</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	I
II <i>Ch. Vaccinio-Piceetalia</i>	+	1.2	+	+	+	+	+	+	+	+	+	+	+	II
<i>Orthilia secunda</i>	4.4	3.3	3.3	4.3	4.3	2.2	4.4	3.3	3.2	3.3	3.3	2.2	2.2	1.1
<i>Vaccinium myrtillus</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	V
III <i>Ch. Vaccinio-Piceetalia</i>	1.1	1.1	1.1	2.2	2.2	2.2	2.1	2.1	2.2	2.2	1.1	1.2	1.1	1.1
<i>Maianthemum bifolium</i>	1.1	1.1	+	+	+	+	+	+	+	+	+	+	+	V
<i>Solidago virgaurea</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
IV <i>Ch. Querceto-Fagetalia</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	V
<i>Rubus hirtus</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	V
<i>Melica nutans</i>	2.1	3.2	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Poa nemoralis</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	V
<i>Carex digitata</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	V
<i>Melilotus alba</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Polygonatum odoratum</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV

Table 3 (continued)

Successive number	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<i>Arenaria nemoralis</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	I
<i>Cephalanthus alba</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	I
<i>Hieracium nobilis</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	I
<i>Lathyrus riger</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	I
<i>Lathyrus rufus</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	I
<i>Viola reichenbachiana</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	I
<i>Galium odoratum</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	I
<i>Lamium galeobdolon</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	I
<i>Festuca gigantea</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	I
<b>V Others</b>														
<i>Anthriscus filix-femina</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Hieracium murorum</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Dryopteris filix-mas</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Oxalis acetosella</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Viola riviniana</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Ajuga reptans</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Convallaria majalis</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Cruciana glabra</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Lucula pilosa</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Monotropa hypopitys</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Mycelis muralis</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Festuca rubra</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Hypericum maculatum</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Valeriana ripensis</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Deschampsia cespitosa</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Galeopsis pubescens</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Senecio fuchsii</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<i>Impatiens parviflora</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
<b>Bryophytes</b>														
III <i>Dicranella heteromalla</i>	2.3	2.2	1.1	+	3.2	2.2	1.2	+	+	+	+	+	+	IV
I <i>Polypodium formosum</i>	1.2	2.2	2.2	2.3	2.2	2.2	1.2	2.2	2.2	2.2	2.2	2.2	2.2	IV
III <i>Polypodium nudum</i>	2.2	2.2	+	1.1	+	1.2	1.2	+	+	+	+	+	+	IV
IV <i>Trichomanes vulgatum</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
V <i>Brachydictyon velutinum</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
V <i>Calypogeia</i> sp.	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
V <i>Cladonia</i> sp.	+	+	+	+	+	+	+	+	+	+	+	+	+	IV
V <i>Plagiothecium denticulatum</i>	1.2	+	+	+	+	+	+	+	+	+	+	+	+	IV
Number of species in the table	37	36	46	42	48	51	44	36	34	50	37	38	31	29
Total number of species	43	41	50	55	61	61	44	37	34	61	40	43	33	33



Table 4 (continued)

[illegible]

Table 4 (continued)

[illegible]



Table 4 (continued)

Successive number	1	2	3	4	5	6	7	8	9	10	11	1'	2'	3'	4'	5'	6'	7'	8'	9'	10'	11'	12	13	14	15
<i>Gymnocarpium dryopteris</i>	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
<i>Impatiens parviflora</i>	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
<i>Senecio nemorosus</i>	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
<i>Urtica dioica</i>	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
<b>Bryophytes</b>																										
I <i>Polystichum formosum</i>	3.2	4.4	1.2	2.3	2.2	2.3	4.4	1.2	2.2	2.2	2.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
II <i>Pleurozium schreberi</i>	2.2	2.2	2.4	4.5	3.2	2.3	3.3	.	2.2	1.2	1.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
III <i>Dicranella heteromalla</i>	2.2	2.2	2.2	2.2	2.2	2.2	2.2	.	2.2	1.2	1.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
III <i>Plagiocheilum curvifolium</i>	2.2	2.2	2.2	2.2	2.2	2.2	2.2	.	2.2	1.2	1.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
III <i>Politis nutans</i>	1.1	1.2	2.2	2.2	2.2	2.2	2.2	.	2.2	1.2	1.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
IV <i>Atrichum undulatum</i>	.	.	2.2	2.2	2.2	2.2	2.2	.	2.2	1.2	1.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
V <i>Calyptrogonia</i> sp.	.	.	2.2	2.2	2.2	2.2	2.2	.	2.2	1.2	1.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
V <i>Plagiomnium affine</i>	2.2	2.2	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
V <i>Plagiomnium rostratum</i>	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Number of species in the table	46	40	36	33	41	44	27	36	40	39	41	26	28	34	40	42	40	36	30	32	34	24	35	34	33	37
Total number of species	52	45	42	39	43	48	30	41	46	39	50	31	30	37	44	44	43	38	31	32	35	24	38	36	38	40

Sporadic species

- II *Moneses uniflora* 9 (1.1), 11 (2.2), *Pyrola chlorantha* 9, *P. minor* 6, 11 (+2), 3', *Sphagnum girgensohnii* 2 (1.2), 7 (1.3).  
 III *Deschampsia flexuosa* 3 (+2), 14, *Thelypteris limbosperma* 4, *Leucobryum glaucum* 4 (+2).  
 IV *Crataegus monogyna* 4b, *Lonicera xylosteum* 9c, *Prunus avium* 12c, *Tilia cordata* 9b, 14c, *Viburnum opulus* 8c, 9c, *Asarum europaeum* 9, *Carex digitata* 13, *C. pilosa* 8 (1.1<sup>0</sup>), 14 (+2), 15, *C. sylvatica* 11 (+2), 1', 6', *Epilobium montanum* 2, *Equisetum sylvaticum* 1, 1', 13 (+2), *Galium odoratum* 1 (+2), 2', 4', *G. schultzei* 11, *Mercuzialis perennis* 9' (+2), *Paris quadrifolia* 4', 9', *Sorophularia nodosa* 8, *Vicia minor* 3 (+3), 3' (+2).  
 V *Alnus glutinosa* 7b, 7a, *Juniperus communis* 4b, 6b, *Larix decidua* subsp. *policarpa* 11a, *Pyrus communis* 4c, 5c, *Quercus rubra* 1b, 2b (1.1), 1'a, *Salix* cf. *aurita* 7b, c, *Salix* sp. 11b, *Sambucus* sp. 8c, *Anthoxanthum odoratum* 1, *Calamagrostis epigaea* 5' (+2), 15 (+2), *Calluna vulgaris* 9, *Campanula rotundifolia* 12 (+2), *Carex bryoides* 6, 7, *Cruciatia glabra* 3, 6, 12, *Danthonia decumbens* 3, *Fragaria vesca* 8, *Galeopsis* sp. 2, *Luella multiflora* 11, *Lycopodium clavatum* 2, *Lysimachia vulgaris* 1, 1' (1.1), *Monotropa hypopitys* 4, *Rubus corylifolius* 8' (1.1), *Rubus* sp. 5, 11, *Stellaria media* 3', *Thelypteris phegopteris* 1 (+2), 15, *Tussilago farfara* 10', *Valeriana tripteris* 11 (+2), *Brachythecium curium* 3, *B. velutinum* 3, 5, *Mnium* sp. 4 (+2), *Plagiocheilum denticulatum* 1' (1.1), 6', *P. lactum* 2, 13 (1.2).

Table 5. *Pino-Quercetum luzuletosum*, *Pinus sylvestris* variant.

Years of records	1958-1959															1987-1988														
	1	2	3	4	5	6	7	8	1'	2'	3'	4'	5'	6'	7'	8'	9	10	11	Presence degree in records										
Successive number	410	392	400	392	441	431	430	429	410	392	400	392	441	431	430	429	424	424	445											
Altitude (m)	NW	SWS	NW	NEN	E	SW	NE	NEN	NW	SWS	NW	NEN	E	SW	NE	NEN	NE	NWN	E											
Exposure	5	15	2	10	5	10	5	10	5	15	2	10	5	10	5	10	5	10	5											
Slope degree (°)	20	20	25	26	14	24	14	16	22	22	26	26	18	25	22	20	18	22	20											
Mean height of trees (m)	a1	70	70	65	30	65	85	70	30	40	35	40	50	55	60	70	75	60	40											
Canopy cover (%)	a2	20	70	65	30	65	85	70	30	40	35	40	50	55	60	70	75	60	40											
Undergrowth cover (%)	b	60	60	60	40	30	60	30	40	50	15	5	20	15	25	15	50	40	40											
Ground layer cover (%)	c	80	70	60	70	80	75	65	65	75	55	5	15	50	70	85	60	50	50											
Bryophytes cover (%)	d	50	20	70	60	40	25	10	10	5	2	2	2	5	5	5	2	20	20											
Area of sample plot (m <sup>2</sup> )	300	400	400	300	500	100	300	500	300	400	400	300	400	100	300	100	300	250	250											
Trees																														
V <i>Abies alba</i>	a	+	2.1	1.1	+	+	+	+	+	1.1 <sup>o</sup>	+	+	2.2	+	+	1.1 <sup>o</sup>	+	+	1.2											
	b	2.2 <sup>o</sup>	2.2	3.2	+	+	3.2	2.2	2.2 <sup>o</sup>	1.1	+	+	1.1	+	+	+	+	3.3 <sup>o</sup>	2.2											
V <i>Fagus sylvatica</i>	a1	+	+	1.1	1.1	1.1	1.1	1.1	1.1	1.1 <sup>o</sup>	+	+	+	+	+	3.2	+	+	2.1											
	b	2.2	+	+	+	+	+	+	1.2	1.2	3.3	4.3	4.4	3.3	1.1	2.2	+	+	+											
V <i>Pinus sylvestris</i>	a	4.3	4.3	4.3	4.2	4.3	2.2	2.1	2.2 <sup>o</sup>	2.2 <sup>o</sup>	2.2	3.3	3.3	3.3	+	2.2	4.3 <sup>o</sup>	3.3	3.3											
	b	+	+	+	2.1	+	+	+	+	+	+	+	+	+	+	+	+	+	+											
V <i>Quercus petraea</i>	a	1.1	+	+	+	+	3.2	2.1	3.3	3.2	+	+	+	+	+	2.1	+	+	3.3											
	b	1.1	2.1	+	+	+	+	+	+	1.1	+	+	+	+	+	+	+	+	1.1											
V <i>Betula pendula</i>	b	+	1.1	+	+	2.1	1.1	+	+	+	+	+	+	+	+	+	+	+	+											
III <i>Picea abies</i>	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+											
I <i>Populus tremula</i>	a	+	+	+	+	2.2	1.1	+	1.1	+	+	+	1.1	+	+	+	+	+	+											
	b	+	+	+	+	+	+	+	+	4 <sup>o</sup>	+	+	+	+	+	+	+	+	+											
V <i>Quercus robur</i>	a	2.1	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+											
	b	1.1	1.1	+	+	1.1	+	+	+	+	+	+	+	+	+	+	2.2	+	+											
IV <i>Acer pseudoplatanus</i>	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+											
IV <i>Carpinus betulus</i>	b	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+											
	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+											
V <i>Larix decidua</i>	b	1.1	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+											
	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+											
	a	+	+	+	+	+	1.1	3.3	+	+	+	+	+	+	+	2.2 <sup>o</sup>	3.3	+	+											



Table 5 (continued)

Successive number	1	2	3	4	5	6	7	8	1'	2'	3'	4'	5'	6'	7'	8'	9	10	11	
<i>Genista tinctoria</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Gymnocarpium dryopteris</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Hypericum perforatum</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Lycopodium clavatum</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Potentilla erecta</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Galeopsis pubescens</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Lysimachia vulgaris</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Senecio fuchsii</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Impatiens parviflora</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Urtica dioica</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<b>Bryophytes</b>																				
III <i>Pleurozium schreberi</i>	3.3	2.3	3.3	3.3	3.3	2.3	2.3	1.2	+	+	+	+	+	+	+	+	+	+	+	+
I <i>Polytrichum formosum</i>	2.3	1.2	3.3	3.3	3.3	2.2	1.2	2.2	+	+	+	+	+	+	+	+	+	+	+	+
V <i>Plagiomnium affine</i>	+	+	2.2	1.1	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
III <i>Pohlia nutans</i>	+	1.2	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
V <i>Calypogeia</i> sp.	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
V <i>Cladonia</i> sp.	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
III <i>Dicranella heteromalla</i>	+	1.2	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
V <i>Plagiothecium curvifolium</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
IV <i>Atrichum undulatum</i>	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Number of species in the table	39	46	39	38	44	50	51	39	30	33	28	17	26	28	27	18	43	33	30	
Total number of species	41	51	43	42	45	56	55	43	31	39	28	18	26	33	29	18	50	35	40	

**Sporadic species**

- II *Dryopteris dilatata* 1' (1.1), *Monotropa hypopitys* var. *hirsuta* 6 (+2), *Ptilium crista-castrensis* 4.  
 III *Calamagrostis arundinacea* 8 (+3), 2' (+2), *Deschampsia flexuosa* 7 (+2), 9, *Leucobryum glaucum* 1 (1.1).  
 IV *Lonicera xylosteum* 2b, *Ribes uva-crispa* 6c (+2), 9c, *Tilia cordata* 2b,c, 2b, *Viburnum opulus* 2c, 2b, *Carex digitata* 8 (+2), 6' (+2), *C. pilosa* 7, *Galium schultesii* 9, *Melampyrum nemorosum* 2, 2', *Moehringia trinervia* 7 (+2), 11 (+2), *Paris quadrifolia* 6, 11.  
 V *Alnus glutinosa* 1a,b, *Malus sylvestris* 4b, 9b, *Quercus rubra* 7a, 11c, *Salix aurita* 4b (1.1), *Sambucus* sp. 8c, *Anthraxanthum odoratum* 4, *Aruncus dioicus* 9, *Calamagrostis epigeios* 11 (+2), *Calamagrostis* sp. 6, *Campanula rotundifolia* 2, *Carex ovalis* 3, *Epilobium montanum* 10, *Fragaria vesca* 3, 6' (+2), *Hieracium pilosella* 5, *Luzula multiflora* 6, *Omalotheca sylvatica* 6, *Poa annua* 9 (+2), *Rubus* sp. 5, 10, *Rumex acetosella* 6, *Thelypteris phegopteris* 7, *Valeriana tripteris* 2, *Veronica chamaedrys* 3, 9, *Brachythecium velutinum* 2 (+2), *Plagiomnium rostratum* 11 (2.2), *Plagiothecium denticulatum* 4', *P. laetum* 7 (1.2), *Polytrichum commune* 3 (2.2), *P. juniperinum* 6 (1.2), *Rhytidadelphus squarrosus* 4.

The group of accompanying species ("others" in the Tables 3, 4, 5), which include plants of various ecological characters, also contributed to the expanding species. The most noticeable among them are the nitrophilous plants of forest clearings (of the *Epilobietea angustifolii* class): *Senecio fuchsii*, *S. nemorensis* and *Urtica dioica*. *Senecio fuchsii* and *S. nemorensis* grow in abundance mainly in the *Abies* variant, in stands with strongly thinned canopy. The newcomer *Impatiens parviflora* also belongs to the successful colo-

**Table 6.** Numbers of occurrences (n) and mean percentages of coverage (%) of selected species in three variants of the *Pino-Quercetum* association. Calculations based on the original phytosociological records from 1958-59 and those repeated in 1986-88. Conversion index from abundance degrees to percentage of coverage: 5 = 87.5%, 4 = 62.5%, 3 = 37.5%, 2 = 15.0%, 1 = 3.0%, + = 0.5%. Roman numerals indicate the syntaxonomic groups of species (as in Tables 3, 4, 5).

		<i>Fagus sylvatica</i> variant 10 records				<i>Abies alba</i> variant 11 records				<i>Pinus sylvestris</i> variant 8 records				
		1958-59		1986-88		1958-59		1986-88		1958-59		1986-88		
		n	%	n	%	n	%	n	%	n	%	n	%	
<b>Trees and shrubs</b>														
V	<i>Fagus sylvatica</i>	a1	10	61.5	10	75.5	2	0.1	4	0.6	2	0.8	2	4.8
		a2	-	-	-	-	1	0.1	7	11.2	1	0.1	8	28.0
V	<i>Abies alba</i>	a	6	17.0	6	7.2	11	58.0	10	32.7	6	2.5	4	2.4
		b	8	15.5	6	4.9	11	20.3	9	17.2	8	21.6	6	4.6
V	<i>Pinus sylvestris</i>	a	5	5.7	4	2.2	9	13.8	8	5.7	8	50.6	8	21.6
V	<i>Quercus petraea</i>	a	2	0.1	2	0.1	5	2.2	6	10.7	4	6.9	4	11.3
III	<i>Sorbus aucuparia</i>	b	9	2.9	7	2.8	11	6.3	8	5.0	6	5.0	6	2.8
V	<i>Sambucus nigra</i>	b	-	-	1	0.1	-	-	5	6.7	1	0.1	3	0.5
V	<i>Sambucus racemosa</i>	b	-	-	-	-	3	0.1	8	14.4	-	-	2	4.8
<b>Ground layer</b>														
I	<i>Ch. Pino-Quercion</i>													
	<i>Luzula luzuloides</i>		10	11.3	10	8.5	11	10.2	5	0.5	8	14.3	2	1.9
II	<i>Ch. Vaccinio-Piceetalia</i>													
	<i>Vaccinium myrtillus</i>		10	56.7	10	15.5	11	55.7	11	5.6	8	59.4	8	8.6
	<i>Orthilia secunda</i>		10	0.8	3	0.2	10	0.9	1	0.1	7	1.1	1	0.4
	<i>Trientalis europaea</i>		-	-	-	-	11	2.3	9	3.7	3	5.6	5	2.4
III	<i>Ch. Vaccinio-Piceetea</i>													
	<i>Maianthemum bifolium</i>		10	10.2	9	4.9	11	35.2	11	7.5	8	15.8	6	3.8
IV	<i>Ch. Querco-Fagetea</i>													
	<i>Rubus hirtus</i>		9	0.5	4	0.2	6	0.5	9	8.0	6	1.6	7	19.6
	<i>Lamiasrtrum galeobdolon</i>		2	0.4	3	0.2	1	0.1	4	2.0	1	0.1	2	4.8
V	<i>Others</i>													
	<i>Anthyrium filix-femina</i>		9	1.0	8	2.1	11	1.4	11	11.5	6	2.8	7	6.8
	<i>Senecio fuchsii</i>		2	0.1	4	0.2	2	0.1	8	15.8	1	0.1	1	0.1
	<i>Impatiens parviflora</i>		-	-	3	1.6	-	-	7	8.1	-	-	3	2.3
	<i>Bryophytes</i>													
I	<i>Polytrichum formosum</i>		10	11.5	3	0.2	11	23.5	8	1.0	8	17.6	2	0.1
III	<i>Pleurozium schreberi</i>		-	-	-	-	10	25.5	-	-	8	22.9	-	-

nizers. *Athyrium filix-femina*, *Dryopteris filix-mas*, *D. carthusiana* and *Oxalis acetosella* seem to be not sensitive or only slightly sensitive to the stress factors. On the contrary, plants of poor pasture and heathland (of the *Nardo-Callunetea* class), which formerly occurred sporadically in the *Pinus* variant of *Pino-Quercetum*, completely disappeared.

**d) Comparison of changes in three variants of *Pino-Quercetum*.** Thirty years ago three variants of *Pino-Quercetum* had (apart from the dominance of different tree species) a number of differential floristic features of their own. It is still possible to recognize these variants in the field, but the rank of differences between them became strongly reduced.

The *Fagus* variant (Table 3) was distinguished by a relatively low share of *Hieracium sabaudum*, as well as by the absence of *Hieracium laevigatum*, *Lycopodium annotinum*, *Melampyrum pratense* and *Pleurozium schreberi*. As most of these species disappeared recently from other variants of the association, this difference ceased to exist. The significant feature of the *Fagus* variant is still, however, the absence of *Trientalis europaea* and the presence of a number of species of mesic deciduous forests (of the *Carpinion* alliance) and of the xerothermic brushwoods (of the *Quercetalia pubescentis* order): *Daphne mezereum*, *Hepatica nobilis*, *Lathyrus niger*, *L. vernus*, *Melittis melissophyllum*, *Polygonatum odoratum*.

The *Abies* variant (Table 4) was distinguished from the two other variants by a more common occurrence of *Hieracium vulgatum*, *Lycopodium annotinum*, *Polygonatum verticillatum*, *Plagiothecium curvifolium*. It was the sole community in which *Sphagnum girgensohnii*, *Dryopteris dilatata* and *Mo-*

**Table 7.** *Abies alba* vitality in some stands of the *Pino-Quercetum* association in the Ojcow National Park (data for 1986).

<sup>x</sup> strict reserves, (1) felled tree

Nos of phytosociological records (Table 4)	Numbers of trees with the share of green needles in the crowns of (%)					Abundance and sociability
	0	5-30	35-60	65-80	above 80	
2 <sup>x</sup>	8	1	12	4	0	2.2
4	0	4	13	8	0	3.3
5	0	(1)	7	13	4	4.3
8	1	5	11	8	0	3.3
11 <sup>x</sup>	5	3	10	7	0	2.1
12 <sup>x</sup>	5	4	11	4	1	4.3

*neses uniflora* occurred. Part of these species have not been found presently. In the *Pinus* variant (Table 5) some plants of relatively warm and dry forest sites occurred thirty years ago (*Cruciata glabra*, *Convallaria majalis*), as well as some species of open, oligotrophic communities of the *Nardo-Callunetea* class (*Calluna vulgaris*, *Genista tinctoria*, *Lycopodium clavatum*, *Potentilla erecta*, *Danthonia decumbens*). At present *Convallaria maialis* became very rare, and the other species of this group do not grow any more in the *Pinus* variant.

**e) Dependence of mixed forest changes on local site conditions.** Evident differences in the intensity and direction of vegetation changes exist between particular stands of *Pino-Quercetum* in the ONP. They seem to depend on an uneven distribution of air pollutants in the study area, as well as on different kinds of forest management intervention. These correlations, already discussed with regard to the tree decline (MEDWECKA-KORNAS et al. 1989, MEDWECKA-KORNAS and GAWRONSKI 1990), are also reflected by the lower vegetation layers. In the *Fagus* variant (Table 3) relatively well preserved stands with *Vaccinium myrtillus* in quite good condition were found in places protected against the western winds, carrying the highest amounts of pollutants (e.g. stand no. 12 with SE exposure in the central part of the Park, as well as stands no. 11 and 13 with S and NNE exposure, respectively, in the northern part of the Park (see Table 3 and Fig. 1). In the *Abies* and *Pinus* variants most significant changes of both the shrub layer and the ground layer were noticed in stands with most severely damaged coniferous trees. In the *Abies* variant (Table 4) such a situation was recorded e.g. in the stands located on the highest point in the central part of the Park (Złota Góra, records 1 and 2), and on a local ridge with western exposure (record 11). A stand with relatively well preserved vegetation was found not far from plot 11, but lower on the slope exposed to the south (record 12). In the *Pinus* variant (Table 5) most stands are located on the plateau, outside of the strict reserves; they have been most severely disturbed by the forest management measures. Small areas with nearly typical character of the community still preserved, were found on southern and northern forest borders (stands with oak, *Quercus robur* or *Q. petraea*, and aspen, *Populus tremula*; records 9 and 10), and on flat areas above the rocks, in the transition zone to the plateau, at the eastern border of the Park (record 2'). The occurrence of the Scots pine in these places seems to be natural.



## 6. SHIFTS IN THE SHARE OF MAJOR SYNTAXONOMIC GROUPS OF SPECIES IN *PINO-QUERCETUM*

Changes in the floristic composition of *Pino-Quercetum* in the ONP may be synthetically expressed by using the taxonomic value of groups of species D (TUXEN and ELLENBERG 1973). The relevant indices have been calculated separately for the records made 30 years ago and those made at present, and for each of the three variants (Table 8). After these data a general decrease in

**Table 8.** Changes in the taxonomic values of the groups of species (Tuxen and Ellenberg 1937) in the *Pino-Quercetum* association of the Ojcow National Park.

A records from 1958-59                      g total of the given group's occurrences  
 B records repeated in 1986-88              n number of records (cfr. Tables 3-5)  
 C mean constancy of the group              t total of occurrences of all the species  
 D taxonomic value of a group of species      in the table  
 G collective participation of the given group      z number of species of the given group

$$C \text{ (in \%)} = \frac{g}{z \cdot n} \cdot 100$$

$$G \text{ (in \%)} = \frac{g}{t} \cdot 100$$

$$D \text{ (in \%)} = \frac{C \cdot G}{100}$$

For further information see PAWLOWSKI et al. 1966.

	z		g		C		G		D	
	A	B	A	B	A	B	A	B	A	B
<b><i>Fagus sylvatica</i></b> <b>variant (n=10)</b>										
<i>Pino-Quercion</i>	7	3	33	15	47.14	50.00	8.13	4.92	3.83	2.46
<i>Vaccinio-Piceetalia</i>	4	4	32	21	80.00	52.50	7.88	6.89	6.30	3.61
<i>Vaccinio-Piceetea</i>	5	5	46	36	92.00	72.00	11.33	11.80	10.42	8.50
<i>Querco-Fagetea</i>	27	24	130	104	48.15	43.33	32.02	34.10	15.42	14.77
Others	25	23	165	129	66.00	56.09	40.64	42.30	26.82	23.72
Total (t)	-	-	406	305	-	-	-	-	-	-
<b><i>Abies alba</i></b> <b>variant (n=11)</b>										
<i>Pino-Quercion</i>	5	4	48	16	87.27	36.36	11.43	4.42	9.97	1.61
<i>Vaccinio-Piceetalia</i>	7	5	52	25	67.53	45.45	12.39	6.91	5.65	3.14
<i>Vaccinio-Piceetea</i>	11	10	75	44	61.98	40.00	17.86	12.15	11.07	4.86
<i>Querco-Fagetea</i>	14	16	37	77	24.03	43.75	8.81	21.27	2.12	9.31
Others	34	34	208	200	55.61	53.48	49.52	55.25	27.54	29.55
Total (t)	-	-	420	362	-	-	-	-	-	-
<b><i>Pinus sylvestris</i></b> <b>variant (n=8)</b>										
<i>Pino-Quercion</i>	8	4	40	8	62.50	25.00	11.63	3.90	7.27	0.97
<i>Vaccinio-Piceetalia</i>	7	6	35	21	62.50	43.75	10.17	10.24	6.36	4.48
<i>Vaccinio-Piceetea</i>	8	7	39	20	60.94	35.71	11.34	9.76	6.91	3.48
<i>Querco-Fagetea</i>	16	17	44	54	34.38	39.71	12.79	26.34	4.40	10.46
Others	42	30	186	102	55.36	42.50	54.07	49.76	29.93	21.15
Total (t)	-	-	344	205	-	-	-	-	-	-

the number of species (z) during the last thirty years is evident (from 68 to 59 species in the *Fagus* variant, from 71 to 69 species in the *Abies* variant and from 81 to 64 species in the *Pinus* variant). The total sum of occurrences (t) decreased parallel. The taxonomic values of group of species (D) decreased evidently with regard to the species of the *Vaccinio-Piceetea* class, and still more dramatically for those of the *Vaccinio-Piceetalia* order and the *Pino-Quercion* alliance. On the contrary, the taxonomic values of the group of species representing the class *Querco-Fagetea* increased in both the *Abies* and *Pinus* variants, and remained unchanged in the *Fagus* variant. These figures strongly emphasize the profound recent transformations of the phytosociological character of *Pino-Quercetum* stands in the ONP.

## 7. POSSIBLE CAUSES OF VEGETATION CHANGES IN THE *PINO-QUERCETUM* ASSOCIATION

A large scale dieback of the forests is being actually observed in Europe and North America (CAPE et al. 1988, KLEIN and PERKINS 1987, MUELLER-DOMBOIS 1987, REBELE 1988, SCHLAEPFER 1989 and others).

The causes of this catastrophe are not yet determined. Relevant investigations concern the trees (e.g. LANDOLT and KELLER 1985, LOEHLE 1988, SMITH 1981, TURCSANYI 1988, ZIEGLER 1986). Studies on changes in the lower forest layers are rare; some of them, however, deal with the forest community as a whole (neutrophilous deciduous forests: FALKENGREN-GRERUP 1986, WILMANN et al. 1986; neutrophilous forests and acidophilous oak-birch forests: KUHN et al. 1987; acidophilous pine forests: KAZMIERCZAKOWA 1987).

In the case of the Ojcow National Park the noxious impact of air pollution upon coniferous trees seems to be evident. There are many premises for such explanation of the rapid decrease in vitality and the death of many individuals of locally most sensitive species: *Abies alba* and *Pinus sylvestris*.

The plants in the lower forest layers are not only exposed to the air pollution, but depend also on other changeable factors, e.g. increased content of pollutants in precipitations dropping from the canopy layer (ZAJAC and GRODZINSKA 1980), or transported with the stemflow (JOCHHEIM 1986), changes in canopy structure and composition (which influence the litter fall and the light and soil conditions) and so on. Soil investigations conducted in the ONP 20 to 30 years ago were unfortunately not detailed enough to document changes in the edaphic conditions in the stands of *Pino-Quercetum*. Data published by GRESZTA and BITKA (1977), as well as those from outside of the Park by MED-




Fig. 2. Indicator values of vascular plants after ELLENBERG 1974, calculated for three variants of the *Pino-Quercetum* association in the Ojcow National Park.  
L - light values, F - moisture values, R - reaction values, N - nitrogen values.  
A - 1958-59 figures for the original set of records, B - 1986-88 figures for records repeated in some stands. For each group of records the mean, maximum and minimum values of the particular indicator are marked as well as standard deviations and the significance levels.

WECKA-KORNAS (1952) seem to indicate that the pH of the soil at that time was the same as at present. The influence of acid rain and soil acidification which is very important in other sites (e.g. in Scandinavia: ANDERSSON 1986, DAHL 1988, FALKENGREN-GRERUP 1986) is not evident in the Ojcow National Park. Changes in soil conditions in the *Pino-Quercetum* stands of the ONP during the last thirty years can be inferred from the ecological indicator values of plants. For this purpose figures from ELLENBERG (1974) have been used (as e.g. by ROST-SIEBERT and JAHN 1988), with some supplementary data taken from ZARZYCKI (1984). For each individual species recorded in the former and/or present set of records the relevant figures of four indicator values (L, F, R, N) have been ascertained; then the mean indicator values for particular records have been calculated, and finally the mean values for complete sets of records for each variant and each of the study periods (Fig. 2). The light indicator values (L) generally decreased within the forests (save one stand of the *Abies* variant) and they do not seem to correlate with the thinning of the canopy layer by dieback of coniferous trees (as indicated in chapter 5a). This discrepancy resulted from the fact, that the shade-tolerant plants of the ground layer were effectively shaded by the increment of beeches or shrubs and tall perennial herbs and prospered well even in stands with recently thinned canopy - A<sub>1</sub>. The soil moisture indicator values (F) pointed out (with a few exceptions) to the increase of soil humidity in the stands studied. These changes were statistically significant in the *Abies* variant. The soil reaction indicator values (R) in the *Fagus* variant remained on an average unchanged. However, in the *Abies* and *Pinus* variants they changed significantly indicating (with one exception) an increase. These figures confirm the lack of acidification processes, as suggested by direct soil measurements (see above). The increase of the pH values does not sufficiently explain the retreat of acidophilous species in the *Pino-Quercetum* of the ONP, because this retreat took place also in the *Fagus* variant, where the soil reaction did not change. A very essential reason for vegetation changes has to be discerned in the accumulation of nitrogen in the forest environment. The indicator values for nitrogen (N) increased significantly in all studied stands, and consequently also in all three variants of *Pino-Quercetum*. In two cases the figures found presently were twice as high as thirty years ago. We certainly have to do with a far-reaching eutrophication of the forest habitats in the Park. It may result from the increased accumulation of dead matter on the forest ground, provided by dead trees, and/or from the increased content of nitrogen oxides in the atmosphere. The last factor is presently considered by many authors as the

main factor of disappearance of many rare (oligotrophic) species (ELLENBERG 1985) and the forest dieback in central Europe (ANDERSSON 1986, MANSFIELD 1988, WANG and SCHAAP 1988). KUHN et al. (1987) are of the opinion, that the present changes in the *Betulo-Quercetum* association in Switzerland, a vicariant community of the Polish *Pino-Quercetum*, resulted first of all from accumulation of nitrogen compounds from the atmosphere. The same is the opinion of WILMANN et al. (1986) in relation to changes in some oak-hornbeam forests (*Quercus-Carpinetum*) in Germany.

Even if eutrophication really is the main factor responsible for the changes in *Pino-Quercetum* of the ONP, other air pollutant cannot be neglected. High sulphur dioxide (SO<sub>2</sub>) concentration in the atmosphere is certainly one of the most important among them (see chapter 2). It causes the accumulation of sulphur in the needles of the conifers (e.g. of *Abies alba*, SZCZEPANOWICZ oral information) and the high sulphate concentration in the water extracts from the bark of *Pinus sylvestris* (MEDWECKA-KORNAS et al. 1989). Another possibly important factor is the impact of heavy metals, first of all of lead and zinc, found in high concentrations in the wood of the beeches (KAZMIERCZAKOWA et al. 1984), in mosses (GRODZINSKA 1980) as well as in the dust fall in the ONP (see chapter 2). Other noxious elements (see chapter 2) may also contribute to the changes (SWIEBODA 1980). The toxic effects of heavy metal pollution were confirmed in a study of acidophilous pine forests (*Vaccinium myrtilli-Pinetum*) in the vicinity of zinc and lead smelting works near Olkusz, some 25 km WNW of Ojcow, where local forest community became completely desintegrated (KAZMIERCZAKOWA 1987). The dust-fall in this area contained not only heavy metals but also alkaline components responsible for the increase of soil pH, even to the neutral values. Experimental studies on the response of the ground layer vegetation in acidophilous forest stands to the environmental pollution were conducted in *Pino-Quercetum* of the Niepolomice Forest, about 40 km SE of Ojcow. Some study plots were treated with mineral fertilizers, the others with industrial dusts. Under the influence of NPK or NPK and dolomite input the share of *Vaccinium myrtillus* was reduced, the share of *Trientalis europaea* remained without distinct changes, and that of *Oxalis acetosella* markedly increased (MITKA 1987). The response of these species was thus similar as in the case of habitat eutrophication in the ONP. In the experiments with industrial dusts *Vaccinium myrtillus* proved to be tolerant to high zinc and cadmium doses, but particularly sensitive to the dust from cement works, which effectively increased the pH of the soil. *Oxalis acetosella* showed a negative reaction to the dust containing alu-

minium, zinc and cadmium in concentrations many times higher than that measured in the soils of the ONP. On the other hand, dust from cement works and from coal power station stimulated expansion of *Oxalis acetosella* (BRANIEWSKI and CHRZANOWSKA 1988). The knowledge of symbiotic connections in forest plants may possibly help to explain the intricate patterns of their reactions towards the environmental stresses. Experiments in the Niepolomice Forest revealed that in stands treated with fertilizers mycotrophism in the ground layer plants was greatly reduced - or even eliminated and the non-mycorrhizal plants, e.g. *Urtica dioica*, became dominant (KEDZIERSKA 1990). In plots treated with alkaline industrial dusts from cement works the mycorrhizal plants proved to be more sensitive than the others (TURNAU 1988). On the other hand, species of the family *Ericaceae* are more resistant to the heavy metals (BRADLEY et al. 1982). This resistance, however, may be reduced through the site eutrophication, which eliminates the mycorrhiza.

## 8. SYNTAXONOMICAL CONCLUSIONS

Due to the disappearance of species characteristic of *Pino-Quercetum* and its higher syntaxonomic units, the acidophilous mixed forest stands in the ONP do not represent any more this plant association. The individual stands became disintegrated to various degrees and in various ways, and do not have the stable character which they possessed before the rise of air pollution and the decline of coniferous trees. Therefore a revision of their syntaxonomic position is needed. The floristic premises, however, are not sufficient for distinction of a new plant association. It seems to be more reasonable to follow the suggestions of FALINSKI (1966) and OLACZEK (1974) and to distinguish only several stages of degradation of *Pino-Quercetum* (e.g. stands slightly, moderately and heavily disturbed). Such a classification indicates the origin of actually existing forest stands and their present site conditions (which still remain in many respects the same as in the former *Pino-Quercetum*). In some situations, however, a different approach seems to be more appropriate. Stands with increased share of neutrophilous forest plants characteristic of the *Quercus-Fagetum* class and its subordinate syntaxa could already be regarded as fragmentary *Fagetalia* communities. Stands with thinned canopy and a large share of plants from the forest clearings could be classified as intermediate between the *Pino-Quercetum* and *Epilobietum angustifolii* communities. Solving all these problems is essential for repeating the vegetation mapping in the Park.

Disintegration of acidophilous forests associations is by no means limited to the Ojcow National Park. It is recorded from many other regions in Poland and abroad. In such situation the proposals concerning a revised classification of acidophilous mixed forests in Poland, with splitting the former *Pino-Quercetum* into numerous associations (MATUSZKIEWICZ 1984) seems to be little substantiated. This is also true of treating the *Fagus* variant of *Pino-Quercetum* as a separate association: *Luzulo nemorosae-Fagetum*, a proposal of MICHALIK (1987) which was not accepted in the present paper.

## 9. OUTLOOK

Preservation of the full diversity of forest communities, formerly existing in the Ojcow National Park, seems to be very difficult. This concerns the last remnants of acidophilous mixed forests, threatened by habitat eutrophication. Reduction of air pollution reaching the Park area seems to be *conditio sine qua non* of survival of the *Pino-Quercetum* association in this area. Such reduction on a general scale (as discussed e.g. by BACH 1985) forms at present the task of uttermost importance for environmental conservation.

## SUMMARY

The Ojcow National Park (ONP), located in the vicinity of the industrial centres of Silesia and Cracow city, is submitted to a very heavy impact of air pollution, which causes a mass-dying of coniferous trees and other changes in the forest. These processes are most notable in the mixed acidophilous forest (*Pino-Quercetum*), studied in the ONP about 30 years ago (MEDWECKA-KORNAS and KORNAS 1963). In the present paper data concerning changes in soil properties in this forest association are submitted (Tables 1, 2), as well as unpublished phytosociological records from the years 1958-59 repeated in 1986-88 (Tables 3-5, Fig. 1). Differences between these two data sets are discussed and their possible causes are suggested. In the tree and shrub layer of the stands of *Pino-Quercetum* the share of silver fir (*Abies alba*) and Scots pine (*Pinus sylvestris*) considerably decreased (Tables 6, 7). The resulting gaps were filled by deciduous woody species. In the field layer, the formerly dominant *Vaccinium myrtillus* and carpet mosses markedly retreated, as also did *Orthilia secunda*, *Maianthemum bifolium* and other acidophilous species. *Lycopodium annotinum* disappeared completely. The share of the neutrophilous species of deciduous forests and cleared forest areas increased, and the newcomer *Impatiens parviflora* emerged in large quantities. In all variants of the association taxonomic values of the groups of species of *Vaccinio-Piceetea* class and its subordinate syntaxonomic units dropped down dramatically, and those of the *Quercio-Fagetea* class increased considerably, especially in the *Abies alba* and *Pinus sylvestris* variants. Indicator values of vascular plants (after ELLENBERG 1974) confirm a significant increase of nitrogen content, i.e. an eutrophication of forest soils (Fig. 2), resulting from the immissions of nitrogen oxides (the NO<sub>2</sub>-concentration in the air in the Ojcow N.P. exceeds e.g. by 10 times that in the Białowieża N.P., NE Poland). Nitrogen pollution in the ONP is running parallel to other pollu-

tants: SO<sub>2</sub>, heavy metals, etc. To prevent further disintegration of the *Pino-Quercetum* association in the Ojcow area an essential reduction of the present air pollution level is necessary.

## REFERENCES

- ANDERSSON F., 1986: Acid deposition and its effects on the forests of nordic Europe. *Water, Air and Soil Pollut.* 30, 17-29.
- BACH W., 1985: Forest dieback: extent of damages and control strategies. *Experientia* 41(9), 1095-1104.
- BRADLEY R., BURT A.J. and READ D.L., 1982: The biology of mycorrhiza in the *Ericaceae*. 8. The role of mycorrhizal infection in heavy metal resistance. *New Phytol.* 91, 197-209.
- BRANIEWSKI S. and CHRZANOWSKA E., 1988: Effect of dusts from electro-filters of different industrial works on the vegetation. *Zesz.Nauk.Akad.Roln. w Krakowie* 226, Lesn. 18, 146-166.
- BRAUN-BLANQUET J., 1964: *Pflanzensoziologie. Grundzüge der Vegetationskunde.* (3rd ed.). Springer, Wien. 865 p.
- CAPE J.N., PATERSON I.S., WELLBURN A.R., WOLFENDEN J., MEHLHORN H., FREER-SMITH P.H. and FINK S., 1988: Early diagnosis of forest decline. Institute of Terrestrial Ecology, Dorchester. 68 p.
- CAPECKI Z. and TUTEJA W., 1977: Appraisal of the health status and threat of the Ojcow National Park by forest pests. (In Polish). *Sylvan* 3, 33-50.
- DAHL E., 1988: Acidification of soils in the Rondane Mountains, South Norway, due to acid precipitation. *Okoforsk Rapp.* 1, 1-53.
- ELLENBERG H., 1974: Zeigerwerte der Gefässpflanzen Mitteleuropas. *Scripta Geobot.* 9, 1-97.
- ELLENBERG H. (jun.), 1985: Veränderungen der Flora Mitteleuropas unter dem Einfluss von Düngung und Immissionen. *Schweiz.Z.Forstwesen* 136(1), 19-39.
- FALINSKI J., 1966: Une définition de la déformation de phytocénose. Un système de phases de dégénération des groupements végétaux. (In Polish with French summary). *Ekol. Pol., Ser. B* 12(1), 31-42.
- FALKENGREN-GRERUP U., 1986: Soil acidification and vegetation changes in deciduous forest in Southern Sweden. *Oecologia* 70, 339-347.
- FAO-UNESCO 1988: Soil map of the world. Revised Legend. *World Soil Resources. Rep.* 60, Rome. 118 p.
- GRESZTA J. and BITKA R., 1977: Soils. In: ZABIEROWSKI K. (ed.), *The nature of the Ojcow National Park.* (In Polish). *Stud.Nat.* B 28, 81-89.
- GRODZINSKA K., 1980: Heavy metals pollution of Polish National Parks. (In Polish with English summary). *Ochr.Przr.* 43, 9-27.
- HARTEL O. and GRILL D., 1972: Die Leitfähigkeit von Fichtenborken-Extrakten als empfindlicher Indikator für Luftverunreinigungen. *Eur.J.Forest.Pathol.* 2(4), 205-215.
- JOCHHEIM H., 1986: Einfluss des Stammablaufwassers auf den Pflanzenbewuchs und den chemischen Bodenzustand im Oberboden von Altbuchenbeständen. *Verh.Ges.Ökol.* 14, 229-308.
- JUSZKIEWICZ J. and PARTYKA J., 1987: Atmospheric air pollution in the Ojcow National Park. (In Polish with English summary). *Parki Nar.Rezerwaty Przr.*, 1987, 41-49.
- KAMIENIECKI F. and SZCZESNY T., 1972: Information about the state of endangered forest stands and vegetation of the Ojcow National Park by industrial air pollution. (In Polish). *Oddziaływanie Przemysłu na Lasy* 5, 15-21.
- KAZMIERCZAKOWA R., 1987: Degradation of pine forest *Vaccinio myrtilli-Pinetum* vegeta-



- tion under influence of zinc and lead smelter. Stud.Nat. A 31, 29-80.
- KAZMIERCZAKOWA R., GRODZINSKA K. and BEDNARZ Z., 1984: Content of heavy metals in xylem of 100-years old beech (*Fagus silvatica* L.) in Southern Poland. Bull.Acad.Polon.Sci., Ser. Sci.Biol. 32, 9-10.
- KEDZIERSKA A., 1990: The influence of mineral fertilization upon the mycorrhiza in the *Pino-Quercetum* association ground layer (Niepolomice Forest, Southern Poland). (In Polish). Disertatio for master degree, Institute of Botany, PAN, Krakow. 37 p. (Poly-copy).
- KLEIN R.M. and PERKINS T.D., 1987: Cascades of causes and effects of forest decline. Ambio 16(2-3), 86-93.
- KROLIKOWSKI L., KOWALINSKI S. and TRZCINSKI W., 1986: The album of Polish soils. (In Polish and English). PWN, Warszawa. 166 p.
- KUHN N., AMIET R. and HUFESCHMID N., 1987: Veränderungen in der Waldvegetation der Schweiz infolge Nährstoffanreicherungen aus der Atmosphäre. Allg.Forst-Jagd-Zeitung 158(5/6), 77-84.
- LANDOLT W. and KELLER Th. 1985: Uptake and effects of air pollutants on woody plants. Experientia 41(3), 301-310.
- LOEHLE C., 1988: Forest decline: endogenous dynamics, tree defenses, and the elimination of spurious correlation. Vegetatio 77(1-3), 65-78.
- MANECKI A., SCHEJBAL-CHWASTEK M. and TARKOWSKI J., 1988: Mineralogical and chemical characteristics of dust air pollutants from areas affected by short- and long-range industrial emissions. Prace Mineralogiczne 78, 27-45.
- MANSFIELD T., 1988: Airborne pollution: how does it kill trees? Plants Today. Jan.-Feb. 1988, 12-15.
- MATUSZKIEWICZ W., 1984: Die Karte der potentiellen natürlichen Vegetation von Polen. Braun-Blanquetia 1, 1-99.
- MEDWECKA-KORNAS A., 1952: Les associations forestières du Jura Cracovien. (In Polish with French summary). Ochr.Przyr. 20, 133-236.
- MEDWECKA-KORNAS A. and GAWRONSKI S., 1990: The dieback of fir *Abies alba* Mill. and changes in the association *Pino-Quercetum* in the Ojcow National Park (Southern Poland). Vegetatio 87, 175-186.
- MEDWECKA-KORNAS A. and KORNAS J., 1963: Vegetation map of the Ojcow National Park. (In Polish with English summary). Ochr.Przyr. 28, 16-87.
- MEDWECKA-KORNAS A., KORNAS J. and PAWLOWSKI B., 1966: Survey of the most important plant associations in Poland. In: SZAFER W.(ed.). The vegetation of Poland. Pergamon Press - PWN, Oxford-Warszawa. 294-510.
- MEDWECKA-KORNAS A., KOZLOWSKA H., GAWRONSKI S. and MATYSIAK E., 1989: Pine bark extracts as indicators of air pollution in the Ojcow National Park. (In Polish with English summary). Fragm.Florist.Geobot. 34(3/4), 425-444.
- MICHALIK S., 1974: The changes induced by man in the vegetation of the Ojcow National Park. (In Polish with English summary). Ochr.Przyr. 39, 65-154.
- MICHALIK S., 1987: Plant communities within permanent study plot "Chelmowa Gora" in the Ojcow National Park. (In Polish with English summary). Parki Nar.Rezerwaty Przyr. 1987, 79-88.
- MITKA J., 1987: The effect of mineral fertilization on quantitative ratios in the herb layer vegetation in the mixed oak-pine forest (*Pino-Quercetum*) in the Niepolomice Forest area. Acta Agrar.Silvestria, ser. Silvestris 26, 75-85.
- MUELLER-DOMBOIS D., 1987: Natural dieback in forests. BioScience, Sep. 1987, 575-583.
- OCHYRA R. and SZMAJDA P., 1978: Annotated list of Polish mosses. Fragm.Florist.Geobot. 24(1), 93-145.
- OLACZEK R., 1974: Trends of forest phytocenoses degeneration and methods of their investigations. (In Polish with English summary). Phytocoenosis 3(3-4), 179-190.

- REBELE F., 1988: Waldschäden in Mitteleuropa. Exkursionsbericht des 14. Intern.Bot. Kongr. Allg.Forstl.Z. 51-53, 1412-1417.
- ROST-SIEBERT K. and JAHN G., 1988: Veränderungen der Waldbodenvegetation während der letzten Jahrzehnte - Eignung zur Bioindikation von Immissionswirkungen? Forst und Holz 43(4), 75-81.
- RYSZKOWSKI L. and BALAZY S., 1988: Environmental hazards in Poland and strategy for conservation of living natural resources. Intecol Bull. 16, 17-28.
- SCHLAEPFER R., 1988: Dépérissement des forêts: une analyse des connaissances fournies par la recherche. Inst.Fed.Rech.Forest. Birmensdorf, Rapp. 306, 1-26.
- SMITH W.H., 1981: Air pollution and forests. Springer, Berlin. 379 p.
- SWIEBODA M., 1980: The impact of industrial air pollution on the Scots pine, *Pinus sylvestris* L., in the area between the "Boleslaw" lead and zinc smelting works and the Ojcow National Park. (In Polish with English summary). Ochr.Przyr. 43, 329-361.
- TUEXEN R. and ELLENBERG H., 1973: Der systematische und ökologische Gruppenwert. Mitt.Florist.-Soziol.Arbeitsgem.Niedersachsen 3, 171-184.
- TURCSANYI G., 1988: Adaptability of trees to the changing environment (a theoretical approach to the possible causes of forest decay). Acta Bot.Acad.Hung. 34(1-2), 65-75.
- TURNAU K., 1988 The influence of industrial dust on the mycoflora of *Pino-Quercetum* forest near Cracow. II. Effects of industrial dust on the mycorrhizal status of herb-layer plants. Zesz.Nauk.Akad.Roln. w Krakowie 226, Lesn. 18, 130-134.
- TUTIN T.C. et al. (eds.), 1964-1980: Flora Europaea vol. 1-5, Cambridge Univ.Press, Cambridge. 1: 464 p., 2: 454 p., 3: 370 p., 4: 505 p., 5: 452 p.
- WANG D. and SCHAAP W. 1988: Air pollution impacts on plants: current research challenges. ISI Atlas Sci., Anim.Plant Sci. 1(1), 33-39.
- WILMANN O., BOGENRIEDER A. and MUELLER W.H. 1986: Der Nachweis spontaner, teils autogener, teils immissionsbedingter Änderungen von Eichen-Hainbuchenwäldern - eine Fallstudie in Kaiserstuhl/Baden. Natur und Landschaft 61(11), 415-422.
- ZAJAC K.P., BIK A. and ZAWODNY Z., in print: Pollution of precipitation on the area of the Ojcow National Park. (In Polish). Ochr.Powietrza 5
- ZAJAC K.P. and GRODZINSKA K., 1980: Contamination and atmospheric precipitation. In: GRODZINSKA K. (ed.), Acidification of forest environment (Niepolomice Forest) caused by SO<sub>2</sub> emissions from steel mills. Final report of investigations. Inst.Bot.Polish Acad. Sci., Cracow, 22-56.
- ZABECKI W., 1984: The extent of weakening and dying of the *Abies alba* stands in the Ojcow National Park as affected by industrial emissions. (In Polish with English summary). Zesz.Nauk.Akad.Roln. w Krakowie 184, 113-123.
- ZIEGLER H., 1986: Pflanzenphysiologische Aspekte der Waldschäden. Rheinisch-Westfälische Akad.Wiss.Vorträge N 347, 7-40.
- ZARZYCKI K., 1984: Indicator values of vascular plants in Poland. (In Polish). Inst.Bot. Polish Acad.Sci., Cracow. 45 p.

Address of authors: Prof. Dr. Anna MEDWECKA-KORNAS  
M.Sci. Stefan GAWRONSKI  
Institute of Botany  
Jagiellonian University  
Lubicz 46  
31-512 Krakow, Poland