



Research centre
for toxic compounds
in the environment

Introduction to Ecotoxicology

linking fundamental science
with environmental risk assessment and management

Ludek Blaha + ecotox colleagues



esf european
social fund in the
czech republic



EUROPEAN UNION



MINISTRY OF EDUCATION,
YOUTH AND SPORTS



OP Education
for Competitiveness




INVESTMENTS IN EDUCATION DEVELOPMENT

Global anthropogenic threats ?

A safe operating space for humanity & the nine planetary boundaries

Rockstrom et al. 2009
(*Ecology and Society* 14(2): 32; *Nature* 461, 472-475)



Research centre
for toxic compounds
in the environment

cetocoen



EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE



OP Research and
Development for Innovation

Environmental pollution

Any examples ???

**CHEMICAL
ENTERS THE
ENVIRONMENT**



**LEVELS, FATE,
PROCESSES**





**Bioavailable
fraction**



"EXPOSURE"

acute

chronic



**CHEMICAL
ENTERS THE
ORGANISM**

biomonitoring




Toxicokinetics

*biotransformation
bioactivation*

excretion / sequestration

Target site

"EFFECT"



Research centre
for toxic compounds
in the environment

ceto

DEVELOPMENT
INVESTING IN YOUR FUTURE


Assessment of chemical hazards

...to...

Humans
(**TOXICOLOGY**)



Other organisms
(**ECO**toxicology)



Chemicals in the environment

*Do you believe that **chemicals in products** sold to consumers have been proven **safe**?*


Think again

Most chemicals in modern use have simply not been tested for their impacts on human, even very basic effects.

... what about the effects in nature, then ?



Chemicals in the environment



- **Rats exposed in the womb to a single low dose of a widespread brominated flame retardant become hyperactive and have decreased sperm counts...**
- *Experiments with dioxin and similar compounds provide support for the assumption that cancer risks mediated by the aryl hydrocarbon receptor are additive. Previously untested for cancer, this assumption underpins a standard way of estimating exposure risks to these compounds. The results reinforce the need to focus health standards on mixtures rather than single compounds.*
- *At exposure levels within the range experienced by the general public, the phthalate **DBP** reduces expression of genes necessary for testosterone synthesis in fetal rats...*
- **Eutrophication of frog ponds is linked to epidemics of frog deformities, because it creates conditions that lead to higher rates of parasitic infections of tadpoles.** *The parasitic infections in turn disrupt normal development of the tadpoles' limb buds during metamorphosis.*

Published online: 21 October 2005; | doi:10.1038/news051017-16

Pollution makes for more girls

The stress of dirty air skews sex ratios in Sao Paulo.

[Erika Check](#)

Toxic fumes favour the fairer sex, a group of researchers in Brazil has found.

Jorge Hallak and his team at the University of Sao Paulo turned up the surprising result by studying babies born in their city. They divided the metropolis of 17 million people into areas of low, medium and high air pollution, using test results from air-quality monitoring stations. They then studied birth registries of children born from 2001 to 2003.

The team found that 48.3% of babies were female in the least polluted areas, but 49.3% were female in the dirtiest parts of town. After measuring the ratio of boys to girls born in all the areas, they calculated that 1,180 more babies would have been boys in the polluted areas if they had the same sex ratios as the cleaner areas. The team reported their findings on 17 October at the American



Babies born in highly polluted areas are more likely to be girls.

© Alamy

Major anthropogenic threats – example: waters

Direct



Indirect



Major impacts

- **Loss of biodiversity**



Changes in biodiversity

NATURE (2012) 482: 20



ATTACK OF THE BLOBS

Blooms of giant Nomura's jellyfish (*Nemopilema nomurai*) have troubled Japanese fishing crews.

increase in the global population of jellyfish — a catch-all term that covers some 2,000 species of true cnidarian jellyfish, ctenophores (or comb jellies) and other floating creatures called tunicates. But many marine biologists are now questioning the idea that jellyfish have started to overrun the oceans.

This week, a group of researchers published preliminary results from what will be the most comprehensive review of jellyfish population data¹. They say that there is not yet enough evi-

NYOUDU/NEWS.COM

Major impacts

Loss of biodiversity



• Impairment of ecosystem services

– Unbalanced water cycles

- Water scarcity
- Draughts/floods

– Impaired water quality

- Drinking waters
- Bathing waters
- Toxicants in food chain


– Shrinking of food supplies

- Direct → lowering fish amounts
- Indirect → crop yield



Impacts on fish → decreased crop yields

NATURE (2005) 437: 880



Research centre
for toxic compounds
in the environment

cetocoen



EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE



OP Research and
Development for Innovation

Impacts on biota → global effects

Mixing oceans

→ cooling the atmosphere

[Nature 447, p.522, May 31, 2007]



Marine life supplies up to 50% of the mechanical energy required worldwide to mix waters from the surface to deeper cool layers

[Dewar, Marine Res 64:541 (2006)]

[Katija a Dabiri, Nature 460:624 (2009)]



Research centre
for toxic compounds
in the environment

cetocoen



EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE



OP Research and
Development for Innovation

Ecotoxicology: ecological hierarchy







Figure 3.1 Biological levels of organization. The dimensions of time and space are less important for the investigation up to the levels of populations and biocoenoses.

From molecules to ecosystem

... and backwards

1962



© Patuxent Wildlife Refuge, MA, USA

"DDT is good for me-e-e!"

The great expectations held for DDT have been realized. During 1946, exhaustive scientific tests have shown that, when properly used, DDT kills a host of destructive insect pests, and is a benefactor of all humanity.

Pennsalt produces DDT and its products in all standard forms and is now one of the country's largest producers of this amazing insecticide. Today, everyone can enjoy added comfort, health and safety through the insect-killing powers of Pennsalt DDT products . . . and DDT is only one of Pennsalt's many chemical products which benefit industry, farm and home.

GOOD FOR STEERS—Beef grows meatier nowadays . . . for it's a scientific fact that—compared to untreated cattle—beef steers gain up to 50 pounds extra when protected from horn flies and many other pests with DDT insecticides.

GOOD FOR THE HOME—helps to make healthier, more comfortable homes . . . protects your family from dangerous insect pests. Use Knox-Out DDT Powders and Sprays as directed . . . then watch the bugs "bite the dust"!

GOOD FOR DAIRIES—Up to 20% more milk . . . more butter . . . more cheese . . . tests prove greater milk production when dairy cows are protected from the annoyance of many insects with DDT insecticides like Knox-Out Stock and Barn Spray.

GOOD FOR FRUITS—Bigger apples, juicier fruits that are free from unsightly worms . . . all benefits resulting from DDT dusts and sprays.

GOOD FOR ROW CROPS—25 more barrels of potatoes per acre . . . actual DDT tests have shown crop increases like that! DDT dusts and sprays help truck farmers pass these gains along to you.

KNOX FOR INDUSTRY—Food processing plants, laundries, dry cleaning plants, hotels . . . dozens of industries gain effective bug control, more pleasant work conditions with Pennsalt DDT products.


PENN SALT
CHEMICALS
87 Years' Service to Industry • Farm • Home
PENNSYLVANIA SALT MANUFACTURING COMPANY
WIDENER BUILDING, PHILADELPHIA 7, PA.

Bitman et al. *Science* 1970, 168(3931): 594



Biochemistry


bird carbonate dehydratase



In vivo: shell thinning



In situ: bioaccumulation
-> bird population decline



ECOTOXICOLOGY by definition

- **Aim:** to maintain the natural structure and function of ecosystems
- **Definitions:**
 - ecotoxicology is concerned with the **toxic effects** of chemical and physical agents on living organisms, especially on populations and communities within defined ecosystems; **it includes the transfer pathways** and their interactions with the environment
 - science of contaminants in the biosphere and their effect on constituents of the biosphere, including humans' (Newman & Unger, 2002)
 - science that provides critical information on effects of toxic compounds on living organisms which SERVE various practical aims (environmental protection)

Ecotoxic effects





Figure 1 The effective concentration of a pollutant in an organism (e.g. fish, daphnia, algae) or at the target site inside the organism is the link between the environmental fate of a pollutant and its toxic effect.

Escher, B. I., Behra, R., Eggen, R. I. L., Fent, K. (1997), "Molecular mechanisms in ecotoxicology: an interplay between environmental chemistry and biology", *Chimia*, **51**, 915-921.

1) From molecules to individuals


MECHANISMS OF TOXICITY



2) From molecules to individuals

ADVERSE OUTCOME PATHWAYS

Mechanistic effect models for ecotoxicology




→ Arrows indicate a causal relationship

See also: Ashauer & Escher *JEM* (2010), Rubach *et al. IEAM* (2011), Jager *et al. ES&T* (2011), Ashauer *et al. ET&C* (2011)

AOP Example: ethinylestradiol


Ethinylestradiol



**Binds to
ESTROGEN
RECEPTOR**

Target genes

- Proliferation/Apoptosis (sexual organs)
- Synthesis of egg yolk (fish, amphibia)





Effects

- Females: reproduction regulation
- Males: feminization
(+ e.g. cancer promotion, development, immunomodulation)

Kidd, K.A. et al. 2007. **Collapse of a fish population** following exposure to **a synthetic estrogen**. *Proceedings of the National Academy of Sciences* 104(21):8897-8901




5 ng/L (!)
7 years



Controls


+ Ethinylestradiol



Effects at different levels

- **Molecular**

- Nonspecific effects
 - Hydrophobic interactions with phospholipid membranes (baseline = narcotic toxicity)
 - Direct reactivity: electrophilic compounds → nucleophilic organism (e.g. oxidation of PROTEINS, lipids (membranes), DNA ...)
- Specific effects
 - Activation of ER, AR and other „nuclear receptors“
 - Inhibition of enzymes (e.g. CN- inhibits hemes in mitochondria/hemoglobin)
 - Neurotoxicity in nontarget organisms (e.g. Insecticides)



Effects at different levels

- **Cellular**

- Effects on structure
- Effects on metabolism (maintenance)
- Effects on regulation


→ Changes in functions (e.g. Ethinylestradiol)

→ Repair, survival, growth

→ Death (apoptosis or necrosis)

→ Proliferation

→ Differentiation



Effects at different levels

- **Organism**

- Effects on structure
- Effects on metabolism (maintenance)
- Effects on regulation


→ Changes in functions (e.g. Ethinylestradiol)

→ Repair, survival, growth

→ Death

→ Proliferation = **Reproduction**

→ Differentiation = **Evolution**



Effects at different levels

- **Population**

(... all the organisms that both belong to the same group or species (i.e. can sexually reproduce) and live in the same time within the same geographical area)

- Effects on structure
 - elderly vs. young, males vs. females
- Effects on maintenance & growth
 - Natality, mortality, reproduction fitness



Effects at different levels

- **Community & Ecosystem**

(... a group of interacting living organisms sharing a populated environment)

- Effects on structure
 - Loss of species, loss of biodiversity
- Effects on functioning
 - (including „ecosystem functions“)




Figure #31: Simplified Food Web (Source Down) similar to warm water lower end of river before entry into Mississippi River System or impoundment. The Flathead acts as a super predator when present as large specimens, and many predators such as walleyes and Gars compete for minnows and shad. Channel Catfish also appear and prey upon mussels and other invertebrates.

(Eco)toxicology – science of „doses“

Paracelsus (1493 - 1541)




*‘What is there which
is not a poison?’*

„Cause-effect paradigm“

- *All things are poison and nothing without poison.*
- *Solely **the dose determines** that a thing is not a **poison**.*

ECOTOXICOLOGY – a synthetic science



Research centre
for toxic compounds
in the environment

Ecology

cetocoen



EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE




OP Research and
Development for Innovation

Ecotoxicology – ultimate goal ?

To identify (or predict)
safe vs hazardous
levels



Ecotoxicology: problems and approaches




Testing ecotoxicity – basics

Bioassays

- single / multiple species
- acute / chronic effects
- standardized (practical)
vs. experimental (research)

Simulation of the ecosystem

- major **trophic levels**
 - producers
 - consumers
 - decomposers





centre
compounds


In the environment




Ecotoxicology methods 1) - standardized assays



Laboratory ecotoxicology – data and results



Threshold:
**No Observed Effect
Concentration (NOEC)**



Ecotoxicology – methods 2: Micro & Mesocosms

Expensive & time consuming (e.g. *Pesticide testing*)
Variable results (natural variability ...)
Higher ecological relevancy




Fig. 5.2 Components of a standardized aquatic microcosm.

Notes on practical testing

- Testing chemicals
 - Traditional / bioassays developed to assess **individual chemicals**
 - Advantage: Standardized approaches
 - Disadvantage: Limited ecological relevance
 - often **acute** tests only
 - „too standardized...“ (? Less representative ?)
 - does not assess/consider bioavailability
 - no consideration of mixture effects
 - no consideration of specific modes of action
 - no consideration of ecological situation
- Example: Acute (96h) fish toxicity assay with ethanol
 - No deaths (but fish are passive – slow swimming) → OK ?
 - Real life: easy prey → population decline


Notes on practical testing

- Testing toxicity of natural **contaminated matrices**
 - Rather new in ecotoxicology – many open challenges
 - Whole effluent toxicity testing (WET)
 - Contact soil toxicity assays
 - More complex and more complicated
 - „cause-effects“ often not clear
 - Natural variability in matrices
 - Algal tests - nutrients (Nitrogen, Phosphorus) >> Toxic compounds

Ecotoxicology in current practice

- Most legislations on chemicals) (e.g. REACH, Pharmaceuticals, Pesticides) have very simple (basic) requirements
 - EC50 from acute toxicity
 - Of 3 basic assays
 - Algae
 - Daphnia
 - Fish

Ecotox database:
www.epa.gov/ecotox




Ecotoxicology in current practice

- How to extrapolate 3 (or few more) EC50 values to get legally binding safe concentration, which is protecting virtually all organisms?

PNEC (Predicted No Effect Concentration)

EQS (Environmental Quality Standard)



Extrapolation approaches

Ecotoxicological data

Assessment / Extrapolation factors

Data	Assessment factor
L(E)C50 short-term toxicity tests	1000
NOEC for 1 long-term toxicity test	100
NOEC for additional long-term toxicity tests of 2 trophic levels	50
NOEC for additional long-term toxicity tests of 3 species of 3 trophic levels	10

PNEC


Extrapolation approaches

Ecotoxicological data

Assessment / Extrapolation factors

Data	Assessment factor
L(E)C50 short-term toxicity tests	1000
NOEC for 1 long-term toxicity test	100
NOEC for additional long-term toxicity tests of 2 trophic levels	50
NOEC for additional long-term toxicity tests of 3 species of 3 trophic levels	10


Species sensitivity distribution (SSD)



PNEC


Species Sensitivity Distribution

EC50 values for Diethylphthalate




Species Sensitivity Distribution

Frequency of EC50 values



Cummulative distribution of EC50 values




Ecotoxicology

WHAT IS IT GOOD FOR ?

SOLVING PRACTICAL PROBLEMS

Environmental policy: Limitations of sources and effects






Cause – effect → Risk assessment

Exposure

(resulting from load)




Predicted Environmental
Concentration (PEC)




Effects

(what exposures cause effects)



Laboratory (and field) studies
Ecotoxicity tests


effective concentrations
(PNEC)



EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE




Risk assessment & management



UNCERTAINTIES & challenges in ecotoxicology

... stay cautious and critical

1) Data availability



2) Bioavailability


EQC expressed as **total concentrations** do not reflect the true environmental risk

→ **Bioavailability** of chemicals depends on number of factors:

- physico-chemical
 - pH, hardness, alkalinity, DOM concentration / quality
- biological factors:
 - species, uptake route, physiological regulation

3) Data quality

Quality and relevance of the (scientific) data used for EQC derivation of metals, in most cases, POOR !



- Literature search: 156 data points
- Application of QC/QA and acceptance criteria [measured, pH and Hum (F), salinity (M)]
- ➔ 34 data points (22 %) used in risk assessment

4) „Real ecotoxicology“ needed

1) Use non-standardized organisms

- Laboratory - aquatic snails, chironomids, soil organisms ...
- Natural – sample natural organisms and test ecotoxicity immediately

2) Assess parameters important for populations

- Reproduction
- Life cycle effects (including early life stages)

3) Consider natural situations

- Addapt test conditions (temperature?, water hardness? ...)
- Simulate real exposures (e.g. peaks during pesticide spraying)

4) „Real ecotoxicology“ needed

4) Work on **models** – answer difficult questions ?

- AOPs (?)
- E.g. ecological impacts of pharmaceuticals ?


4) „Real ecotoxicology“ needed

4) Work on **models** – answer difficult questions ?

- AOPs (?)
- E.g. ecological impacts of pharmaceuticals ?

Example - antiparasitic ivermectin

- Used (for example) 2-times per season per sheep/cow
- Kills 100% parasites in sheep
- Released in dung - kills 80% larvae of dung flies
- High concentrations in dung (released 2 days post application)
- Fairly persistent in the soil (half-life 30 days)
- May be washed into adjacent streams (highly toxic to water insects)



4) „Real ecotoxicology“ needed

4) Work on **models** – answer difficult questions ?

- AOPs (?)
- E.g. ecological impacts of pharmaceuticals ?

Example - antiparasitic ivermectin


- Used (for example) 2-times per season per sheep/cow
- Kills 100% parasites in sheep
- Released in dung - kills 80% larvae of dung flies
- High concentrations in dung (released 2 days post application)
- Fairly persistent in the soil (half-life 30 days)
- May be washed into adjacent streams (highly toxic to water insects)

- **What are the indirect impacts on soil biota ?**

- Soil texture and quality ? Will plants grow on the pastures ?

- **Any impacts on bats, birds?**

- Dung flies and aquatic invertebrates serve as food



Practical example for ecotoxicologist

European strategy how to deal with chemicals



- **± 40 Directives** or Regulations concerning the evaluation and management of the dangers/risks associated with chemical substances
 - Regulation EEC 793/93 – **Existing substances**
 - Dir. 67/548/EEC – **New substances**
 - Dir. 98/8/EC – Biocides / Plant Protection Products
 - Further Directives – E.R.A. of new pharmaceuticals

Existing substances

- > 95,000,000 known chemicals
(...and counting <http://www.cas.org/>)
- 100,000 substances in EINECS (i.e. commercial use)
- 2747 HPVCs (High Production Volume Chemicals)
 - 14% minimum data-set (base-set)
 - 65% less than base-set
 - 21% no toxicity data
- Various priority lists
 - Aquatic hazard (EU Water framework directive)
 - Endocrine disruptors
 -

REACH

Registration, Evaluation and Authorisation of Chemicals

- 27-2-2001: White Paper on the Strategy for Future Chemicals Policy
- 23-10-2003: Commission's proposal REACH
- December 2008: Pre-registration mandatory (all chemicals in EU must be registered at ECHA)



European Chemicals Agency

HOME

SIEF

REACH

CONSULTATIONS

ECHA CHEM

REACH-IT

CLASSIFICATION

HELP

European Chemicals Agency (ECHA)

The Agency, located in Helsinki, Finland will manage the registration, evaluation, authorisation and restriction processes to ensure consistency across the European Union. These REACH processes are designed to provide additional information on the safe use of chemicals, and to ensure competitiveness of the European industry.

In its decision-making the Agency will take the best available scientific and technical data and socio-economic information on chemicals and technical and scientific advice. By assessing and approving testing proposals, the Agency will ensure that animal testing is reduced to a minimum.

During the first 12 months the Agency is building up its organisation and recruiting personnel to be ready to accept applications for registration.


[More](#)

European Chemicals
Agency
(<http://echa.europa.eu>)



EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE







* Substances do not have to be registered or evaluated to be placed under authorisation or restriction. They can be identified in other ways.

** Can cause cancer or mutations, or is toxic to reproduction; or is persistent, bio-accumulative and toxic, or very persistent and very bio-accumulative.

REACH





- **Major goals**

- Protection of man and the environment
- Increase competitiveness of EU chemical industry
- Increase transparency
- Avoid fragmentation of market
- Integration with international policies
- Reduction use of test animals

- **Approach**

- Industry is responsible – provides data

- **30000 existing substances**

- 0-3 year (2010): all HPVC and CMR substances (~ 3000)
- 4-6 year (2013): all 100-1000 t/y substances
- 7-11 year (2018'): all 10-100 and 1-10 t/y substances



- **Physico-chemical properties, e.g.:**
 - Vapour pressure, boiling point, Kow,...
- **Human toxicology, e.g.:**
 - Acute and chronic toxicity, skin irritation, carcinogenicity,...
- **Environment/ Ecotoxicological information, e.g.:**
 - Acute and/or chronic toxicity for aquatic organisms, biodegradation, ...

REACH: how many substances



Table 6. Estimated testing needs (% of total number of substances)

Endpoint	Minimum	Average	Maximum
6.3 Skin sensitisation	7486 (25.5)	10293 (35.1)	13728 (46.8)
6.2 Eye irritation (incl. <i>in vivo</i>)	5923 (20.1)	6910 (23.5)	8182 (27.9)
6.4.4 <i>In vivo</i> mutagenicity study	6580 (22.4)	6580 (22.4)	6580 (22.4)
7.1.2 Growth inhibition algae	2638 (9.0)	5277 (18.0)	11466 (39.1)
7.1.4 Active sludge respiration test	4616 (15.7)	4616 (15.7)	4616 (15.7)
7.1.1 Short-term <i>Daphnia</i> toxicity	2321 (7.9)	4096 (14.0)	8798 (30.0)
6.1 Skin irritation/corrosion (incl. <i>in vivo</i>)	1974 (6.7)	3949 (13.4)	5817 (19.9)
7.2.2.1 Hydrolysis	2691 (9.2)	3425 (11.7)	4518 (15.4)
6.4.1 Gene mutation study in bacteria	875 (3.0)	2916 (9.9)	6424 (21.9)
6.4.2 Cytogenicity study in mammalian cells	875 (3.0)	2916 (9.9)	6424 (21.9)
6.7.2 Development toxicity study	2408 (8.2)	2893 (9.9)	3711 (12.6)
7.2.1.1 Ready biodegradability test	1574 (5.4)	2624 (8.9)	5752 (19.6)
6.7.3 Two-generation reproduction toxicity	1665 (5.7)	2135 (7.3)	2699 (9.2)



REACH: testing



Classification categories	Test requirements in REACH			
	>1t		>10t	>100t
	New or prioritised substance			
Reproductive toxicity (a generation test)	no	no	no	no
Chronic toxicity and cancer	no	no	no	(yes)
90-day study	no	no	no	(yes)
28-day study	no	no	(yes)	yes
Acute toxicity (a second route of exposure)	no	no	yes	yes
Acute toxicity	no	yes	yes	yes
Skin allergy	no	yes	yes	yes
Skin and eye irritation	no	yes	yes	yes
Mutagenicity (in vitro)	no	yes	yes	yes
Further ecotoxicity studies (incl long term tests)	no	no	no	yes
Acute toxicity: fish	no	no	yes	yes
Acute toxicity: algae	no	yes	yes	yes
Acute toxicity: Daphnia	no	yes	yes	yes
Biotic degradation	no	yes	yes	yes



REACH: costs

	>1t/y	>10t/y	>100t/y	>1000t/y	Total
Registration costs	€ 100 mn	€ 100 mn	€ 100 mn	€ 200 mn	€ 500 million
Testing costs	€ 150 mn	€ 300 mn	€ 350 mn	€ 450 mn	€ 1250 million
Safety data sheet costs					€ 250 million
Authorisation procedures					€ 100 million
Reduced costs for new substances below 1t etc.					(benefit of € 100 million)
Total testing and registration costs					€ 2, 000 million
Agency fees (paid by chemicals sector)					€ 300 million
Total costs (including Agency fees)					€ 2, 300 million

REACH: testing costs




Table 8. Estimated testing costs for most costly endpoints (Million EURO)

Endpoint	Minimum	Average	Maximum
6.7.2 Development toxicity study	396	476	611
6.7.3 Two-generation reproduction toxicity	293	376	475
6.4.4 <i>In vivo</i> mutagenicity study	129	129	129
6.6.2 Sub-chronic toxicity	76	111	210
6.6.3 Long-term repeated dose toxicity study (incl. 6.9 Carcinogenicity study)	44	52	73
6.6.1 Short-term repeated dose toxicity study	13	49	189
6.4.2 Cytogenicity study in mammalian cells	16	52	116
6.3 Skin sensitisation	29	40	54
7.2.1.1 Ready biodegradability test	19	32	71
7.3.2 Accumulation	14	28	67
7.1.2 Growth inhibition algae	13	26	57
6.7.1 Development toxicity screening	12	26	101
7.2.2.1 Hydrolysis	16	21	28



REACH: test and cost reduction?



**MODELS,
QSAR**

Figure 6.1. Ecotoxicology is a multi-disciplinary study into the toxic effects of substances on species in complex systems [1].



All European Economic Area (EEA*) Countries

Data as of: 31/10/2012



Overall Summary

Table 1: REACH dossiers (registrations by companies from 1st June 2008 - NONS excluded)

(NONS are substances notified to Member State Competent Authorities under the previous European chemicals legislation - Directive 67/548/EEC)

The following table shows the total number of new registration numbers granted by ECHA following submission of a registration dossier.

Note that registrations resulting from notifications (NONS) made by companies under the previous chemicals legislation (Directive 67/548/EEC) are reported separately below in Table 2. The allocation of substances and dossiers across Tables 1 and 2 is described in detail in Table 3 at the end of this report.

	# Registrations	# Unique Substances
TOTAL	27 684	4 734
phase-in	26 131	4 004
non phase-in	1 553	730

Table 2: Notified substances (NONS, notified to Member State Competent Authorities under the previous European chemicals legislation - Directive 67/548/EEC)

Substances notified under Directive 67/548/EEC (NONS) prior to the introduction of REACH are considered as registered. The table below shows (a) the number of NONS which were granted a registration number by ECHA (b) the number of these claimed by the notifier and (c) the number for which an update has been submitted under REACH.

	# Registrations	# Unique Substances
(a) Number of NONS Notifications	9 962	5 292
(b) of which have been claimed	5 091	3 715
(c) of which have been updated under REACH	1 475	1 305

* *Non-phase in – nebyly předregistrovány 2008 = nesmí být vůbec použity*

* *NONS – zvláštní pravidla: apriori považovány za „registrované“*



Registered Substances by Total Tonnage Band

Total Tonnage Band: This is calculated by summing the latest year values for actual tonnages in all full registrations (i.e. not including intermediates) for a given substance and converting it to a band



Tonnage Band	# Substances
100 000 000 - 1 000 000 000 tonnes per annum	5
10 000 000 - 100 000 000 tonnes per annum	45
1 000 000 - 10 000 000 tonnes per annum	156
100 000 - 1 000 000 tonnes per annum	325
10 000 - 100 000 tonnes per annum	594
1 000 - 10 000 tonnes per annum	938
100 - 1 000 tonnes per annum	323
10 - 100 tonnes per annum	173
1 - 10 tonnes per annum	237
Intermediate Use Only	1 938
TOTAL	4 734

REACH: XI-2012



Most frequently registered Substances # Registrations

calcium dihydroxide	323
ethylene oxide	318
ethanol	310
iron	301
calcium sulphate	266
Fuels, diesel	241
calcium oxide	229
Ashes (residues), coal	217
methyloxirane	214
aluminium	180
ethylene	180
propene	176
Fuel oil, residual	175
aluminium oxide	174
Gasoline	168
Kerosine (petroleum)	162
silicon	160
buta-1,3-diene	159

REACH: XI-2012



Overview of all Countries	# Registrations	# Substances
Germany	6 789	2 708
United Kingdom	3 182	1 172
Netherlands	2 444	1 059
France	2 408	1 081
Belgium	2 101	1 069
Italy	2 020	949
Spain	1 842	854
Poland	866	370
Sweden	764	445
Finland	734	394
Ireland	628	361
Czech Republic	531	304

REACH ... Current situation: check ECHA



Search the ECHA Website

Advanced search >

- About Us
- Regulations
- Addressing Chemicals of Concern
- Information on Chemicals**
- Chemicals in our Life
- Support

ECHA > Information on Chemicals



Information on Chemicals

This is unique source of information on the chemicals manufactured and imported in Europe. It covers their hazardous properties, classification and labelling, and information on how to use them safely. This information is a valuable resource for advancing the safe use of chemicals and for the replacement of the most hazardous ones by safer alternatives.

Search for Chemicals

I have read and I accept the legal notice

Name, EC or CAS No

REACH



- > [Registered substances](#)
- > [Pre-registered substances](#)
- > [EC Inventory](#)
- > [Dossier Evaluation decisions](#)
- > [Testing Proposals Consultation](#)
- > [Substance Evaluation - CoRAP](#)
- > [Information on Candidate List substances in articles](#)

See also under the *Addressing Chemicals of Concern* section

- > [Candidate List of Substances of Very High Concern for Authorisation](#)
- > [Substances requiring Authorisation](#)
- > [Substances restricted under REACH](#)
- > [Public Activities Coordination Tool \(PACT\)](#)



REACH: implications




- Total: 2,8 to 5,6 billion €
- Industry pays
- Test costs (50-60% of total cost):
 - 86% for HH tests
 - 14% for environment tests
 - 0% for analyses
- Manpower and expertise?
 - Tests
 - Risk assessments
 - Evaluations
- Financial and time pressure:
danger for 'hazard-based' instead of 'risk-based' approach

Risks of chemicals: a balancing act


between perception, uncertainties,
science and pragmatism?

Final considerations

Ecological risks of chemicals



Risks vs. Benefits



Society a balancing act ...



Scientist

Closing remarks



- Ecotoxicology is exciting **science!**
- Interface: science and society
- Many opportunities
- Science is a hard work
 - 10% inspiration and 90% „perspiration“
- Be creative: move frontiers
- Keep the purpose in mind
- Be critical: do not accept perceptions as facts
- Speak up: you have something to say!

Introduction to ecotoxicology

Ludek Blaha

blaha@recetox.muni.cz

<http://www.recetox.cz>



INVESTMENTS IN EDUCATION DEVELOPMENT