

Multinukleární NMR spektroskopie C6800

- Jiří Pinkas, A12-224
- Materiály z přednášky v Isu
- Řešené úlohy ze spektroskopie nukleární magnetické resonance


<http://nmr.sci.muni.cz>

- Úlohy – vyřešit a odevzdat
- Prezentace (na konci semestru) 10-15 min na vybrané téma NMR
- Závěrečná písemná zkouška

NMR – Historical Perspective

- 1922 Electron spin is observed (Stern-Gerlach)
- 1926 Nuclear spin - David Dennison (H_2)
- 1938 I. I. Rabi observes NMR
in a molecular beam of H_2
- Isidor I. Rabi awarded Nobel prize in physics 1944

"for his resonance method for recording the magnetic properties of atomic nuclei"




(1898 – 1988)

NMR – Historical Perspective

- 1945 Purcell, Torrey, Pound @ Harvard solid paraffin
- 1945 Bloch, Hansen, Packard @ Stanford liquid H₂O
- Varian Bros. & Russell klystron for radars (WWII)
- 1948 Pake, van Vleck solid state NMR
- 1950 W. G. Proctor, F. C. Yu @ Stanford
 δ - chemical shift in $^{14}\text{NH}_4^{14}\text{NO}_3$
- 1950 W. C. Dickinson @ MIT
 δ - chemical shift in ^{19}F
- 1952 Commercial NMR instruments used at DuPont, Shell, Humble Oil

NMR – Historical Perspective



Edward M. Purcell (1912-1997) & Felix Bloch (1905-1983)
NP in physics 1952




"for their development of new methods for nuclear magnetic precision measurements and discoveries in connection therewith"

NMR – Historical Perspective


- 1951 Proctor , Yu - the first observed J scalar coupling $^{121}\text{Sb}-^{19}\text{F}$ in NaSbF_6
- 1951 Gutowsky, McCall, Slichter @ U. of IL - J scalar coupling $^{31}\text{P}-^{19}\text{F}$
- 1952 Hahn, Maxwell @ Berkeley - J scalar coupling
- 1955 Bloom, Shoolery spin decoupling
- 1960 Shoolery integration
- 1966 Ernst, Anderson FT NMR at Varian
- 1968 Waugh @ MIT HR, multipulse NMR in solids

NMR – Historical Perspective



- 1971 Jeener - 2D NMR
- 1971 Damadian - different NMR relaxation times of tissues and tumors
- 1972 CP, HP decoupling
- 1972 The first routine ^{13}C NMR spectrometer (before mainly ^1H , ^{19}F , and ^{31}P NMR)
- 1973 Lauterbur - MRI



MRI-Magnetic Resonance Imaging



**Paul C. Lauterbur
(1929-)**




**Sir Peter Mansfield
(1933-)**

NP in physiology and medicine 2003

NMR – Historical Perspective

- 1974/1979 R. R. Ernst 2D COSY, NOESY
- 1977 MAS
- 1981 Bax, Freeman INADEQUATE
- 1982 APT
- 1983 Freeman BB decoupling, MLEV, WALTZ
- 1990 3D and $^1\text{H}/^{15}\text{N}/^{13}\text{C}$ Triple resonance
- 1991 R. R. Ernst NP in chemistry
- 2001 The first commercial 900 MHz instrument
- 2002 K. Wüthrich NP in chemistry

NMR – Historical Perspective



Richard R. Ernst
(1933-)
NP in chemistry 1991

"for his contributions to the development of the methodology of high resolution nuclear magnetic resonance (NMR) spectroscopy"




Kurt Wüthrich
(1938-)
NP in chemistry 2002

"for his development of nuclear magnetic resonance spectroscopy for determining the three-dimensional structure of biological macromolecules in solution"



NMR – Historical Perspective




Nuclear Magnetic Resonance

- High resolution liquid state NMR spectroscopy
- Solid state NMR spectroscopy
- High-pressure NMR
- NMR in the gas phase
- NMR spectroscopy in liquid crystalline media
- Magnetic resonance imaging (MRI)

Hyperfine Interactions

- Interactions of nuclei with the electric and magnetic fields
- Interactions between a nucleus and electrons
- Transfer of chemical (electronic) information from bonds and lone pairs to a nucleus:
 - Indirect
 - Direct




Hyperfine Interactions

Indirect

- Electric field gradient (EFG) with nuclear electric quadrupole
- Induced magnetic field with nuclear magnetic moments (shielding)

Direct


- s-electrons within nuclei, polarization of bonding spins by nuclear spin (J-coupling)



Direct Interactions

ONLY s-electrons can interact with nuclei

ONLY s-electrons have non-zero electron density at a nucleus



Which quantum number determines the number of angular nodes?
Which quantum number determines the number of radial nodes?

Relationship Between Wavelength, Frequency and Energy

- Speed of light (c) is the same for all wavelengths

$$c = 2.9979 \times 10^8 \text{ m s}^{-1}$$

- Frequency (ν), the number of wavelengths per second, is inversely proportional to wavelength:


$$\nu = c/\lambda$$

- Energy of a photon is directly proportional to frequency and inversely proportional to wavelength:


$$E = h\nu = hc/\lambda$$

$$h = \text{Planck's constant} = 6.626176 \times 10^{-34} \text{ J s}$$

Electromagnetic Radiation



Method Energy Scale



Energy Scale Conversion Factors

	Hz	eV	J mol ⁻¹
Hz	1	4.136×10^{-15}	3.990×10^{-10}
eV	2.418×10^{14}	1	9.649×10^4
J mol ⁻¹	2.506×10^9	1.036×10^{-5}	1

Isotopes

**Isotopes = a set of nuclides of an element, same Z, different A
there is about 2600 nuclides (stable and radioactive)**

340 nuclides found in nature

270 stable and 70 radioactive

Monoisotopic elements:

${}^9\text{Be}$, ${}^{19}\text{F}$, ${}^{23}\text{Na}$, ${}^{27}\text{Al}$, ${}^{31}\text{P}$, ${}^{59}\text{Co}$, ${}^{127}\text{I}$, ${}^{197}\text{Au}$

Polyisotopic elements:

${}^1\text{H}$, ${}^2\text{H}$ (D), ${}^3\text{H}$ (T)

${}^{10}\text{B}$, ${}^{11}\text{B}$


Sn has the highest number of stable isotopes – 10

112, 114, 115, 116, 117, 118, 119, 120, 122, ${}^{124}\text{Sn}$

Natural Abundance, %

Isotopic Compositions of the Elements

${}^A\text{Hg}$	I	NA%
196	0	0.146
198	0	10.02
199	1/2	16.84
200	0	23.13
201	3/2	13.22
202	0	29.80
204	0	6.850



Mass number, A

I = Nuclear Spin

Natural Abundance, %

Isotopic Compositions of the Elements

^1H	99.985	^{16}O	99.759
^2H	0.015	^{17}O	0.037
		^{18}O	0.204
^{12}C	98.89		
^{13}C	1.11	^{32}S	95.00
		^{33}S	0.76
^{14}N	99.63	^{34}S	4.22
^{15}N	0.37	^{36}S	0.014

Variability in Isotopic Compositions

Natural Abundance, %

Isotope	Range	Average
^{10}B	18.927 - 20.337	19.9 (7)
^{11}B	81.073 - 79.663	80.1 (7)
^{16}O	99.7384 - 99.7756	99.757 (16)
^{17}O	0.0399 - 0.0367	0.038 (1)
^{18}O	0.2217 - 0.1877	0.205 (14)

Nuclear Spin

electron spin $s = 1/2$


proton and neutron $I = 1/2$

nuclear spin $I = z \cdot 1/2$ $z = \text{integer } 0, 1, 2, 3, \dots$

Number of protons, Z	Number of neutrons, N	I
even	even	0
odd	odd	integer
even	odd	multiples of $1/2$
odd	even	

Nuclear Spin

protons and neutrons are Fermions, obey Pauli exclusion principle



Nuclear Spin

■ **even – even: $I = 0$** ${}^4\text{He}$, ${}^{12}\text{C}$, ${}^{16}\text{O}$, ${}^{20}\text{Ne}$,
 ${}^{24}\text{Mg}$, ${}^{28}\text{Si}$, ${}^{32}\text{S}$, ${}^{36}\text{Ar}$, ${}^{40}\text{Ca}$

■ **odd – odd: $I = \text{integer}$**

ONLY ${}^2\text{H}$, ${}^6\text{Li}$, ${}^{10}\text{B}$, ${}^{14}\text{N}$, ${}^{40}\text{K}$, ${}^{50}\text{V}$, ${}^{138}\text{La}$, ${}^{176}\text{Lu}$

■ **even – odd and odd – even:**

$I = \text{multiples of } \frac{1}{2}$

${}^{13}\text{C } \frac{1}{2}$, ${}^{17}\text{O } \frac{5}{2}$, ${}^{33}\text{S } \frac{3}{2}$

Nuclear Spin

Number of protons Z	Number of neutrons N	Number of nuclides
even	even	168
odd	odd	8
odd	even	50
even	odd	57

Allowed Nuclear Multipole Moments as a function of Spin I

	$l = 0$	$l = 1$	$l = 2$	$l = 3$	$l = 4$
Spin	monopole	dipole	quadrupole	octapole	hexadecapole
$I = 0$	electric	0	0	0	0
$I = \frac{1}{2}$	electric	magnetic	0	0	0
$I = 1$	electric	magnetic	electric	0	0
$I = \frac{3}{2}$	electric	magnetic	electric	magnetic	0
$I = 2$	electric	magnetic	electric	magnetic	electric

Nuclear Magnetic Dipole Moment couples to magnetic field




H																	He
Li	Be											B	C	N	O	F	Ne
Na	Mg											Al	Si	P	S	Cl	Ar
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
Cs	Ba	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
Fr	Rd	Ac															
		Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu		
		Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr		

Spin = $\frac{1}{2}$
Spin > $\frac{1}{2}$

Nuclear Electric Quadrupole Moment couples to electric field gradient





Elements Accessible by NMR

1 IA												13 IIIA 14 IVA 15 VA 16 VIA 17 VIIA						18 VIIIA					
Hydrogen 1 H 1.007 940 8762 1.007 825 0322 1.007 570 2471																		Helium 2 He 4.002 603 2541					
Lithium 3 Li 6.941 150 83 7.016 199 808	Beryllium 4 Be 9.012 183 878											Boron 5 B 10.811 187 10 11.009 305 508	Carbon 6 C 12.010 738 10 12.010 738 10	Nitrogen 7 N 14.006 439 8 15.004 869 4	Oxygen 8 O 15.999 032 23 15.999 032 23	Fluorine 9 F 18.998 403 23 18.998 403 23	Neon 10 Ne 20.179 7 20.179 7						
Sodium 11 Na 22.989 769 28	Magnesium 12 Mg 24.304 67 24.304 67											Aluminum 13 Al 26.981 538 6 26.981 538 6	Silicon 14 Si 28.085 579 6 28.085 579 6	Phosphorus 15 P 30.973 761 2 30.973 761 2	Sulfur 16 S 32.065 034 4 32.065 034 4	Chlorine 17 Cl 35.453 064 4 35.453 064 4							
Potassium 19 K 39.098 309 1 39.098 309 1	Calcium 20 Ca 40.078 4 40.078 4	Scandium 21 Sc 44.955 908 6 44.955 908 6	Titanium 22 Ti 47.88 710 4 47.88 710 4	Vanadium 23 V 50.941 5 50.941 5	Chromium 24 Cr 51.996 15 51.996 15	Manganese 25 Mn 54.938 044 54.938 044	Iron 26 Fe 55.845 06 55.845 06	Cobalt 27 Co 58.933 195 58.933 195	Nickel 28 Ni 58.693 4 58.693 4	Copper 29 Cu 63.546 38 63.546 38	Zinc 30 Zn 65.38 65.38	Gallium 31 Ga 69.723 17 69.723 17	Germanium 32 Ge 72.630 08 72.630 08	Arsenic 33 As 74.921 60 74.921 60	Selenium 34 Se 78.96 78.96	Bromine 35 Br 79.904 18 79.904 18	Krypton 36 Kr 83.80 83.80						
Rubidium 37 Rb 85.467 8 85.467 8	Sr Strontium 38 Sr 87.62 87.62	Yttrium 39 Y 88.905 848 88.905 848	Zirconium 40 Zr 91.224 91.224	Niobium 41 Nb 92.906 38 92.906 38	Molybdenum 42 Mo 95.94 95.94	Technetium 43 Tc 98.906 254 98.906 254	Ruthenium 44 Ru 101.07 101.07	Rhodium 45 Rh 102.905 50 102.905 50	Palladium 46 Pd 106.363 10 106.363 10	Silver 47 Ag 107.868 2 107.868 2	Cadmium 48 Cd 112.411 8 112.411 8	Indium 49 In 114.818 38 114.818 38	Tin 50 Sn 118.710 4 118.710 4	Antimony 51 Sb 121.757 1 121.757 1	Tellurium 52 Te 127.603 127.603	Iodine 53 I 126.905 47 126.905 47	Xenon 54 Xe 131.294 4 131.294 4						
Cesium 55 Cs 132.905 45 132.905 45	Ba Barium 56 Ba 137.327 137.327	Lanthanum 57 La 138.905 47 138.905 47	Hafnium 72 Hf 178.49 178.49	Tantalum 73 Ta 180.947 88 180.947 88	Tungsten 74 W 183.84 183.84	Rhenium 75 Re 186.207 186.207	Osmium 76 Os 190.23 190.23	Iridium 77 Ir 192.222 192.222	Platinum 78 Pt 195.084 195.084	Gold 79 Au 196.966 57 196.966 57	Mercury 80 Hg 200.59 200.59	Thallium 81 Tl 204.383 3 204.383 3	Lead 82 Pb 207.2 207.2	Bismuth 83 Bi 208.980 4 208.980 4									
		Lanthanum 57 La 138.905 47 138.905 47	Thulium 69 Tm 168.930 4 168.930 4	Ytterbium 70 Yb 173.054 7 173.054 7																			

element — Hydrogen 1 — atomic number — 1
 symbol — **H**
 isotope — 1, 2, 3
 atomic weight — 1.007 940 8762, 2.014 101 778 1, 3.016 049 2
 spin number — 1/2, 0, 5/2
 frequency (MHz) — 14.324 036 1, 10.132 484 4, 27.045 047 9



□ I=1/2 nuclei
 ■ I=1/2 nuclei
 ■ I=1/2 and I=1/2 nuclei

Nuclear Spin

- NO stable nucleus has spin 2
- the highest value of spin for a stable nucleus is 7
 ^{176}Lu
- unstable nuclei
 - highest integral spin 16 - isomer ^{178}Hf
 - highest half-integer $37/2$ - isomer ^{177}Hf

Nuclear Spin

- Nuclei with spin $\frac{1}{2}$ - a spherical charge distribution
- Nuclei with $I > \frac{1}{2}$ - nonspherical charge distributions (prolate or oblate)





- Nuclei with a non-zero spin \rightarrow magnetic moment (μ)
- Nonspherical nuclei \rightarrow electric quadrupole moment (eQ)

Nuclear Spin

Properties of Nuclei:


Mass, Charge, Spin and Magnetism.



“The concept of spin is difficult. It was forced upon scientists by the experimental evidence”. Malcolm Levitt.

Rotating positive charge generates magnetic field

Nuclear Spin



Nuclear Spin

Nuclear spin = Spin angular momentum, P (*vector*)
(*moment hybnosti*)

Spin quantum number I


Magnetic quantum number m_I

Magnitude of P is quantized:

$$P = \frac{h}{2\pi} \sqrt{I(I+1)}$$

Direction with respect to the magnetic field B_0 is quantized:

$$P_z = \frac{h}{2\pi} m_I$$



Spin Angular Momentum, P

$$P = \frac{h}{2\pi} \sqrt{I(I+1)}$$

$$P_z = \frac{h}{2\pi} m_I$$


I = Nuclear spin quantum number
 $I = 0, 1/2, 1, 3/2, 5/2, 3, 7/2, \dots$

m_I = Nuclear spin magnetic quantum number

Multiplicity, M $2I + 1$ values
 $m_I = I, I - 1, I - 2, \dots, -I + 2, -I + 1, -I$

$$\cos \theta = \frac{P_z}{P} = \frac{m_I}{\sqrt{I(I+1)}}$$

\uparrow B_0 $^{59}\text{Co}, I = 7/2$



Spin Angular Momentum, P

$$P = \frac{h}{2\pi} \sqrt{I(I+1)}$$

I	$[I(I+1)]^{1/2}$
$1/2$	0.866
1	1.414
$3/2$	1.936
$5/2$	2.958
3	3.464
$7/2$	3.969
4	4.472
$9/2$	4.975


Spin Magnetic Moment, μ

The electrons, nucleons (protons, neutrons) and some nuclei possess **intrinsic magnetism**, which is not due to a circulating current.

Permanent magnetic moment similarly as spin angular momentum.

Magnetic moment, μ , is directly proportional to the spin angular momentum, P :

$$\mu = \gamma P$$



γ is the gyromagnetic (magnetogyric) ratio

Magnetogyric Ratio

γ - the magnetogyric ratio is the ratio of the nuclear magnetic moment μ to the nuclear angular momentum P .

$$\mu = \gamma P$$

γ - Important characteristic of nuclei

$$[\text{rad T}^{-1} \text{s}^{-1}]$$

Spin Magnetic Moment, μ


$$\mu = \gamma P = \gamma \hbar [I(I + 1)]^{1/2}$$

$$\mu_z = \gamma P_z = \gamma \hbar m_I$$


Nucleus	^1H	^2H	^{13}C	^{15}N	^{19}F	^{29}Si	^{31}P
$\gamma [10^{-7} \text{ rad T}^{-1}\text{s}^{-1}]$	26.75	4.11	6.73	-2.71	25.18	-5.32	10.84

electron

$$\gamma_e = 17\,609\,10^7 = 658 \gamma(\text{H})$$




Nuclear Spin in Magnetic Field



Nuclear Spin in Magnetic Field

Random orientation


Zeeman splitting to $2I + 1$ levels
Alignment of spins



No Field

Magnetic Field


Nuclear Spin in Magnetic Field



magnetic dipole

Nuclear Spin in Magnetic Field

- An angular momentum is associated with each rotating object
- A nuclear spin possesses a magnetic moment μ arising from the angular momentum of the nucleus
- The magnetic moment μ is a vector perpendicular to the current loop
- In a magnetic field (B) the magnetic moment behaves as a magnetic dipole




$$\mu = i A$$

Nuclear Spin in Magnetic Field

In B_0 , a magnetic moment μ is directed at some angle w.r.t. B_0 direction




the B_0 field will exert a torque on the magnetic moment. This causes μ to precess about the magnetic field direction

Torque is the rate of change of the nuclear spin angular momentum



Nuclear Spin in Magnetic Field

Spin precession in the external magnetic field.



Quantum description of precession shows that the frequency of the motion is:

$$\omega_0 = -\gamma B_0 \text{ [rad s}^{-1}\text{]} \text{ or } \nu_0 = -\gamma B_0 / 2\pi \text{ [Hz]}$$


It is called the Larmor frequency (if $\gamma > 0$ then $\nu_0 < 0$)

Larmor Frequency

$$\omega_0 = -\gamma B_0 \text{ [rad s}^{-1}\text{]}$$

$$\nu_0 = -\gamma B_0 / 2\pi \text{ [Hz]}$$


$$\nu_0 = -\frac{\gamma B_0}{2\pi}$$



Larmor Frequency



Sir Joseph Larmor
(1857-1942)



$$\omega_0 = -\gamma B_0 \text{ [rad s}^{-1}\text{]} \text{ or } \nu_0 = -\gamma B_0 / 2\pi \text{ [Hz]}$$

Nuclear Zeeman Effect - Splitting

Nuclei are charged and if they have spin, they are magnetic



No Field

Higher energy state: magnetic field opposes applied field



$m_I - 1/2$

Energy of transition = energy of radiowaves

B

Applied Magnetic Field = B_0

Lower energy state: magnetic field aligned with applied field



$m_I + 1/2$

Nuclear Spin in Magnetic Field

The magnetic energy depends on the interaction between the magnetic moment and B_0 field:




$$E_{\text{mag}} = -\boldsymbol{\mu} \cdot \mathbf{B}_0 \quad (\text{a scalar product of 2 vectors})$$

$$E_{\text{mag}} = -\mu_z B = -\gamma P_z B$$

$$E_{\text{mag}} = -m_I \hbar \gamma B$$

NMR selection rule $\Delta m_I = \pm 1$


CALVIN AND HOBBS By Bill Watterson



Spin in Magnetic Field

$$I = 1/2 \quad E_{m = -1/2}$$


$$E_{m = 1/2}$$





$$\Delta E_{\text{mag}} = E_{m=-1/2} - E_{m=1/2} = \Delta m_I \hbar \gamma B = h \nu \Rightarrow \nu = \gamma B / 2\pi$$


The frequency of the electromagnetic radiation that corresponds to the energy difference between the two energy levels is equal to the precessional frequency of the nuclei.

Spin in Magnetic Field





Excitation of NMR Spin





Energy Levels for $I = 1/2$




$$\Delta E_{\text{mag}} = E_{m=-1/2} - E_{m=1/2} = \Delta m_I \hbar \gamma B = \hbar \gamma B / 2\pi$$

Protons


$$\Delta E = (6.626 \times 10^{-34} \text{ J s } \cdot 26.75 \times 10^7 \text{ rad T}^{-1}\text{s}^{-1} \cdot 11.743 \text{ T}) / 2\pi = 3.313 \times 10^{-25} \text{ J}$$

very small energy difference

Energy Levels for $I = 1/2$



Energy Levels for $I = \frac{1}{2}$



Boltzmann Distribution

The excess of nuclei on the lower energy level is given by Boltzmann distribution:

$$\frac{N_{\uparrow\downarrow}}{N_{\uparrow\uparrow}} = \exp\left(-\frac{\Delta E}{k_B T}\right) = \exp\left(-\frac{\hbar \gamma B}{k_B T}\right) =$$

$$= \exp(-3.313 \cdot 10^{-25} / 4.101 \cdot 10^{-21}) = \exp(-8.078 \cdot 10^{-5}) = 0.99991922$$

If $N_{\uparrow\uparrow} = 1\,000\,000$ then $N_{\uparrow\downarrow} = 999919$

Only 81 out of 2 million ^1H nuclei contribute to NMR signal at 500 MHz!

$$\hbar = 1.055 \cdot 10^{-34} \text{ J s}$$

$$\gamma_{\text{H}} = 26.75 \cdot 10^7 \text{ rad T}^{-1}\text{s}^{-1}$$





$$B = 11.7433 \text{ T (500 MHz)}$$

$$k_B = 1.3807 \cdot 10^{-23} \text{ J K}^{-1}$$

$$T = 297 \text{ K}$$

Boltzmann Distribution

$$N_{\uparrow\downarrow} / N_{\uparrow\uparrow} = \exp(-\Delta E / k_B T) = \exp(-\hbar \gamma B / k_B T)$$




the stronger the field and the higher the magnetogyric ratio, the larger the population difference

the higher the temperature, the smaller the population difference

Boltzmann Distribution


The higher the field B ,
the larger the energy difference,
the larger the population difference,
the larger the net magnetization,
and the bigger the NMR signal



Nuclear Magnetic Resonance (NMR)

- **Nuclear** – spin $\frac{1}{2}$ nuclei (e.g. protons) behave as tiny bar magnets.
- **Magnetic** – a strong magnetic field causes a small energy difference between $+\frac{1}{2}$ and $-\frac{1}{2}$ spin states.
- **Resonance** – photons of radio waves can match the exact energy difference between the $+\frac{1}{2}$ and $-\frac{1}{2}$ spin states resulting in absorption of photons as the protons change spin states.

Magnetization




More nuclei point in parallel to the static magnetic field.

The macroscopic magnetic moment, M_0


$$M_0 = \sum \mu_i$$

In-Field




Bloch equations: the nuclear magnetization $M = (M_x, M_y, M_z)$ as a function of time and relaxation times T_1 and T_2


Longitudinal Magnetization



No magnetic field




Magnetic field



$$\frac{d}{dt} M_z = R_1 (M_{eq} - M_z) = \frac{1}{T_1} (M_{eq} - M_z)$$

$$M_z(t) = M_{eq} \left(1 - e^{\frac{-(t-t_{on})}{T_1}} \right)$$



Spin-Lattice Relaxation Time



$R_1 = 1/T_1$ [Hz] longitudinal relaxation rate constant

T_1 [s] longitudinal relaxation time

spin-lattice relaxation time

enthalpy

Transverse Magnetisation



Spin coherence

$$\frac{d}{dt} M_{xy} = -R_2 M_{xy} = \frac{-1}{T_2} M_{xy} \quad M_{xy}(t) = M_{eq} e^{\frac{-t}{T_2}}$$

Spin-Spin Relaxation Time


$R_2 = 1/T_2$ [Hz] transverse relaxation rate constant

T_2 [s] transverse relaxation time constant



spin-spin relaxation time

entropy


One RF Pulse






Effect of a 90°_x pulse
The magnetization vector is rotated to the y axis





Fourier transform



Relaxation



Free Induction Decay FID



$$\begin{aligned}
 S(t) &= S_{Re}(t) + iS_{Im}(t) \\
 &= [\cos(\omega_L t) + i\sin(\omega_L t)] \exp(-t/T_2) \\
 &= \exp(i\omega_L t) \exp(-t/T_2) = \exp[-(1/T_2 - i\omega_L)t]
 \end{aligned}$$

