

Inorganic Chemistry III

C4010

Prof. RNDr. Jiří Příhoda, CSc.

Prof. RNDr. Jiří Pinkas, Ph.D.


Syllabus

Část I. Prof. Příhoda

1. Koordinační chemie
2. Chelatující ligandy
3. Ionty v roztoku
4. Makroseparační metody kovů
5. Transurany













Část II. Prof. Pinkas

6. Periodic Table
7. + 8. Chemical Bonding
9. Acid-base Chemistry
10. + 11. Rings and Polyhedra
12. + 13. Magnetochemistry, Moessbauer spectroscopy








Textbooks

Reading assignments = FIND ERRORS



Journals

Reading assignments



IUPAC Periodic Table of the Elements

1																	18
1 H hydrogen 1.008 (1.00784, 1.00811)																	2 He helium 4.0026
3 Li lithium 6.94 (6.938, 6.991)	4 Be beryllium 9.0122											5 B boron 10.81 (10.806, 10.821)	6 C carbon 12.011 (12.009, 12.012)	7 N nitrogen 14.007 (14.005, 14.009)	8 O oxygen 15.999 (15.998, 16.003)	9 F fluorine 18.998	10 Ne neon 20.180
11 Na sodium 22.990	12 Mg magnesium 24.305 (24.304, 24.307)											13 Al aluminium 26.982	14 Si silicon 28.086 (28.084, 28.089)	15 P phosphorus 30.974	16 S sulfur 32.06 (32.059, 32.075)	17 Cl chlorine 35.45 (35.446, 35.457)	18 Ar argon 39.948
19 K potassium 39.098	20 Ca calcium 40.078(4)	21 Sc scandium 44.956	22 Ti titanium 47.867	23 V vanadium 50.942	24 Cr chromium 51.996	25 Mn manganese 54.938	26 Fe iron 55.845(2)	27 Co cobalt 58.933	28 Ni nickel 58.693	29 Cu copper 63.546(3)	30 Zn zinc 65.38(2)	31 Ga gallium 69.723	32 Ge germanium 72.630(8)	33 As arsenic 74.922	34 Se selenium 78.971(8)	35 Br bromine 79.904 (79.901, 79.907)	36 Kr krypton 83.798(2)
37 Rb rubidium 85.468	38 Sr strontium 87.62	39 Y yttrium 88.906	40 Zr zirconium 91.224(2)	41 Nb niobium 92.906	42 Mo molybdenum 95.94	43 Tc technetium	44 Ru ruthenium 101.07(2)	45 Rh rhodium 102.91	46 Pd palladium 106.42	47 Ag silver 107.87	48 Cd cadmium 112.41	49 In indium 114.82	50 Sn tin 118.71	51 Sb antimony 121.76	52 Te tellurium 127.60(3)	53 I iodine 126.90	54 Xe xenon 131.29
55 Cs caesium 132.91	56 Ba barium 137.33	57-71 Lanthanoids	72 Hf hafnium 178.49(2)	73 Ta tantalum 180.95	74 W tungsten 183.84	75 Re rhenium 186.21	76 Os osmium 190.23(2)	77 Ir iridium 192.22	78 Pt platinum 195.08	79 Au gold 196.97	80 Hg mercury 200.59	81 Tl thallium 204.38 (204.38, 204.39)	82 Pb lead 207.2	83 Bi bismuth 208.98	84 Po polonium	85 At astatine	86 Rn radon
87 Fr francium	88 Ra radium	89-103 actinoids	104 Rf rutherfordium	105 Db dubnium	106 Sg seaborgium	107 Bh bohrium	108 Hs hassium	109 Mt meitnerium	110 Ds darmstadtium	111 Rg roentgenium	112 Cn copernicium	113 Nh nihonium	114 Fl flerovium	115 Mc moscovium	116 Lv livermorium	117 Ts tennessine	118 Og oganesone

Key:
 atomic number
Symbol
 name
 conventional atomic weight
 standard atomic weight



57 La lanthanum 138.91	58 Ce cerium 140.12	59 Pr praseodymium 140.91	60 Nd neodymium 144.24	61 Pm promethium	62 Sm samarium 150.36(2)	63 Eu europium 151.96	64 Gd gadolinium 157.25(2)	65 Tb terbium 158.93	66 Dy dysprosium 162.50	67 Ho holmium 164.93	68 Er erbium 167.26	69 Tm thulium 168.93	70 Yb ytterbium 173.05	71 Lu lutetium 174.967
89 Ac actinium	90 Th thorium 232.04	91 Pa protactinium 231.04	92 U uranium 238.03	93 Np neptunium	94 Pu plutonium	95 Am americium	96 Cm curium	97 Bk berkelium	98 Cf californium	99 Es einsteinium	100 Fm fermium	101 Md mendelevium	102 No nobelium	103 Lr lawrencium

For notes and updates to this table, see www.iupac.org. This version is dated 28 November 2016.
 Copyright © 2016 IUPAC, the International Union of Pure and Applied Chemistry.

Periodic Table of the Elements

1 H Hydrogen 1.008																	2 He Helium 4.003
3 Li Lithium 6.941	4 Be Beryllium 9.012											5 B Boron 10.811	6 C Carbon 12.011	7 N Nitrogen 14.007	8 O Oxygen 15.999	9 F Fluorine 18.998	10 Ne Neon 20.180
11 Na Sodium 22.990	12 Mg Magnesium 24.305											13 Al Aluminum 26.982	14 Si Silicon 28.086	15 P Phosphorus 30.974	16 S Sulfur 32.066	17 Cl Chlorine 35.453	18 Ar Argon 39.948
19 K Potassium 39.098	20 Ca Calcium 40.078	21 Sc Scandium 44.956	22 Ti Titanium 47.867	23 V Vanadium 50.942	24 Cr Chromium 51.996	25 Mn Manganese 54.938	26 Fe Iron 55.845	27 Co Cobalt 58.933	28 Ni Nickel 58.693	29 Cu Copper 63.546	30 Zn Zinc 65.38	31 Ga Gallium 69.723	32 Ge Germanium 72.631	33 As Arsenic 74.922	34 Se Selenium 78.971	35 Br Bromine 79.904	36 Kr Krypton 84.798
37 Rb Rubidium 84.468	38 Sr Strontium 87.62	39 Y Yttrium 88.906	40 Zr Zirconium 91.224	41 Nb Niobium 92.906	42 Mo Molybdenum 95.95	43 Tc Technetium 98.907	44 Ru Ruthenium 101.07	45 Rh Rhodium 102.906	46 Pd Palladium 106.42	47 Ag Silver 107.868	48 Cd Cadmium 112.414	49 In Indium 114.818	50 Sn Tin 118.711	51 Sb Antimony 121.760	52 Te Tellurium 127.6	53 I Iodine 126.904	54 Xe Xenon 131.294
55 Cs Cesium 132.905	56 Ba Barium 137.328	57-71 Lanthanides	72 Hf Hafnium 178.49	73 Ta Tantalum 180.948	74 W Tungsten 183.84	75 Re Rhenium 186.207	76 Os Osmium 190.23	77 Ir Iridium 192.217	78 Pt Platinum 195.085	79 Au Gold 196.967	80 Hg Mercury 200.592	81 Tl Thallium 204.383	82 Pb Lead 207.2	83 Bi Bismuth 208.980	84 Po Polonium [209]	85 At Astatine 209	86 Rn Radon 222.018
87 Fr Francium 223.020	88 Ra Radium 226.025	89-103 Actinides	104 Rf Rutherfordium [261]	105 Db Dubnium [262]	106 Sg Seaborgium [266]	107 Bh Bohrium [264]	108 Hs Hassium [269]	109 Mt Meitnerium [268]	110 Ds Darmstadtium [269]	111 Rg Roentgenium [272]	112 Cn Copernicium [277]	113 Nh Nihonium unknown	114 Fl Flerovium [289]	115 Mc Moscovium unknown	116 Lv Livermorium [298]	117 Ts Tennessine unknown	118 Og Oganesson unknown

57 La Lanthanum 138.905	58 Ce Cerium 140.116	59 Pr Praseodymium 140.908	60 Nd Neodymium 144.243	61 Pm Promethium 144.913	62 Sm Samarium 150.36	63 Eu Europium 151.964	64 Gd Gadolinium 157.25	65 Tb Terbium 158.925	66 Dy Dysprosium 162.500	67 Ho Holmium 164.930	68 Er Erbium 167.259	69 Tm Thulium 168.934	70 Yb Ytterbium 173.055	71 Lu Lutetium 174.967
89 Ac Actinium 227.028	90 Th Thorium 232.038	91 Pa Protactinium 231.036	92 U Uranium 238.029	93 Np Neptunium 237.048	94 Pu Plutonium 244.064	95 Am Americium 243.061	96 Cm Curium 247.070	97 Bk Berkelium 247.070	98 Cf Californium 251.080	99 Es Einsteinium [254]	100 Fm Fermium 257.095	101 Md Mendelevium 258.1	102 No Nobelium 259.101	103 Lr Lawrencium [262]

Alkali Metal	Alkaline Earth	Transition Metal	Basic Metal	Semimetal	Nonmetal	Halogen	Noble Gas	Lanthanide	Actinide
--------------	----------------	------------------	-------------	-----------	----------	---------	-----------	------------	----------

© 2020
Todd Helmenstine
www.ck12.org

Periodic Table of the Elements

Long form

		Group																																	
		1	2	3											4	5	6	7	8	9	10	11	12	13	14	15	16	17	18						
1	1 H																									2 He									
2	3 Li	4 Be																			5 B	6 C	7 N	8 O	9 F	10 Ne									
3	11 Na	12 Mg																			13 Al	14 Si	15 P	16 S	17 Cl	18 Ar									
4	19 K	20 Ca	21 Sc																		22 Ti	23 V	24 Cr	25 Mn	26 Fe	27 Co	28 Ni	29 Cu	30 Zn	31 Ga	32 Ge	33 As	34 Se	35 Br	36 Kr
5	37 Rb	38 Sr	39 Y																		40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe
6	55 Cs	56 Ba	57 La	58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu	72 Hf	73 Ta	74 W	75 Re	76 Os	77 Ir	78 Pt	79 Au	80 Hg	81 Tl	82 Pb	83 Bi	84 Po	85 At	86 Rn			
7	87 Fr	88 Ra	89 Ac	90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr	104 Rf	105 Db	106 Sg	107 Bh	108 Hs	109 Mt	110 Ds	111 Rg	112 Cn	113 Nh	114 Fl	115 Mc	116 Lv	117 Ts	118 Og			

Periodic Table of the Elements



IUPAC 2017

The image shows a blank periodic table grid. The main body of the table is a grid of 18 columns and 7 rows. The first two columns are on the left, and the last two columns are on the right. The middle 14 columns are connected. A thick horizontal line is drawn below the 7th row, separating the main body from the lanthanide and actinide series. Below this line, there are two columns of two cells each, representing the lanthanide and actinide series. The top two cells of each of these columns contain diagonal lines, indicating they are not part of the main grid.



Orbital Energies in Polyelectronic Atoms

Schroedinger equation

$$\hat{H} \Psi = E \Psi$$




Radial Density Distribution Function



Number of nodal surfaces = $n - l - 1$

Aufbau Principle

The order of orbital filling – not the order of atomic energies



the Madelung rule:

Electrons (e) fill orbitals starting at the lowest available energy state before filling higher states.

The higher the total number of nodes in the atomic orbital, $n + l$, the higher is its energy.

If $n + l$ is the same, then lower n

Hund's Rules

State with **the largest value of S is most stable** and stability decreases with decreasing S
= electrons always enter an empty orbital before they pair up.

The **e** in singly occupied orbitals are less effectively screened or shielded from the nucleus = more tightly bound.

A Coulomb repulsion between two **e** in the same orbital = a spin pairing energy.


For states with same values of S, the state with the largest value of L is the most stable.

The total (scalar) angular momentum, the *relative* momentum vectors of the various **e**.

If states have same values of L and S then, for a subshell that is less than half filled, state with smallest J is most stable; for subshells that are more than half filled, state with largest value of J is most stable.



Coulomb repulsion



Exchange energy



Exchange Energy

Exchange energy (K) = the energy released when two or more e with **the same spin** exchange their positions in the degenerate orbitals of a subshell


The result of spin correlation = e do not move independently

A pair of e with parallel spins $\uparrow\uparrow$ has less repulsion than a pair with anti-parallel spins $\uparrow\downarrow$

$\uparrow\downarrow$ occupy smaller volume than $\uparrow\uparrow$ \Rightarrow $\uparrow\downarrow$ experience more repulsion than $\uparrow\uparrow$

Exchange energy **decreases** orbital energy - lower orbital energy = the orbital is closer to nucleus - electrons are more tightly bound which gives extra stability of electrons of valence shell

Relative exchange energy = number of pairs with parallel spins $\times K$





Exchange Energy

Calculate exchange energies for configurations p^1 to p^6



Calculate exchange energies for configurations d^1 to d^{10}




exchange energy for $d^5 = 10 K$

For every pair of electrons with parallel spins, there is a corresponding exchange energy K that reduces the value of the orbital energy

Which configuration is the most stable $\text{Cr (Ar) } 3d^4 4s^2$ or $3d^5 4s^1$

Orbital Energies



Cu Ag, Au
 Cr, Mo, W
 Ru, Rh, Pd
 Nb
 La, Ce, Pr, Gd
 Cm, Th, Pa, Np

Ni [Ar] 3d⁹ 4s¹ (4p⁰) free atom in vacuum


Ni [Ar] 3d⁸ 4s² (4p⁰) very close in energy

Ni [Ar] 3d¹⁰ (4s⁰ 4p⁰) in molecules, Ni(CO)₄



Lanthanides

Xe	$[\text{Kr}] 4d^{10} 5s^2 5p^6$	$E(4f) > E(6s)$
Cs	$[\text{Xe}] 6s^1 4f^0 5d^0$	
Ba	$[\text{Xe}] 6s^2 4f^0 5d^0$	
La	$[\text{Xe}] 4f^0 5d^1 6s^2$	transition metal
Ce	$[\text{Xe}] 4f^1 5d^1 6s^2$	$E(4f) < E(6s), E(5d)$
Pr	$[\text{Xe}] 4f^3 6s^2$	
Eu	$[\text{Xe}] 4f^7 5s^2 5p^6 5d^0 6s^2$	
Gd	$[\text{Xe}] 4f^8 5s^2 5p^6 5d^0 6s^2$	
Gd	$[\text{Xe}] 4f^7 5s^2 5p^6 5d^1 6s^2$	4f half-filled
Lu	$[\text{Xe}] 4f^{14} 5d^1 6s^2$	4f full



Ionization Energies




Electron Affinities




Atomic Radii




Atomic Radii



Atomic Volumes



Metallic Radii CN = 12



Metallic radii for 12-coordination (Å)

Radius vs. CN:

CN = 12 Rel. R = 1.00

CN = 8 Rel. R = 0.97

CN = 6 Rel. R = 0.96


CN = 4 Rel. R = 0.88

solid solutions: a linear relationship = Vegard

Li	Be														
1.57	1.12														
Na	Mg	Al													
1.91	1.60	1.43													
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge		
2.35	1.97	1.64	1.47	1.35	1.29	1.37	1.26	1.25	1.25	1.28	1.37	1.53	1.39		
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	
2.50	2.15	1.82	1.60	1.47	1.40	1.35	1.34	1.34	1.37	1.44	1.52	1.67	1.58	1.61	
Cs	Ba	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	
2.72	2.24	1.88	1.59	1.47	1.41	1.37	1.35	1.36	1.39	1.44	1.55	1.71	1.75	1.82	
4f elements:			Ce (1.82)–Lu (1.72) but Eu, 2.06, Yb, 1.94												
5f elements:			Th	Pa	U	Np	Pu	Am	Cm						
			1.80	1.63	1.56	1.56	1.64	See Table 29.4							

Ionic Radii

r, Å



Oxidation State

1835 Wöhler Unorganische Chemie – oxydationsstufe

IUPAC “Gold Book” **Compendium of Chemical Terminology**

The oxidation state is the atom's charge after ionic approximation of its bonds.


1. Atoms in an element have oxidation state 0.
2. The sum of the oxidation states for atoms in a compound is 0, in ions the algebraic sum of the oxidation states of the constituent atoms must be equal to the charge on the ion.
3. Fluorine in compounds has the oxidation state -1 .
4. Alkaline metals in compounds have the oxidation state $+1$, alkaline-earth metals $+2$.
5. Hydrogen in compounds has the oxidation state $+1$ (except hydrides -1).
6. Oxygen in compounds has the oxidation state -2 (except peroxides -1).

Oxidation State


The oxidation state of an atom in a compound is given by the hypothetical charge of the corresponding atomic ion that is obtained by **heterolytically** cleaving its bonds such that the atom with the higher electronegativity in a bond is allocated all electrons in this bond. Bonds between like atoms (having the same formal charge) are cleaved **homolytically**.

Criteria for the ionic approximation:

- 1) Extrapolation of the bond's polarity
 - a) from the electronegativity difference
 - b) from the dipole moment
 - c) from quantum-chemical calculations of charges
- 2) Assignment of electrons according to the atom's contribution to the molecular orbital (MO)



Oxidation State



	NBO/DF-BP86/def2-SVP	Lewis Character
1	$\cdot\ddot{\text{N}}-\ddot{\text{N}}=\ddot{\text{O}}\cdot$	90.6 %
2	$:\text{N}\equiv\text{N}=\ddot{\text{O}}\cdot$	88.9 %*
3	$:\text{N}\equiv\overset{\oplus}{\text{N}}-\ddot{\text{O}}\ominus$	96.4 %
4	$\cdot\overset{\ominus}{\text{N}}=\overset{\oplus}{\text{N}}=\ddot{\text{O}}\cdot$	94.0 %
5	$\ominus\ominus\cdot\ddot{\text{N}}-\overset{\oplus}{\text{N}}\equiv\overset{\oplus}{\text{O}}\cdot$	91.7 %

* if treated as excited state.

Electronegativity - the 3rd Dimension of the Periodic Table

Pauling 1932

electronegativity χ_p based on bond energies, ability of an atom in a molecule to attract electrons for itself

Mulliken 1934

orbital, “absolute” electronegativity χ_M is an average of ionization energy and electron affinity

Allred + Rochow 1958

χ_A electronegativity is “a force” needed to remove an electron from the atom’s ground state (similarity to Coulomb’s Law).

Electronegativity - the 3rd Dimension of the Periodic Table

Allen 1992

average valence-electron energy of the free atom, χ_{spec}


Sanderson

ratio of average electron density to that of corresponding rare gas atom, tendency of an atomic orbital to attract electrons from another atomic orbital with which it has combined to form a bond, chemical potential

Mulliken-Jaffe

electronegativity is the slope of the energy-vs-charge curve as it passes through the origin

Electronegativity - the 3rd Dimension of the Periodic Table



Mulliken Electronegativity

		in eV											
H													
s	7.2												
Li		Be		B		C		N		O		F	
s	3.1	di ²	4.8	tr ³	6.4	di ² π ²	10.4, 5.7	di ³ π ²	15.7, 7.9	tr ⁵ π	17.1, 20.2	s	31.3
p	1.8	te ²	3.9	te ³	6.0	tr ³ π	8.8, 5.6	tr ⁴ π	12.9, 8.0	di ² π ⁴	19.1	p	12.2
						te ⁴	8.0	te ⁵	11.6	te ⁶	15.3		
Na		Mg		Al		Si		P		S		Cl	
s	2.9	di ²	4.1	tr ³	5.5	di ² π ²	9.0, 5.7	di ³ π ²	11.3, 6.7	tr ⁴ π ²	10.9	s	19.3
p	1.6	te ²	3.3	te ³	5.4	tr ³ π	7.9, 5.6	tr ⁴ π	9.7, 6.7	te ⁶	10.2	p	9.4
						te ⁴	7.3	te ⁵	8.9				
K		Ca		Ga		Ge		As		Se		Br	
s	2.9	di ²	3.4	tr ³	6.0	di ² π ²	9.8, 6.5	di ³ π ²	9.0, 6.5	tr ⁴ π ²	10.6	s	18.3
p	1.8	te ²	2.5	te ³	6.6	tr ³ π	8.7, 6.4	tr ⁴ π	8.6, 7.0	te ⁶	9.8	p	8.4
						te ⁴	8.0	te ⁵	8.3				
Rb		Sr		In		Sn		Sb		Te		I	
s	2.1	di ²	3.2	tr ³	5.3	di ² π ²	9.4, 6.5	di ³ π ²	9.8, 6.3	tr ⁴ π ²	10.5	s	15.7
p	2.2	te ²	2.2	te ³	5.1	tr ³ π	8.4, 6.5	tr ⁴ π	9.0, 6.7	te ⁶	9.7	p	8.1
								te ⁵	8.5				

Values can be computed only for orbitals holding 1 electron. For the carbon and nitrogen families it is possible to have both hybrid and π atomic orbitals half-filled. *digonal* ≡ *sp* hybrid, *trigonal* ≡ *sp²* hybrid, *tetrahedral* ≡ *sp³* hybrid.

$$\chi_M = \frac{1}{2}(IE + EA) = -\mu$$

Chemical potential of the electrons


Mulliken Electronegativity

$$\chi_M = 0.178(IE + EA) + 0.17$$

Pauling scale, IE and EA in eV

$$\chi_M = 1.97 \times 10^{-3}(IE + EA) + 0.19$$

Pauling scale, IE and EA in kJ mol⁻¹



Allen Electronegativities

Electronegativity = the *average* one-electron energy of the valence-shell electrons in ground-state free atoms.

L.C. Allen, J. Am. Chem. Soc., 1989, **111**, 9003-9014

Only the Allen electronegativity is truly independent of the oxidation state, as it relates to the average valence-electron energy of the free atom.

Spectroscopic electronegativity:

m, n = number of p and s electrons, respectively

ϵ_s, ϵ_p = multiplet-averaged ionization potentials of s and p electrons (experimental data).

$$\chi_{spec} = \frac{m\epsilon_p + n\epsilon_s}{m + n}$$

Allen Electronegativities





Table 1: Allen electronegativities^[18–20] (in Pauling units).

H										He
2.300										4.16
Li	Be	B	C	N	O	F				Ne
0.912	1.576	2.051	2.544	3.066	3.610	4.193				4.787
Na	Mg	Al	Si	P	S	Cl				Ar
0.912	1.293	1.613	1.916	2.253	2.589	2.869				3.242
K	Ca	Ga	Ge	As	Se	Br				Kr
0.734	1.034	1.756	1.994	2.211	2.424	2.685				2.966
Rb	Sr	In	Sn	Sb	Te	I				Xe
0.706	0.963	1.656	1.834	1.984	2.158	2.359				2.582
Cs	Ba	Tl	Pb	Bi	Po	At				Rn
0.659	0.881	1.789	1.854	2.01	2.19	2.39				2.60
Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	
1.19	1.38	1.53	1.65	1.75	1.80	1.84	1.88	1.85	1.59	
Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	
1.12	1.32	1.41	1.47	1.51	1.54	1.56	1.58	1.87	1.52	
Lu ^[a]	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	
1.09	1.16	1.34	1.47	1.60	1.65	1.68	1.72	1.92	1.76	

[a] The variation across the lanthanoid series has not been evaluated.

Sanderson Electronegativities

The ability of an atom or ion to attract electrons to itself is dependent upon the **effective nuclear charge** felt by the outermost valence electrons.

As the effective nuclear charge increases the size of the atom decreases.

The Sanderson electronegativity S is proportional to the compactness of an atom:

$$S \approx \frac{Z}{\frac{4}{3}\pi r^3} \approx \frac{D}{D_0}$$

D is the electron density of an atom (its atomic number divided by its atomic volume (covalent radius cubed)) and D_0 is the expected electron density of an atom, calculated from extrapolation between the noble gas elements.

$$\sqrt{\chi_P} = a \times S + b$$

Electronegativity Equalization

When two or more atoms initially different in electronegativity combine chemically, they adjust to have the same intermediate electronegativity within the compound.

This intermediate electronegativity is given by **the geometric mean** of the individual electronegativities of the component atoms.

The electron density will flow from the more electropositive atom to the more electronegative atom, creating a partial positive charge on the former and a partial negative charge on the latter. As the positive charge on the electropositive atom increases, its effective nuclear charge increases, hence its electronegativity increases. The same trend happens in the opposite direction for the more electronegative atom, until the two have the same electronegativity – electronegativities are equalized and charge transfer stops.


The partial ionic charge and ionic radius of an ion are not constants, but vary depending upon the electronegativity of the surrounding atoms. Furthermore, the partial charge on an atom never exceeds +1 or -1.

The oxidation states do not reflect the true charge of an ion.

Electronegativity Equalization

$$S(\text{SnO}_2) = (S_{\text{Sn}} \times S_{\text{O}}^2)^{1/3} = (4.28 \times 4.90^2)^{1/3} = 4.68$$

$$S(\text{SrTiO}_3) = (S_{\text{Sr}} \times S_{\text{Ti}} \times S_{\text{O}}^3)^{1/5} = (1.28 \times 2.09 \times 4.90^3)^{1/5} = 3.16$$



Electronegativity Equalization

Electronegativity of an atom in an A–B bond depends on its partial charge, δ

$$\chi_A = a_A + b_A \delta_A \qquad \chi_B = a_B + b_B \delta_B$$

Charge transfer causes that electronegativities are equalized

$$\chi_A = \chi_B \qquad \delta_A = -\delta_B$$

$$\delta_A = \frac{a_B - a_A}{b_A + b_B}$$

Sanderson Electronegativities

A–B

$$dG = dU + TdS - pdV + \Sigma\mu_i dN_i$$

$$dS = 0, pdV = 0$$

$$dG = dU + \Sigma\mu_i dN_i$$

$$dN_A = -dN_B = dN$$

$$dG = dU + (\mu_A - \mu_B)dN$$

$$\mu = \left(\frac{\partial G}{\partial N} \right)_{V,S,n_i}$$

Mullikan-Jaffe Electronegativity

When the ionization energies and electron affinities of atoms are plotted against the oxidation state a smooth curve results and the total energy can be described by a quadratic formula

$$E = \alpha q + \beta q^2$$


Jaffe said that the Mullikan definition of electronegativity is the slope of this curve as it passes through the origin

E = total energy of an atom

q = ionic charge

$q = 0$ – neutral atom $E = 0$.

$$\chi_M = \left(\frac{\partial E}{\partial q} \right) = \alpha + 2\beta q$$



Mulliken-Jaffe Electronegativity

For $q = +1$ in the equation, the energy of the system corresponds to **the first ionization energy IE**

and
$$E = \alpha q + \beta q^2$$

for $q = -1$ the energy of the system corresponds to the **negative of electron affinity EA**.

$$IE_V = \alpha + \beta \quad (2) \quad \text{and} \quad EA_V = -\alpha + \beta \quad (3)$$

Therefore on adding equation (2) and (-3): $IE_V + EA_V = 2\alpha$

where α stands for **Mulliken electronegativity χ_M**

Mulliken-Jaffe Electronegativity

χ_M involves valence state ionization energy and electron affinities (calculated for atoms in the valence state as they exist in a molecule), takes into account the electronegativity of partially charged atoms emphasizing the effect the partial charge can have on the electronegativity of the atom.

$$E = \alpha q + \beta q^2 \quad \chi_M = \left(\frac{\partial E}{\partial q} \right) = \alpha + 2\beta q = a + b\delta$$

δ = the partial ionic charge

a = inherent or neutral atom electronegativity

b = charge coefficient (rate of change of electronegativity with charge)

Large soft atoms – low b

Small hard nonpolarizable atoms – high b

$1/b$ = charge capacity, ability to donate/absorb electrons

Hybridization and Electronegativity

Hybridization has a profound effect on electronegativity:

The s orbitals have lower energy hence greater electron attracting power

The greater the s character - the greater the electronegativity

The factor affecting the acidity of hydrogen is the difference in electronegativity between hydrogen and the atom to which it is bonded
Acetylene has highly acidic hydrogen as the carbon atom is sp hybridized (50% s character) and has almost the same electronegativity as chlorine atom.


This is in contrast to methane in which the carbon atom is sp³ hybridized (25% s character) and is not reactive

As the s character in hybridization of nitrogen increases the basicity of nitrogen will decrease. In ethane nitrile N is sp hybridized and hence very electronegative. The more electronegative the nitrogen atom less is the tendency to share the lone pair of electron. The basicity therefore decreases.


Hybridization and Electronegativity

Hybridization	sp^3	sp^2	sp
C	2.48	2.75	3.29
N	3.68	3.94	4.67

Electronegativity - the 3rd Dimension of the Periodic Table




Electronegativity and Chemical Properties



Chemical Properties

$$\frac{r}{IE}$$



Chemical Properties

V·T·E

Periodic table of electronegativity by Pauling scale

→ Atomic radius decreases → Ionization energy increases → Electronegativity increases →

Group →	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	
↓ Period																			
1	H 2.20																		He
2	Li 0.98	Be 1.57											B 2.04	C 2.55	N 3.04	O 3.44	F 3.98		Ne
3	Na 0.93	Mg 1.31											Al 1.61	Si 1.90	P 2.19	S 2.58	Cl 3.16		Ar
4	K 0.82	Ca 1.00	Sc 1.36	Ti 1.54	V 1.63	Cr 1.66	Mn 1.55	Fe 1.83	Co 1.88	Ni 1.91	Cu 1.90	Zn 1.65	Ga 1.81	Ge 2.01	As 2.18	Se 2.55	Br 2.96		Kr 3.00
5	Rb 0.82	Sr 0.95	Y 1.22	Zr 1.33	Nb 1.6	Mo 2.16	Tc 1.9	Ru 2.2	Rh 2.28	Pd 2.20	Ag 1.93	Cd 1.69	In 1.78	Sn 1.96	Sb 2.05	Te 2.1	I 2.66		Xe 2.60
6	Cs 0.79	Ba 0.89	*	Hf 1.3	Ta 1.5	W 2.36	Re 1.9	Os 2.2	Ir 2.20	Pt 2.28	Au 2.54	Hg 2.00	Tl 1.62	Pb 1.87	Bi 2.02	Po 2.0	At 2.2		Rn 2.2
7	Fr 0.7	Ra 0.9	**	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Cn	Uut	Fl	Uup	Lv	Uus		Uuo
			*	La 1.1	Ce 1.12	Pr 1.13	Nd 1.14	Pm 1.13	Sm 1.17	Eu 1.2	Gd 1.2	Tb 1.1	Dy 1.22	Ho 1.23	Er 1.24	Tm 1.25	Yb 1.1		Lu 1.27
			**	Ac 1.1	Th 1.3	Pa 1.5	U 1.38	Np 1.36	Pu 1.28	Am 1.13	Cm 1.28	Bk 1.3	Cf 1.3	Es 1.3	Fm 1.3	Md 1.3	No 1.3		Lr 1.3

Metal/metalloid line

Chemical Properties

