




Lecture 11: INEPT, HSQC

Simultaneous spin echo

D




INEPT




$$\hat{\rho}(a) = \frac{1}{2} \mathcal{I}_t + \frac{1}{2} \kappa_1 \mathcal{I}_z + \frac{1}{2} \kappa_2 \mathcal{I}_z$$

INEPT




$$\hat{\rho}(b) = \frac{1}{2} \mathcal{I}_t - \frac{1}{2} \kappa_1 \mathcal{I}_y + \frac{1}{2} \kappa_2 \mathcal{I}_z$$

INEPT




$$\hat{\rho}(e) = \frac{1}{2} \mathcal{I}_t + \frac{1}{2} \kappa_1 \cos \frac{\pi J}{2J} \mathcal{I}_y - \frac{1}{2} \kappa_1 \sin \frac{\pi J}{2J} (2 \mathcal{I}_x \mathcal{I}_z) - \frac{1}{2} \kappa_2 \mathcal{I}_z$$

INEPT





$$\hat{\rho}(e) = \frac{1}{2} \mathcal{I}_t - \frac{1}{2} \kappa_1 (2 \mathcal{I}_x \mathcal{I}_z) - \frac{1}{2} \kappa_2 \mathcal{I}_z$$


INEPT



$$\hat{\rho}(f) = \frac{1}{2} \mathcal{I}_t + \frac{1}{2} \kappa_1 (2 \mathcal{I}_z \mathcal{I}_z) - \frac{1}{2} \kappa_2 \mathcal{I}_z$$


C

INEPT



$$\hat{\rho}(g) = \frac{1}{2} \mathcal{I}_t - \frac{1}{2} \kappa_1 (2 \mathcal{I}_z \mathcal{I}_y) + \frac{1}{2} \kappa_2 \mathcal{I}_y$$

INEPT



$$\hat{\rho}(g) = \frac{1}{2} \mathcal{I}_t - \frac{1}{2} \kappa_1 (2 \mathcal{I}_z \mathcal{I}_y) + \frac{1}{2} \kappa_2 \mathcal{I}_y$$

INEPT

$$\begin{array}{l}
 \mathcal{I}_t \longrightarrow \mathcal{I}_t \longrightarrow \mathcal{I}_t \\
 -2\mathcal{I}_z\mathcal{I}_y \longrightarrow \left\{ \begin{array}{l} -c_2 2\mathcal{I}_z\mathcal{I}_y \longrightarrow \left\{ \begin{array}{l} -c_2c_J 2\mathcal{I}_z\mathcal{I}_y \\ +c_2s_J \mathcal{I}_x \end{array} \right. \\ +s_2 2\mathcal{I}_x\mathcal{I}_z \longrightarrow \left\{ \begin{array}{l} +s_2c_J 2\mathcal{I}_z\mathcal{I}_x \\ +s_2s_J \mathcal{I}_y \end{array} \right. \end{array} \right. \\
 \mathcal{I}_y \longrightarrow \left\{ \begin{array}{l} +c_2\mathcal{I}_y \longrightarrow \left\{ \begin{array}{l} +c_2c_J \mathcal{I}_y \\ -c_2s_J 2\mathcal{I}_x\mathcal{I}_z \end{array} \right. \\ -s_2\mathcal{I}_x \longrightarrow \left\{ \begin{array}{l} -s_2c_J \mathcal{I}_x \\ -s_2s_J 2\mathcal{I}_y\mathcal{I}_z \end{array} \right. \end{array} \right.
 \end{array}$$

Relaxation with J -coupling

- \hat{H}_J : $\mathcal{I}_{1x} \rightarrow 2\mathcal{I}_{1y}\mathcal{I}_{2z}$ $\mathcal{I}_{1y} \rightarrow -2\mathcal{I}_{1x}\mathcal{I}_{2z}$
 $\Rightarrow \mathcal{I}_{1+} = \mathcal{I}_{1x} + i\mathcal{I}_{1y} \rightarrow -i2\mathcal{I}_{1+}\mathcal{I}_{2z}$ different R_2
- $\mathcal{I}_{1+} \leftrightarrow 2\mathcal{I}_{1+}\mathcal{I}_{2z} \Rightarrow \bar{R}_2$
- relaxation of \mathcal{I}_{1+} depends on $2\mathcal{I}_{1+}\mathcal{I}_{2z}$
relaxation of $2\mathcal{I}_{1+}\mathcal{I}_{2z}$ depends on \mathcal{I}_{1+}
cross-correlated cross-relaxation (ingnored here)
cf. cross-relaxation of $\Delta\langle M_{1z} \rangle$ and $\Delta\langle M_{2z} \rangle$ (NOE)



INEPT

$$\langle M_+ \rangle \propto \frac{\kappa_1}{4} e^{-\bar{R}_2 t} \left(e^{-i(\Omega_2 - \pi J)t} - e^{-i(\Omega_2 + \pi J)t} \right) + \frac{\kappa_2}{4} e^{-\bar{R}_2 t} \left(e^{-i(\Omega_2 - \pi J)t} + e^{-i(\Omega_2 + \pi J)t} \right)$$


$$\Re\{Y(\omega)\} =$$

$$\frac{\mathcal{N} \gamma_1^2 \hbar^2 B_0}{16k_B T} \left(\frac{\bar{R}_2}{\bar{R}_2^2 + (\omega - \Omega_2 + \pi J)^2} - \frac{\bar{R}_2}{\bar{R}_2^2 + (\omega - \Omega_2 - \pi J)^2} \right) + \frac{\mathcal{N} \gamma_2^2 \hbar^2 B_0}{16k_B T} \left(\frac{\bar{R}_2}{\bar{R}_2^2 + (\omega - \Omega_2 + \pi J)^2} + \frac{\bar{R}_2}{\bar{R}_2^2 + (\omega - \Omega_2 - \pi J)^2} \right)$$


INEPT




INEPT:



anti-phase vs. in phase coherences



Phase cycling




$$\phi = +90^\circ, \quad y : \quad \hat{\rho}(g) = \frac{1}{2} \mathcal{I}_t - \frac{1}{2} \kappa_1 (2 \mathcal{I}_z \mathcal{I}_y) + \frac{1}{2} \kappa_2 \mathcal{I}_y$$

$$\phi = -90^\circ, \quad -y : \quad \hat{\rho}(g) = \frac{1}{2} \mathcal{I}_t + \frac{1}{2} \kappa_1 (2 \mathcal{I}_z \mathcal{I}_y) + \frac{1}{2} \kappa_2 \mathcal{I}_y$$

$$\text{difference :} \quad \hat{\rho}(g) = - \kappa_1 (2 \mathcal{I}_z \mathcal{I}_y)$$


Phase cycling





$$\Re\{Y(\omega)\} =$$

$$\frac{\mathcal{N}\gamma_1^2\hbar^2 B_0}{16k_B T} \left(\frac{\bar{R}_2}{\bar{R}_2^2 + (\omega - \Omega_2 + \pi J)^2} - \frac{\bar{R}_2}{\bar{R}_2^2 + (\omega - \Omega_2 - \pi J)^2} \right)$$

INEPT with phase cycle:



INEPT vs. direct excitation



INEPT vs. direct excitation

INEPT (phase cycled): $\Re\{Y(\omega)\} =$

$$\frac{\mathcal{N}\gamma_1^2\hbar^2 B_0}{16k_B T} \left(\frac{\bar{R}_2}{\bar{R}_2^2 + (\omega - \Omega_2 + \pi J)^2} - \frac{\bar{R}_2}{\bar{R}_2^2 + (\omega - \Omega_2 - \pi J)^2} \right)$$

Direct excitation: $\Re\{Y(\omega)\} =$


$$\frac{\mathcal{N}\gamma_2^2\hbar^2 B_0}{16k_B T} \left(\frac{\bar{R}_2}{\bar{R}_2^2 + (\omega - \Omega_2 + \pi J)^2} + \frac{\bar{R}_2}{\bar{R}_2^2 + (\omega - \Omega_2 - \pi J)^2} \right)$$

$$\gamma_1^2/\gamma_2^2 \approx \mathbf{16} \text{ for } ^{13}\text{C}$$


$$\gamma_1^2/\gamma_2^2 \approx \mathbf{100} \text{ for } ^{15}\text{N}$$

Insensitive **N**uclei **E**nhanced by **P**olarization **T**ransfer

INEPT vs. direct excitation:



HSQC Spectroscopy (Heteronuclear Single-Quantum Coherence)



COMPLEX EXPERIMENT

ANALYSIS FACILITATED BY SIMPLIFICATIONS

Using results of already analyzed building blocks (echoes)

Ignoring components of $\hat{\rho}$ that cannot produce signal

HSQC Spectroscopy

Measured quantity: M_{1+}

(M_{2+} does not pass the frequency filters)

Only $\mathcal{I}_x \hat{M}_{1+}$ and $\mathcal{I}_y \hat{M}_{1+}$ have non-zero traces:

$$\begin{aligned}\text{Tr} \left\{ \mathcal{I}_x (\mathcal{I}_{1x} + i\mathcal{I}_{1y}) \right\} &= 1 \\ \text{Tr} \left\{ \mathcal{I}_y (\mathcal{I}_{1x} + i\mathcal{I}_{1y}) \right\} &= i\end{aligned}$$

Directly measurable: $\mathcal{I}_x, \mathcal{I}_y$ (in-phase single-quantum of nucleus 1)

Evolve to measurable due to J coupling:

$2\mathcal{I}_x\mathcal{I}_z, 2\mathcal{I}_y\mathcal{I}_z$ (anti-phase single-quantum of nucleus 1)

Need 90° pulse + J coupling:


\mathcal{I}_z (90° pulse), $\mathcal{I}_z, 2\mathcal{I}_z\mathcal{I}_z$ (populations, longitudinal polarization)

$\mathcal{I}_x, \mathcal{I}_y, 2\mathcal{I}_z\mathcal{I}_x, 2\mathcal{I}_z\mathcal{I}_y$ (single-quantum of nucleus 2)

$2\mathcal{I}_x\mathcal{I}_x, 2\mathcal{I}_y\mathcal{I}_y, 2\mathcal{I}_x\mathcal{I}_y, 2\mathcal{I}_y\mathcal{I}_x$ (multiple-quantum)

Never measurable: \mathcal{I}_t (unit matrix)

HSQC Spectroscopy




BLOCK 1: INEPT

$$\hat{\rho}(a) = \frac{1}{2}\mathcal{I}_t + \frac{1}{2}\kappa_1 (\mathcal{I}_z) + \frac{1}{2}\kappa_2 \mathcal{I}_z \rightarrow$$

$$\hat{\rho}(e) = \frac{1}{2}\mathcal{I}_t - \frac{1}{2}\kappa_1 (2\mathcal{I}_z \mathcal{I}_y) + \frac{1}{2}\kappa_2 \mathcal{I}_y$$

HSQC Spectroscopy




BLOCK 2: DECOUPLING ECHO, INCREMENTED t_1

$$\hat{\rho}(e) = \frac{1}{2}\mathcal{I}_t - \frac{1}{2}\kappa_1 (2\mathcal{I}_z\mathcal{I}_y) + \frac{1}{2}\kappa_2\mathcal{I}_y \rightarrow$$

$$\hat{\rho}(f) = \frac{1}{2}\mathcal{I}_t + \frac{1}{2}\kappa_1 (c_{21}2\mathcal{I}_z\mathcal{I}_y - s_{21}2\mathcal{I}_z\mathcal{I}_x) + \frac{1}{2}\kappa_2 (c_{21}\mathcal{I}_y - s_{21}\mathcal{I}_x)$$

HSQC Spectroscopy – Real




BLOCK 3: TWO 90° PULSES, PHASE x (^{13}C or ^{15}N)

$$\hat{\rho}(f) = \frac{1}{2}\mathcal{I}_t + \frac{1}{2}\kappa_1 (c_{21}2\mathcal{I}_z\mathcal{I}_y - s_{21}2\mathcal{I}_z\mathcal{I}_x) + \frac{1}{2}\kappa_2 (c_{21}\mathcal{I}_y - s_{21}\mathcal{I}_x)$$

$$\hat{\rho}(g) = \frac{1}{2}\mathcal{I}_t - \frac{1}{2}\kappa_1 (c_{21}2\mathcal{I}_y\mathcal{I}_z - s_{21}2\mathcal{I}_y\mathcal{I}_x) + \frac{1}{2}\kappa_2 (c_{21}\mathcal{I}_z - s_{21}\mathcal{I}_x)$$

HSQC Spectroscopy – Real




BLOCK 3: TWO 90° PULSES, PHASE x (^{13}C or ^{15}N)

$$\hat{\rho}(f) = \frac{1}{2}\mathcal{I}_t + \frac{1}{2}\kappa_1 (c_{21}2\mathcal{I}_z\mathcal{I}_y - s_{21}2\mathcal{I}_z\mathcal{I}_x) + \frac{1}{2}\kappa_2 (c_{21}\mathcal{I}_y - s_{21}\mathcal{I}_x)$$

$$\hat{\rho}(g) = -\frac{1}{2}\kappa_1 c_{21}2\mathcal{I}_y\mathcal{I}_z + \text{unmeasurable (no more } 90^\circ \text{ pulses)}$$

HSQC Spectroscopy – Real




BLOCK 4: SIMULTANEOUS ECHO

$$\hat{\rho}(g) = -\frac{1}{2}\kappa_1 c_{21} 2\mathcal{I}_y \mathcal{I}_z + \text{unmeasurable} \rightarrow$$


$$\hat{\rho}(h) = \frac{1}{2}\kappa_1 c_{21} \mathcal{I}_x + \text{unmeasurable}$$

HSQC Spectroscopy – Real



$$\frac{1}{2} \kappa_1 c_{21} \mathcal{I}_x \rightarrow \begin{cases} \frac{1}{2} \kappa_1 c_{21} c_{12} \mathcal{I}_x \\ \frac{1}{2} \kappa_1 c_{21} s_{12} \mathcal{I}_y \end{cases} \rightarrow \begin{cases} +\frac{1}{2} \kappa_1 c_{21} c_{12} c_J \mathcal{I}_x \\ +\frac{1}{2} \kappa_1 c_{21} c_{12} s_J 2\mathcal{I}_y \mathcal{I}_z \\ +\frac{1}{2} \kappa_1 c_{21} s_{12} c_J \mathcal{I}_y \\ -\frac{1}{2} \kappa_1 c_{21} s_{12} s_J 2\mathcal{I}_x \mathcal{I}_z \end{cases}$$

HSQC Spectroscopy – Imaginary




BLOCK 3: TWO 90° PULSES, PHASE y (^{13}C or ^{15}N)

$$\hat{\rho}(f) = \frac{1}{2}\mathcal{I}_t + \frac{1}{2}\kappa_1 (c_{21}2\mathcal{I}_z\mathcal{I}_y - s_{21}2\mathcal{I}_z\mathcal{I}_x) + \frac{1}{2}\kappa_2 (c_{21}\mathcal{I}_y - s_{21}\mathcal{I}_x)$$

$$\hat{\rho}(g) = \frac{1}{2}\mathcal{I}_t - \frac{1}{2}\kappa_1 (c_{21}2\mathcal{I}_y\mathcal{I}_y + s_{21}2\mathcal{I}_y\mathcal{I}_z) + \frac{1}{2}\kappa_2 (c_{21}\mathcal{I}_y + s_{21}\mathcal{I}_z)$$

HSQC Spectroscopy – Imaginary




BLOCK 3: TWO 90° PULSES, PHASE y (^{13}C or ^{15}N)

$$\hat{\rho}(f) = \frac{1}{2}\mathcal{I}t + \frac{1}{2}\kappa_1 (s_{21}2\mathcal{I}_z\mathcal{I}_y - s_{21}2\mathcal{I}_z\mathcal{I}_x) + \frac{1}{2}\kappa_2 (c_{21}\mathcal{I}_y - s_{21}\mathcal{I}_x)$$

$$\hat{\rho}(g) = -\frac{1}{2}\kappa_1 s_{21}2\mathcal{I}_y\mathcal{I}_z + \text{unmeasurable (no more } 90^\circ \text{ pulses)}$$

HSQC Spectroscopy – Imaginary




BLOCK 4: SIMULTANEOUS ECHO

$$\hat{\rho}(g) = -\frac{1}{2}\kappa_1 s_{21} 2\mathcal{I}_y \mathcal{I}_z + \text{unmeasurable} \rightarrow$$


$$\hat{\rho}(h) = \frac{1}{2}\kappa_1 s_{21} \mathcal{I}_x + \text{unmeasurable}$$

HSQC Spectroscopy – Imaginary




$$\frac{1}{2}\kappa_1 s_{21} \mathcal{I}_x \rightarrow \begin{cases} \frac{1}{2}\kappa_1 s_{21} c_{12} \mathcal{I}_x \\ \frac{1}{2}\kappa_1 s_{21} s_{12} \mathcal{I}_y \end{cases} \rightarrow \begin{cases} +\frac{1}{2}\kappa_1 s_{21} c_{12} c_J \mathcal{I}_x \\ +\frac{1}{2}\kappa_1 s_{21} c_{12} s_J 2\mathcal{I}_y \mathcal{I}_z \\ +\frac{1}{2}\kappa_1 s_{21} s_{12} c_J \mathcal{I}_y \\ -\frac{1}{2}\kappa_1 s_{21} s_{12} s_J 2\mathcal{I}_x \mathcal{I}_z \end{cases}$$

HSQC Spectroscopy – Hypercomplex




$$\frac{1}{2}\kappa_1 e^{i\Omega_2 t_1} \mathcal{I}_x \rightarrow \begin{cases} \frac{1}{2}\kappa_1 e^{i\Omega_2 t_1} c_{12} \mathcal{I}_x \\ \frac{1}{2}\kappa_1 e^{i\Omega_2 t_1} s_{12} \mathcal{I}_y \end{cases} \rightarrow \begin{cases} +\frac{1}{2}\kappa_1 e^{i\Omega_2 t_1} c_{12} c_J \mathcal{I}_x \\ +\frac{1}{2}\kappa_1 e^{i\Omega_2 t_1} c_{12} s_J 2\mathcal{I}_y \mathcal{I}_z \\ +\frac{1}{2}\kappa_1 e^{i\Omega_2 t_1} s_{12} c_J \mathcal{I}_y \\ -\frac{1}{2}\kappa_1 e^{i\Omega_2 t_1} s_{12} s_J 2\mathcal{I}_x \mathcal{I}_z \end{cases}$$

HSQC Spectroscopy – Hypercomplex




$$\Re\{Y(\omega)\} = \frac{\mathcal{N}\gamma_1^2\hbar^2 B_0}{16k_B T} \times \frac{\overline{R}_{2,2}^2}{\overline{R}_{2,2}^2 + (\omega - \Omega_2)^2} \left(\frac{\overline{R}_{2,1}^2}{\overline{R}_{2,1}^2 + (\omega - \Omega_1 + \pi J)^2} + \frac{\overline{R}_{2,1}^2}{\overline{R}_{2,1}^2 + (\omega - \Omega_1 - \pi J)^2} \right)$$

Decoupling in direct dimension




Decoupling in direct dimension





$$\Re\{Y(\omega)\} = \frac{\mathcal{N}\gamma_1^2\hbar^2 B_0}{8k_B T} \frac{\overline{R}_{2,2}^2}{\overline{R}_{2,2}^2 + (\omega - \Omega_2)^2} \frac{R_{2,1}^2}{R_{2,1}^2 + (\omega - \Omega_1)^2}$$

Decoupling in direct dimension




$$\Re\{Y(\omega)\} = \frac{\mathcal{N}\gamma_1^2\hbar^2 B_0}{8k_B T} \frac{\overline{R}_{2,2}^2}{\overline{R}_{2,2}^2 + (\omega - \Omega_2)^2} \frac{R_{2,1}^2}{R_{2,1}^2 + (\omega - \Omega_1)^2}$$

Decoupling in direct dimension



HSQC spectrum of a 20 kDa protein



Benefits of HSQC

- *High sensitivity* for ^{13}C or ^{15}N
(higher by $(\gamma_1/\gamma_2)^{5/2}$ than by the direct detection)
- *High resolution*
Second dimension and less peaks in spectrum
(only $^{13}\text{C}/^{15}\text{N}$ -bonded protons and protonated $^{13}\text{C}/^{15}\text{N}$ visible)
- *Important structural information*
 ^1H - ^{13}C and ^1H - ^{15}N correlation
(it tells us which proton is attached to which ^{13}C or ^{15}N).

HOMework:

COSY

Section 11.3

