

C8953
NMR strukturní analýza
seminář

Identification of an unknown compound

Jan Novotný
176003@is.muni.cz

April 1, 2020

Task 0: Classification of an unknown substance

Assign the general name to displayed substances:

CARBOHYDRATE

PEPTIDE


STEROID

TERPENE


ALKALOID

COUMARINE


VISNADINE




QUININE




CARVONE




SACHAROSA




DIHYDROCHOLESTEROL




SGGLRLHLGLS



Task 0: Classification of an unknown substance



Task 0: Classification of an unknown substance




Task 1: $C_{10}H_{13}NO$

δ [ppm]	Multiplicity	Integral
1.05	triplet	3
1.75	singlet	3
3.70	quartet	2
7-7.60	complex multiplet	5

Task 1: C₁₀H₁₃NO

δ [ppm]	Multiplicity	Integral
1.05 a	triplet	3
1.75 c	singlet	3
3.70 b	quartet	2
7-7.60 d-f	complex multiplet	5




Task 2: $C_{11}H_{15}NO_2$

δ [ppm]	Multiplicity	J (Hz)	Integral
1.30	triplet	7	3
3.00	singlet	-	6
4.25	quartet	7	2
6.65	dublet	8	2
7.80	dublet	8	2


Task 2: $C_{11}H_{15}NO_2$

δ [ppm]	Multiplicity	J (Hz)	Integral
1.30 b	triplet	7	3
3.00 e	singlet	-	6
4.25 a	quartet	7	2
6.65 d	dublet	8	2
7.80 c	dublet	8	2



Task 3: $C_{12}H_{17}NO$ - $^1H/COSY$


1H NMR (300 MHz, $CDCl_3$)



SOLUTION


Task 3: $C_{12}H_{17}NO$ - $^1H/COSY$

COSY (300 MHz, $CDCl_3$)




Task 3: $C_{12}H_{17}NO$ - 1H - ^{13}C /HSQC, HMBC

HSQC (300 MHz, $CDCl_3$)




Task 3: $C_{12}H_{17}NO$ - 1H - ^{13}C /HSQC, HMBC



HMBC (300 MHz, $CDCl_3$)



^1H NMR (300 MHz, CDCl_3)



COSY (300 MHz, CDCl_3)




General comments

- inspect molecular formula $C_mH_hO_oN_nX_x$:
Degree of unsaturation $m + 1 - 0.5(h + x - n)$
- identify signals of CH_3 and exchangeable protons in 1D 1H spectrum
- arbitrary numbering (e.g., from lower to higher value of chemical shift) of resolved resonances in all spectra
- identification of the individual spin systems using DQF-COSY
- resolve geminal protons using HSQC
- connect molecular fragments/isolated spins using HMBC, NOESY
- specify the stereochemistry (relative configuration) by means of J - and NOE interaction

- in 1D spectrum bottom blue numbers are integrals, labels in violet frames contains the arbitrary label (A-N), multiplet specification (use with caution, automatically determined), and position of a signal in ppm
- UnHa-UnHb in 2D refers to correlation of protons a and b of unknown compound Un

1D ^1H of $\text{C}_{10}\text{H}_{20}\text{O}$



1H 1D

c=0.8M DMSO 25C TXI





06 1.04 1.02 1.00 0.98 0.96 0.94 0.92 0.90 0.88 0.86 0.84 0.82 0.80 0.78 0.76 0.74 0.72 0.70 0.68


f1 (ppm)

Navigation icons: back, forward, search, zoom, etc.


^1H - ^{13}C HSQC and NOESY



Task 1: J -connectivity of $C_{10}H_{20}O$





Task 1: J -connectivity of $C_{10}H_{20}O$




Task 1: J -connectivity of $C_{10}H_{20}O$

- CH **9g** has crosspeaks with deshielded **10m** \Rightarrow OH group (**n**)
- CH **9g** has two crosspeaks with diastereotopic protons **4if**





Task 1: J -connectivity of $C_{10}H_{20}O$




Task 1: J -connectivity of $C_{10}H_{20}O$

- CH **10m** connected with CH₂ **8ck**
- CH₂ **4if** connected with CH₂ **7bj**





Task 1: J -connectivity of $C_{10}H_{20}O$




Task 1: J -connectivity of $C_{10}H_{20}O$

- CH_2 **8ck** weakly coupled with CH_2 **7bj \Rightarrow closing ring**
- protons **b** and **k** coupled to CH **6h** which is connected to methyl **3e**
- other expected crosspeaks in DQ-COSY crowded/overlapped, found topology confirmed in HMBC ($^3/4 J_{HC}$)





Task 1: J -connectivity of $C_{10}H_{20}O$



Task 1: J -connectivity of $C_{10}H_{20}O$



- CH_2 **8ck** weakly coupled with CH_2 **7bj \Rightarrow closing ring**
- protons **b** and **k** coupled to CH **6h** which is connected to methyl **3e**
- other expected crosspeaks in DQ-COSY crowded/overlapped, found topology confirmed in HMBC ($^{3/4}J_{HC}$)



Task 1:

Stereochemistry of menthol $C_{10}H_{20}O$

- $1 \leftrightarrow 2$:
homonuclear/heteronuclear couplings
 - large couplings preserved in 1D slices of HSQC:
axial H - 2 visible interactions (geminal and vicinal) \times
equatorial H - only geminal
 - 1D TOCSY: selective decoupling \Rightarrow simplification of complex multiplets
 - DQF-COSY: analysis of phase sensitive spectrum
- $1 \leftrightarrow 3$: NOE contacts (axial strong)



Task 1:

Stereochemistry of menthol $C_{10}H_{20}O$

- $1 \leftrightarrow 2$:
homonuclear/heteronuclear couplings
 - large couplings preserved in 1D slices of HSQC:
axial H - 2 visible interactions (geminal and vicinal) \times
equatorial H - only geminal
 - 1D TOCSY: selective decoupling \Rightarrow simplification of complex multiplets
 - DQF-COSY: analysis of phase sensitive spectrum
- $1 \leftrightarrow 3$: NOE contacts (axial strong)

