



Orion Nebula, Distance about 450 pc, Total Mass about 5000  $M(\text{sun})$ , Diameter about 3 pc

M11, NGC 6705: Total Mass About 10000  $M(\text{sun})$ , 200 Myr



# Cluster formation

- Observations versus Models
- Important parameters
  1. Time scale
  2. Total mass
  3. Initial Mass Function
  4. Velocity distribution
  5. Binary fraction
  6. Diameter
  7. Density distribution


# Heuristic Approach

- We know of 14 Open Clusters which are younger than 10 Myrs within 1000 pc around the Sun (Source: WEBDA)
- There are also five star forming regions
- Open Clusters still have to form within the solar vicinity
- Total masses: up to 40 000 M(sun)
- Stable for some Gyrs
- Evolutionary theory has to explain these facts


## Clusters selected

Cluster_name	RA_2000_Dec		l	b	Dist	Mod	EB-V	Age	ST	Z	Diam	Fe/H	MRV	pm RA	pm Dec
<a href="#">Mamajek 1</a>	08 42 06	-79 01 38	292.482	-21.654	97	4.93	0.00	6.9		-35.8	40.0		+16.1	-30.00	+27.80
<a href="#">Collinder 70</a>	05 35 31	-01 06 00	205.03	-17.35	391	8.09	0.04	6.71		-116.6	180.0		19.49	0.36	-0.68
<a href="#">ASCC 24</a>	06 28 44	-07 01 11	216.64	-8.23	400	8.44	0.14	6.96		-57.3	42.0		16.35	-5.55	-4.05
<a href="#">ASCC 16</a>	05 24 35	+01 47 59	200.98	-18.35	460	8.59	0.09	6.93		-144.8	74.4		0.75	+0.75	-0.18
<a href="#">NGC 1980</a>	05 35 24	-05 54 35	209.51	-19.60	550	8.86	0.05	6.67	B1	-184.5	25.2		25.34	0.83	-0.36
<a href="#">Bochum 14</a>	18 02 00	-23 41 00	6.388	-0.499	578	13.48	1.508	6.996		-5.0	2.0				
<a href="#">NGC 2264</a>	06 40 58	+09 53 42	202.936	2.196	667	9.28	0.051	6.954	O7	25.6	39.0	-0.15	+25.5	-1.13	-3.80
<a href="#">ASCC 122</a>	22 33 14	+39 36 36	95.91	-15.90	700	9.53	0.10	6.98		-191.8	86.4		-8.17	-0.29	-4.19
<a href="#">Collinder 419</a>	20 17 59	+40 43 12	78.07	2.79	740	10.40	0.34	6.85	B2	36.0	30.0		-8.19	-2.56	-6.99
<a href="#">ASCC 79</a>	15 19 11	-60 43 47	320.04	-2.86	800	10.01	0.16	6.86		-39.9	62.4		4.03	-2.67	-4.10
<a href="#">IC 5146</a>	21 53 24	+47 16 00	94.383	-5.495	852	11.49	0.593	6.00	B1	-81.6	20.0			-1.77	-1.70
<a href="#">Lynga 14</a>	16 55 04	-45 14 00	340.919	-1.089	881	14.15	1.428	6.712		-16.7	3.0				
<a href="#">Ruprecht 119</a>	16 28 15	-51 30 00	333.276	-1.879	956	11.67	0.570	6.853		-31.3	8.0			-1.15	-1.80
<a href="#">NGC 6383</a>	17 34 48	-32 34 00	355.690	0.041	985	10.89	0.298	6.962	O7	0.7	20.0		+7.00	+1.58	-2.00

Several spurious entries like the “ASCC clusters”



Distribution of young open clusters and star forming regions from Alfaro et al., 2009, Ap&SS, 324, 141



Stars hotter  
than B0 and  
B0 to B2


Distribution of star forming regions from Preibisch & Mamajek, 2008, Handbook of Star Forming Regions, Volume II

# Giant Molecular Clouds

- Star Clusters can only form within „Giant Molecular Clouds“ (GMC) with a high enough initial mass
- The stellar formation rate in the solar neighborhood is very low
- But still there have to exist several GMCs to form Star Clusters
- Is the formation process the same for all observed Galaxy types?

# Giant Molecular Clouds

Stark & Lee, 2006, ApJ, 641, L116



Recent investigation of the  $^{13}\text{CO}$  Gas within 2000 pc around the Sun

The number of young OCLs can be very well explained

Formation rate of 0.45 OCLs per  $\text{kpc}^{-2} \text{Myr}^{-1}$  in the galactic disk within 2 kpc around the Sun

Battinelli & Capuzzo-Dolcetta, 1991, MNRAS, 248, 76




NGC 6611 (M16)

$d = 1750 \text{ pc}$

$t = 8 \text{ Myr}$

Star formation  
„live“



**Gaseous Pillars • M16**

**HST • WFPC2**

PRC95-44a • ST Sci OPO • November 2, 1995  
J. Hester and P. Scowen (AZ State Univ.), NASA

# Initial Mass Function

- The „Initial Mass Function“ (IMF) describes the mass distribution for a population of stars when they are formed together
- Relevant astrophysics:
  1. Size, total mass and metallicity of the initial GMC
  2. Fragmentation of the GMC
  3. Conservation of the angular momentum
  4. Local and global magnetic fields
  5. Accretion in the Pre-Main Sequence phase
- The **only** observational parameter for the test of stellar formation and evolution models
- We observe a luminosity function which has to be transformed to the IMF

# Initial Mass Function

- Several most important questions are still not solved
  1. Is the IMF homogeneous within the Milky Way?
  2. Is the IMF constant throughout time?
  3. What is the influence of the local and global magnetic field on the IMF?
  4. What is the influence of the local and global metallicity on the IMF?

# Initial Mass Function


The IMF  $\theta(m)$ , often called „Present-Day Mass Function“ (PDMF), is defined as:

$$dN = \theta(m) dm$$

$dN$  is the number of all stars per cubic parsec on the *main sequence* with a mass between  $M$  and  $(M + dm)$ .

But we observe not the masses of stars but their magnitudes (relative and absolute) or luminosities.

So we have to define the luminosity function and transform it into the IMF.



In each row ( $M_V + dM$ ) there is a mixture of main sequence and evolved objects. For the IMF, we need the main sequence only.

# Luminosity function

The luminosity function  $\Psi(M_V)$ , is defined as:

$$dN = -\Psi(M_V) dM_V$$


$dN$  is the number of all stars per cubic parsec on the *main sequence* with an absolute magnitude between  $M_V$  and  $(M_V + dM_V)$ .

The transformation to the IMF is given as:

$$\theta(m) = -\Psi(M_V)[dm(M_V)/dM_V]^{-1}$$

The second term is the derivation of the Mass-Luminosity function  $m(M_V)$ . It is depending on the age ( $t$ ), metallicity ( $Z$ ) and rotation ( $v_{\text{rot}}$ )

$$m(M_V) = m(M_V, Z, t, v_{\text{rot}})$$



# Correction of the observations

We have to correct the complete observations for the evolved objects. There are three possibilities:

1. Take a statistical sample with a well known luminosity function (clusters)
2. Take a statistical sample with well known photometric magnitudes and distances
3. Take isochrones = theoretical star evolution = models based on observations = circular argument

All these methods are not self consistent and always introduce an unknown error to the analysis



FRACTION  $f$  OF MAIN-SEQUENCE STARS (TYPE EARLIER THAN  $Sp_d$ )

	$M_v$								
	-4.5	-3.5	-2.5	-1.5	-0.5	+0.5	+1.5	+2.5	+3.5
$Sp_d$ .....	B0	B3	B6	B9	A1	A6	F0	F8	G7
$f$ .....	0.10	0.25	0.48	0.51	0.43	0.40	0.60	0.70	0.90

Salpeter, 1955, ApJ, 121, 161

Results of classical spectral classification, only 10% of stars with  $M_v = -4.5$  mag are on the main sequence!

These values are depending on the chosen sample for the spectral classification and which classification scheme is applied.

The errors are rather large.


All observations have to be normalized to one “standard system” which means essentially to one “time scale”.

The observations show, that this heuristic law describes them very well

$$\theta(m) \approx m^{-\Gamma} \quad \text{Salpeter law (1955)}$$

Star cluster are one of the most important observational test for the IMF because they, normally, have well defined ages, distances and metallicities. However, the errors are still quiet large.


But there is still no homogeneous IMF determination for open clusters taking into account the available data.



# TYCHO2 data

cluster	$(m - M)_0$ [mag]	$E_{B-V}$ [mag]	$t$ Myr	$d$ [']
Blanco 1*	6.8	0.03	50	105
Stock 2	7.5	...	100	260
$\alpha$ Per*	6.3	0.09	20	255
Pleiades*	5.6	0.05	75	300
NGC 2451 A*	6.4	0.00	20	140
IC 2391*	5.8	0.00	20	110
Praesepe*	6.0	0.00	650	195
IC 2602*	5.8	0.03	10	185
NGC 7092	7.6	0.12	70	170

cluster	# stars	$\Gamma$	mass range [ $M_\odot$ ]	$V_T$ range [mag]
Blanco 1	34	$-2.27 \pm 0.70$	[1.1; 4.8]	[6.1; 11.4]
Stock 2	204	$-2.01 \pm 0.40$	[1.5; 4.1]	[7.6; 11.0]
$\alpha$ Per	70	$-1.57 \pm 0.44$	[1.1; 6.8]	[5.0; 10.5]
Pleiades	127	$-1.99 \pm 0.39$	[1.0; 4.1]	[5.0; 10.9]
NGC 2451 A	27	$-0.69 \pm 0.63$	[1.3; 6.8]	[4.8; 10.0]
IC 2391	29	$-1.07 \pm 0.53$	[1.1; 8.1]	[3.5; 10.7]
NGC 7092	25	$-1.93 \pm 1.24$	[1.4; 3.4]	[6.5; 9.9]




Mass-Function Slope  $\Gamma$  for Two Subregions and for the Whole-Cluster Region  
in the Given Mass Range

Cluster	Mass range ( $M_{\odot}$ )	Mass function slopes ( $\Gamma \pm \sigma$ )		
		Inner region	Outer region	Whole cluster
Be 62	11.17–1.14	$-0.89 \pm 0.17$	$-2.10 \pm 0.74$	$-1.88 \pm 0.34$
NGC 1528	2.55–0.73	$-1.96 \pm 0.42$	$-2.17 \pm 0.43$	$-2.10 \pm 0.35$
NGC 1960	6.82–1.01	$-1.25 \pm 0.24$	$-1.99 \pm 0.15$	$-1.80 \pm 0.14$
NGC 2287	2.70–0.83	$-1.35 \pm 0.86$	$-1.22 \pm 0.27$	$-1.22 \pm 0.19$
NGC 2301	2.78–0.82	$-0.85 \pm 0.33$	$-1.56 \pm 0.54$	$-1.34 \pm 0.32$
NGC 2323	4.22–0.67	$-1.69 \pm 0.09$	$-2.28 \pm 0.31$	$-2.01 \pm 0.17$
NGC 2420	1.44–0.67	$-0.93 \pm 0.32$	$-1.50 \pm 0.56$	$-1.30 \pm 0.39$
NGC 2437	3.51–1.02	$-1.72 \pm 0.13$	$-2.30 \pm 0.62$	$-2.03 \pm 0.42$
NGC 2548	2.46–0.82	$-1.11 \pm 0.85$	$-1.02 \pm 0.36$	$-1.12 \pm 0.70$

Typical values and errors


# Magnetic field – star formation

- Price & Bate, 2009, MNRAS, 398, 33
- Effects of magnetic pressure on fragmentation




Increasing magnetic field strength

Increasing magnetic field strength ↓



# Star formation



Gravitation „wins“

Magnetic field, Shock wave

Protostar

← **FREE GAS**



**NO FREE GAS**





# Star formation

- The detection of free Gas in a Star Cluster is an excellent indicator for the time scale of continuous stellar formation

STAR-FORMING REGIONS

Region	$\langle t \rangle^a$ (Myr)	Molecular Gas?	Ref. (age)
Coalsack .....	...	Yes	...
Orion Nebula .....	1	Yes	1
Taurus .....	2	Yes	1, 2, 3
Oph .....	1	Yes	1
Cha I, II .....	2	Yes	1
Lupus .....	2	Yes	1
MBM 12A .....	2	Yes	4
IC 348 .....	1-3	Yes	1, 4, 5, 6
NGC 2264 .....	3	Yes	1
Upper Sco .....	2-5	No	1, 6, 7
Sco OB2 .....	5-15	No	8
TWA .....	~10	No	9
$\eta$ Cha .....	~10	No	10

<sup>a</sup> Average age in Myr.

Star formation lasts  
3 to 4 Myrs and is  
**continuous**

This is also the  
“intrinsic” error of an  
age determination

# Numerical simulation of star formation in Giant Molecular Clouds

- Hypothesis: the formation of all members of a star cluster is continuous for 3 to 4 Myrs within one GMCs
- Is this a realistic approach?
- Is it possible to simulation the formation of star clusters and compare the results with observational data within the solar vicinity?

# Numerical simulation of star formation in Giant Molecular Clouds

- Detailed paper by Bate & Bonnell, 2005, MNRAS, 356, 1201
- Basis: Orion Nebula and Taurus star forming region
- “Complete” astrophysical numerical simulation including Shock Waves, dynamical parameters and 3D-Hydrodynamics, Jeans Mass  $< 1 M(\text{sun})$
- The numerical simulations are astonishing close to the observations

# Numerical simulation of star formation in Giant Molecular Clouds

Input parameter:

1. Mass (GMC) = 50 M(sun), limited by CPU time
2. Diameter = 0.375 pc, limited by CPU time
3. Time for the gravitational collapse: 19000 years
4. Random turbulence field with a 3D Gaussian distribution


Core	Initial Gas Mass $M_{\odot}$	Initial Size pc	Final Gas Mass $M_{\odot}$	No. Stars Formed	No. Brown Dwarfs Formed	Mass of Stars and Brown Dwarfs $M_{\odot}$	Star Formation Efficiency %
1	1.50 (0.15)	$0.04 \times 0.04 \times 0.03$	2.03 (1.04)	$\geq 13$	$\leq 52$	6.33	76 (86)
2	0.92 (0.16)	$(0.03 \times 0.01 \times 0.01)$	1.18 (0.50)	$\geq 4$	$\leq 8$	1.33	53 (73)
3	0.17 (0.06)	$(0.02 \times 0.01 \times 0.01)$	0.32 (0.08)	1	0	0.18	36 (69)
4	0.31 (0.07)	$(0.03 \times 0.01 \times 0.01)$	0.32 (0.06)	1	0	0.09	22 (60)
Cloud	50.0	$0.38 \times 0.38 \times 0.38$	42.1	$\geq 19$	$\leq 60$	7.92	16

„Stars“: Mass  $> 0.084 M(\text{sun})$

Brown Dwarfs: Mass  $< 0.084 M(\text{sun})$ , no Hydrogen burning


More low mass stars formed due to the IMF


For star clusters it is essential to know the internal velocity distribution because of their evolution (see later)



The formation of  
Binary systems

Continuous star formation  
in time





Binaries are  
connected with  
a line

The rms velocity dispersion of the simulations is  $4.3 \text{ km s}^{-1}$   
Such observational data for  $d > 500 \text{ pc}$  are still not  
available => Gaia satellite mission

# Evolution of Star Clusters

- Star Clusters form with the following characteristics
  1. **Total Mass: IMF**
  2. **Metallicity**
  3. **Kinematics of the Cluster center:** location within the Galaxy
  4. **Internal velocity dispersion**
- How does a Star Cluster evolve with these starting parameters?



- Each member (= star) evolve “as an individual”, some important topics
  1. Binary Evolution
  2. Mass Loss (hot stars)
  3. AGB Evolution
  4. Planetary Nebula (cool stars)
  5. Supernovae explosions
- In Star Clusters, collisions are very uncommon (see later), almost no new multiple (binary) systems form during the later evolution
- Star Clusters, normally, follow Galactic Rotation

