

Liam.Keegan@ceitec.muni.cz

Lewin, GENES XI

Chapter 26

Operons, prokaryotic gene regulation and
yeast GAL4

[Lewin's GENES XII \(PDF\)](#)

<https://pdfroom.com/books/lewins-genes-xii/Wx5aDYKI2BJ>

[biotecher.ir](#)

<https://biotecher.ir/wp-content/uploads/2018/06/bio...>

LEWIN'S GENES XI

JOCELYN E. KREBS
ELLIOTT S. GOLDSTEIN
STEPHEN T. KILPATRICK

Chapter 26 The Operon

26.2 Operons are Structural Gene Clusters that Are Coordinately Controlled

- Genes coding for proteins that function in the same pathway may be located adjacent to one another and controlled as a single unit that is transcribed into a **polycistronic mRNA**.





Figure 26.05: The lac operon occupies ~6000 bp of DNA.

Sequence-specific DNA recognition is the key to differential gene regulation.



Mark Ptashne with graduate students Cynthia Wohlberger, Liam Keegan, Ed Giniger at Harvard, 1982

26.1 Introduction

- In **negative regulation**, a repressor protein binds to an **operator** to prevent a gene from being expressed.
- In **positive regulation**, a transcription factor is required to bind at the promoter in order to enable RNA polymerase to initiate transcription.




Figure 26.02: In negative control, a trans-acting repressor binds to the cis-acting operator to turn off transcription.




Figure 26.03: In positive control, a trans-acting factor must bind to cis-acting site in order for RNA polymerase to initiate transcription at the promoter.

26.1 Introduction

- In **inducible regulation**, the gene is regulated by the presence of its substrate (the **inducer**).
- In **repressible regulation**, the gene is regulated by the product of its enzyme pathway (the **corepressor**).

26.3 Famous inducible genes in *E. coli*

The *lac* Operon Is under Negative control by a repressor







Figure 26.06: *lac* repressor and RNA polymerase bind at sites that overlap around the transcription startpoint of the *lac* operon.

- Transcription of the *lacZYA* operon is controlled by a repressor protein (the ***lac* repressor**) that binds to an operator that overlaps the promoter at the start of the cluster.
- **constitutive expression** – A state in which a gene is expressed continuously.
- In the absence of β -galactosides, the *lac* operon is expressed only at a very low (basal) level.

lacZ (β -galactosidase) plate tests and liquid culture assays for beta-gal activity units.

X-gal identifies cells with beta-galactosidase



ONPG hydrolysis by cells gives a soluble blue product for spectrophotometric measurement of *lacZ* (β -galactosidase) activity units

26.3 The *lac* Operon Is under negative transcriptional control by repressor and is inducible by galactosides

- The repressor protein is a tetramer of identical subunits coded by the *lacI* gene.
- β -galactoside sugars, the substrates of the *lac* operon, are its inducer.
- Addition of specific β -galactosides induces transcription of all three genes of the *lac* operon.
- The *lac* mRNA is extremely unstable; as a result, induction can be rapidly reversed.

26.3 The *lac* Operon Is Negative Inducible




Figure 26.07: Addition of inducer results in rapid induction of *lac* mRNA, and is followed after a short lag by synthesis of the enzymes.

26.4 *lac* Repressor Is Controlled by a Small-Molecule Inducer




Figure 26.08: *lac* repressor maintains the *lac* operon in the inactive condition by binding to the operator.

- An inducer functions by converting the repressor protein into a form with lower operator affinity.
- Repressor has two binding sites, one for the operator DNA and another for the inducer.
- **gratuitous inducer** – Inducers that resemble authentic inducers of transcription, but are not substrates for the induced enzymes.

26.4 *lac* Repressor Is Controlled by a Small-Molecule Inducer

- Repressor is inactivated by an allosteric interaction in which binding of inducer at its site changes the properties of the DNA-binding site (**allosteric control**).
- The true inducer is **allolactose**, not the actual substrate of β -galactosidase. In experiments we induce with artificial IPTG, not metabolized, a gratuitous inducer.




Figure 26.09: Addition of inducer converts repressor to a form with low affinity for the operator. This allows RNA polymerase to initiate transcription.

26.5 *cis*-Acting Constitutive Mutations Identify the Operator

- Mutations in the operator cause constitutive expression of all three *lac* structural genes.
- These mutations are *cis*-acting and affect only those genes on the contiguous stretch of DNA.
- Mutations in the promoter prevent expression of *lacZYA* and are **uninducible** and *cis*-acting.

26.5 *cis*-Acting Constitutive Mutations

Identify the Operator

- ***cis*-dominant** – A site or mutation that affects the properties only of its own molecule of DNA, often indicating that a site does not code for a diffusible product. O^C means ‘operator constitutive’ mutation




Figure 26.10: Operator mutations are constitutive because the operator is unable to bind repressor protein.

26.6 *trans*-Acting Mutations Identify the Regulator Gene

- Mutations in the *lacI* gene are *trans*-acting and affect expression of all *lacZYA* clusters in the bacterium.
- Mutations that eliminate *lacI* function cause constitutive expression and are recessive (*lacI⁻*).
- Mutations in the DNA-binding site of the repressor are constitutive because the repressor cannot bind the operator.




Figure 26.11: Mutations that inactivate the *lacI* gene cause the operon to be constitutively expressed.

26.6 *trans*-Acting Mutations Identify the Regulator Gene

- Mutations in the inducer-binding site of the repressor prevent it from being inactivated and cause uninducibility. Super-repressor mutants.
- When mutant and wild-type subunits are present, a single *lacI^d* mutant subunit can inactivate a tetramer whose other subunits are wild-type.
 - It is **dominant negative**.

26.6 *trans*-Acting Mutations Identify the Regulator Gene




Figure 26.12: A *lacI*-d mutant gene makes a monomer that has a damaged DNA binding.

- **negative complementation**
 - This occurs when interallelic complementation allows a mutant subunit to suppress the activity of a wild-type subunit in a multimeric protein.
- *lacI*^{-d} mutations occur in the DNA-binding domain. Their effect is explained by the fact that repressor activity requires all DNA-binding sites in the tetramer to be active.

26.7 *lac* Repressor Is a Tetramer Made of Two Dimers

- A single repressor subunit can be divided into the N-terminal DNA-binding domain, a hinge, and the core of the protein.
- The DNA-binding domain contains two short α -helical regions that bind the major groove of DNA.
- The inducer-binding site and the regions responsible for multimerization are located in the core.

26.7 *lac* Repressor Is a Tetramer Made of Two Dimers





Figure 26.13: The structure of a monomer of Lac repressor identifies several independent domains.

Structure from Protein Data Bank 1LBG. M. Lewis, et al., Science 271 (1996): 1247-1254. Photo courtesy of Hongli Zhan and Kathleen S. Matthews, Rice University.

Copyright © 2013 by Jones & Bartlett Learning, LLC an Ascend Learning Company
www.jblearning.com


26.7 *lac* Repressor Is a Tetramer Made of Two Dimers



- Monomers form a dimer by making contacts between core subdomains 1 and 2.
- Dimers form a tetramer by interactions between the tetramerization helices.

Figure 26.15: The repressor tetramer consists of two dimers.


26.7 *lac* Repressor Is a Tetramer Made of Two Dimers



- Different types of mutations occur in different domains of the repressor protein.

Figure 26.16: The locations of three type of mutations in lactose repressor are mapped on the domain structure of the protein.

26.8 *lac* Repressor Binding to the Operator Is Regulated by an Allosteric Change in Conformation



- Inducer binding causes a change in repressor conformation that reduces its affinity for DNA and releases it from the operator.

Figure 26.18: The inducer changes the structure of the core.

26.8 *lac* Repressor Binding to the Operator Is Regulated by an Allosteric Change in Conformation

- *lac* repressor protein binds to the double-stranded DNA sequence of the operator.
- The operator is a **palindromic** sequence of 26 bp.
- Each inverted repeat of the operator binds to the DNA-binding site of one repressor subunit.




Figure 26.17: The *lac* operator has a symmetrical sequence.

26.9 *lac* Repressor Binds to Three Operators and Interacts with RNA Polymerase

- Each dimer in a repressor tetramer can bind an operator, so that the tetramer can bind two operators simultaneously.
- Full repression requires the repressor to bind to an additional operator downstream or upstream as well as to the primary operator at the *lacZ* promoter.
- Binding of repressor at the operator stimulates binding of RNA polymerase at the promoter but precludes transcription.




Figure 26.21: If both dimers in a repressor tetramer bind to DNA, the DNA between the two binding sites is held in a loop. Third operator not shown.

26.11 Positive regulators of transcription

The *lac* Operon Has a Second Layer of Control: Catabolite Repression

- **catabolite repression** – The ability of glucose to prevent the expression of a number of genes.
 - In bacteria this is a positive control system; in eukaryotes, it is completely different.
- **Catabolite repressor protein (CRP)** is an activator protein that binds to a target sequence at a promoter.
- CRP is or was also known as cAMP-dependent, **Catabolite Activator Protein (CAP)**, a newer name I find clearer.

26.11 The *lac* Operon Has a Second Layer of Control: Catabolite Repression




Figure 26.25: cAMP converts an activator protein CRP to a form that binds the promoter and assists RNA polymerase in initiating transcription.

26.11 The *lac* Operon Has a Second Layer of Control: Catabolite Repression




Figure 26.27: By reducing the level of cyclic AMP, glucose inhibits the transcription of operons that require CRP activity.

- A dimer of CRP is activated by a single molecule of **cyclic AMP (cAMP)**.
- cAMP is controlled by the level of glucose in the cell; a low glucose level allows cAMP to be made.
- CRP interacts with the C-terminal domain of the α subunit of RNA polymerase to activate it.

27.1 Introduction

Famous inducible genes and transcriptional repressors in *E. coli*

- **bacteriophage** (or **phage**) – A bacterial virus.
- **lytic infection** – Infection of a bacterium by a phage that ends in the destruction of the bacterium with release of progeny phage.
- **lysis** – The death of bacteria at the end of a phage infective cycle when they burst open to release the progeny of an infecting phage (because phage enzymes disrupt the bacterium's cytoplasmic membrane or cell wall).

27.1 Introduction





Figure 27.01: Lytic development involves the reproduction of phage particles with destruction of the host bacterium.

λ bacteriophage **plaques** on a lawn of *E. coli* are **clear** where all cells were killed by virus infection and cell lysis (- like a cold sore).

Cloudy plaques contain λ **lysogen** cells that are **immune** to further lambda virus infection leading directly to cell lysis



27.9 The Lambda Repressor and Its Operators Define the Immunity Region

- **immunity** – In phages, the ability of a prophage to prevent another phage of the same type from infecting a cell.
- **virulent mutations** – Phage mutants that are unable to establish lysogeny.

27.9 The Lambda Repressor and Its Operators Define the Immunity Region

- Several lambdoid phages have different **immunity regions**.
- A lysogenic phage confers immunity to further infection by any other phage with the same immunity region.





Figure 27.16: In the absence of repressor, RNA polymerase initiates at the left and right promoters.

27.8 Lysogeny Is Maintained by the Lambda Repressor Protein




- The lambda repressor, encoded by the *cl* gene, is required to maintain lysogeny.
- The lambda repressor acts at the O_L and O_R operators to block transcription of the immediate early genes.
- The immediate early genes trigger a regulatory cascade; as a result, their repression prevents the lytic cycle from proceeding.

Figure 27.15: Repressor acts at the left operator and right operator to prevent transcription of the immediate early genes (*N* and *cro*).

Need to purify scarce gene regulator proteins drove biotechnology of protein overexpression

- Take *E. coli* cell as a cube 1 micron on each side. A liter is a cube 10 cm on each side. What is the concentration of the lac operator? Just how scarce can repressor be in a cell? 10-100 molecules if the k_d for operator-binding is 10^{-10} M or higher.
- Lambda repressor was first isolated from an overproducer mutant virus. Later the repressor gene was expressed from *lac* or *tac* promoters. Also, a hybrid ribosome binding site upstream of ATG gave strong translation initiation.

27.10 The DNA-Binding Form of the Lambda Repressor Is a Dimer



- A repressor monomer has two distinct domains.
- The N-terminal domain contains the DNA-binding site.
- The C-terminal domain dimerizes.
- Binding to the operator requires the dimeric form so that two DNA-binding domains can contact the operator simultaneously.
- Cleavage of the repressor between the two domains reduces the affinity for the operator and induces a lytic cycle.

Figure 27.18: Repressor dimers bind to the operator.

27.11 Lambda Repressor Uses a Helix-Turn-Helix Motif to Bind DNA

- Each DNA-binding region in the repressor contacts a half-site in the DNA.
- The DNA-binding site of the repressor includes two short α -helical regions that fit into the successive turns of the major groove of DNA (**helix-turn-helix**).
- A DNA-binding site is a (partially) palindromic sequence of 17 bp.



Figure 27.19: The operator is a 17-bp sequence with an axis of symmetry through the central base pair.


Lambda repressor binds as a symmetrical protein dimer to a nearly symmetrical 17 bp DNA sequence.

Symmetric repressor site

5' TATCACCGCCGGTGATA
ATAGTGGCGGCCACTAT 5'

FIGURE 16-11 Binding of a protein with a helix-turn-helix domain to DNA.

The protein, as is typically the case, binds as a dimer, and the two subunits are indicated by the shaded circles. The helix-turn-helix motif on each monomer is indicated; the "recognition helix" is labeled R.



27.11 Lambda Repressor Uses a Helix-Turn-Helix Motif to Bind DNA

- The amino acid sequence of the **recognition helix** makes contacts with particular bases in the operator sequence that it recognizes.




Figure 27.22: Two proteins that use the two-helix arrangement to contact DNA recognize lambda operators with affinities determined by helix-3.

Recognition helix amino acid side-chains make many sequence-specific contacts to base pairs.







FIGURE 16-12 Hydrogen bonds between λ repressor and base pairs in the major groove of its operator. Diagram of the repressor-operator complex, showing hydrogen bonds (in dotted lines) between amino acid side chains and bases in the consensus half-site. Only the relevant amino acid side chains are shown. In addition to Gln44 and Ser45 in the recognition helix, Asn55 in the loop following the recognition helix also makes contact with a specific base. Furthermore (and unusual to this case, see later in the text) Lys4 in the N-terminal arm of the protein makes a contact in the major groove on the opposite face of the DNA helix. Gln33 contacts the backbone. (Source: Redrawn from Jordan, S. and Pabo, C. *Science* 242: 896, Fig. 3B.)

Example of an amino acid contact that can discriminate between bases in a binding site.



The position of the glutamine is fixed when repressor binds.

Glutamine side chain needs to find the hydrogen donor and acceptor sites in just the right places.

It cannot reach the paired base on the other strand.

It cannot reach the other bases on the same strand.

Only an A base meets the criteria.

27.12 Lambda Repressor Dimers Bind Cooperatively to the Operator

- Repressor binding to one operator increases the affinity for binding a second repressor dimer to the adjacent operator.
- The affinity is $10 \times$ greater for O_L1 and O_R1 than other operators, so they are bound first.
- Cooperativity allows repressor to bind the O_L2/O_R2 sites at lower concentrations.




Figure 27.25: When two lambda repressor dimers bind cooperatively, each of the subunits of one dimer contacts a subunit in the other dimer.

27.13 Lambda Repressor Maintains an Autoregulatory Circuit

- The DNA-binding region of repressor at O_{R2} contacts RNA polymerase and stabilizes its binding to P_{RM} .
- This is the basis for the autoregulatory control of repressor maintenance.
- Repressor binding at O_L blocks transcription of gene N from P_L .





Figure 27.26: Positive control mutations identify a small region at helix-2 that interacts directly with RNA polymerase.

27.13 Lambda Repressor Maintains an Autoregulatory Circuit



- Repressor binding at O_R blocks transcription of cro , but also is required for transcription of cI .
- Repressor binding to the operators therefore simultaneously blocks entry to the lytic cycle and promotes its own synthesis.

Figure 27.27: Lysogeny is maintained by an autoregulatory circuit.

27.14 Cooperative Interactions Increase the Sensitivity of Regulation

- Repressor dimers bound at O_L1 and O_L2 interact with dimers bound at O_R1 and O_R2 to form octamers.
- These cooperative interactions increase the sensitivity of regulation.




Figure 27.29: O_L3 and O_R3 are brought into proximity by formation of the repressor octamer.

Transcription attenuation in the leader of the *E. coli trp* operon

26.12 The *trp* Operon Is a Repressible Operon with Three Transcription Units

- The *trp* operon is negatively controlled by the level of its product, the amino acid tryptophan (**autoregulation**).
- The amino acid tryptophan activates an inactive repressor encoded by *trpR*.
- A repressor (or activator) will act on all loci that have a copy of its target operator sequence.




Figure 26.32: Operators may lie at various positions relative to the promoter.

26.13 The *trp* Operon Is Also Controlled by Attenuation

- **attenuation** – The regulation of bacterial operons by controlling termination of transcription at a site located before the first structural gene.




Figure 26.33: Termination can be controlled via changes in RNA secondary structure that are determined by ribosome movement.

26.13 The *trp* Operon Is Also Controlled by Attenuation

- An **attenuator** (intrinsic terminator) is located between the promoter and the first gene of the *trp* cluster.
- The absence of Trp-tRNA suppresses termination and results in a 10× increase in transcription.




Figure 26.34: An attenuator controls the progression of RNA polymerase into the *trp* genes.

26.14 Attenuation Can Be Controlled by Translation

- The leader region of the *trp* operon has a 14-codon open reading frame that includes two codons for tryptophan.
- The structure of RNA at the attenuator depends on whether this reading frame is translated.
- In the presence of Trp-tRNA, the leader is translated to a **leader peptide**, and the attenuator is able to form the hairpin that causes termination.

26.14 Attenuation Can Be Controlled by Translation




Figure 26.35: The *trp* operon has a short sequence coding for a leader peptide that is located between the operator and the attenuator.

26.14 Attenuation Can Be Controlled by Translation




Figure 26.36: The trp leader region can exist in alternative base-paired conformations.




Figure 26.37: The alternatives for RNA polymerase at the attenuator depend on the location of the ribosome.

26.14 Attenuation Can Be Controlled by Translation

- In the absence of Trp-tRNA, the **ribosome stalls** at the tryptophan codons and an alternative secondary structure prevents formation of the hairpin, so that transcription continues.






Figure 26.38: In the presence of tryptophan tRNA, ribosomes translate the leader peptide and are released.

Mechanism of transcription activation of the yeast *GAL* genes.



B



A classical model of inducible gene expression in a eukaryote.


- ***GAL1*** galactokinase (mutant allele is *gal1*)
- ***GAL2*** galactose permease
- ***GAL7*** galactose uridyl transferase
- ***GAL10*** galactose-glucose epimerase

- ***GAL4*** positive regulator (mutant is *gal4*)
- ***GAL80*** negative regulator (mutant is *gal80*)

Yeast *GAL* genes are positively regulated at the level of transcription activation.

- A *gal4*, *gal80* double mutant fails to induce expression of galactose enzymes.
- Interpretation is that **GAL4** targets the structural genes to activate transcription.
- **GAL80** interacts with **GAL4** to control its activity.

Deletion analysis of yeast *GAL1* upstream region defined an Upstream Activating Site (UAS_G).



Four copies of a nearly symmetric 17 bp GAL4 site.


GAL1-lacZ and GAL10-lacZ fusions used for deletion analysis.

UAS_G had many properties of the Enhancer defined in SV40.


Experiments on the mechanism of transcription activation by GAL4.

Separation of transcription activation from DNA-binding function.


DNA-binding and transcription activation are separable functions.



DNA-binding and transcription activation are separable functions.




Genes and Signals, © 2002 by Cold Spring Harbor Laboratory Press, Chapter 2, Figure 5.




Genes and Signals, © 2002 by Cold Spring Harbor Laboratory Press, Chapter 2, Figure 4.

Current understanding of yeast *GAL* gene activation



- GAL1-GAL80 complex DOES bind UAS_G DNA (a is wrong on this).
- GAL3 is the galactose sensor for GAL gene induction
- GAL3 is closely related to GAL1, also binds galactose and ATP
- GAL1 itself acts instead of GAL3 to bind GAL80 in *Kluveromyces lactis*, the whey (skimmed milk) yeast, which hydrolyses lactose and then converts the galactose to glucose

Deletion analysis of yeast *GAL1* upstream region defined an Upstream Activating Site (UAS_G) and Mig1 site.



Four copies of a nearly symmetric 17 bp GAL4 site.


Mig1 protein mediates glucose repression of *GAL1*

Mig1 recruits Ssn6 and Tup1

Mig1 also binds *GAL4* promoter and lowers *GAL4* protein levels


Eukaryotic gene activators recruit RNA polymerase to the promoter.

- Activating regions associate with mediator which in turn associates with RNA polymerase.
- Screen for yeast mutants with increased activation by GAL4(1-100) identified a potentiator mutant in GAL11, a component of Mediator.
- Gal11 when fused to LexA activates transcription. This is an activator bypass experiment.
- Results favour the idea that GAL4 recruits polymerase to the promoter by helping it bind as CAP/CRP does in *E. coli*.



Genes and Signals, © 2002 by Cold Spring Harbor Laboratory Press, Chapter 2, Figure 13.


Uses of GAL4 Number 1: The yeast two hybrid system for identification of interacting proteins.




Can be used on a genomic scale to test all proteins against all others. Target protein fused to DNA-binding domain in cells of one mating type mated to a library of activating region fusion proteins.



Uses of GAL4 Number 2: GAL4 activates transcription from UAS_G in *Drosophila*.

Widely used for targeted protein expression in *Drosophila* (GAL4-UAS two-component system).





Induction of ectopic eyes by GAL4-targeted expression of the *eyeless* gene in *Drosophila*



The GAL4-UAS binary system is used to map brain neurons involved in memories and behaviours and to target gene expression there.

- *rutabaga* mutant flies lack an adenyl-cyclase required for synaptic plasticity and cannot learn a variety of training tasks.
- A UAS-RUTABAGA construct was expressed under the control of different GAL4 drivers in a *rutabaga* mutant fly to see which brain cells are needed to learn particular tasks.
- Mushroom body cells (somewhat similar to mammalian hippocampus) need RUTABAGA protein at synapses to learn to avoid particular odours and central complex cells need it for visual learning.
- *Drosophila* connectome is all the neuron types and pathways in the whole CNS

Liam.Keegan@ceitec.muni.cz

Lewin, GENES XI

Chapter 26

Operons, prokaryotic gene regulation and
yeast GAL4

[Lewin's GENES XII \(PDF\)](#)

<https://pdfroom.com/books/lewins-genes-xii/Wx5aDYKI2BJ>



[biotecher.ir](#)

<https://biotecher.ir/wp-content/uploads/2018/06/bio...>

Showing that GAL4 binds the *GAL UAS*.

**Methods for identifying and characterising
sequence-specific DNA-binding proteins.**

DNase I protection assay to precisely define binding sites of sequence-specific DNA binding proteins and to measure DNA-binding affinity, (DNase I Footprinting)




The GAL4-binding site is CGG-N(11)-CCG.

The GAL4 DNA-binding domain has a Zn_2Cys_6 binuclear cluster structure found in over 60 proteins but restricted to fungi.

B

| | $\alpha 1$ | Linker | $\alpha 2$ | 3_{10} | DNA Site |
|-------|-------------------------------------|-------------------|-------------------------------|----------|---|
| Gal4p | A C D I C R L K K L R C S K | -- E K P K -- | C A K C L K N N W E | -- C R Y | CGG (N ₁₁) CCG |
| Put3p | A C L S C R K R H I K C P G | -- G N P -- | C Q K C V T S N A I | -- C E Y | CGG (N ₁₀) CCG |
| Ppr1p | A C K R C R L K K I K C D Q | -- E F P S -- | C K R C A K L E V P | -- C V Y | CGG (N ₆) CCG |
| Leu3p | A C V E C R Q Q K S K C D A H E R A | -- E P F C T K -- | C A K K N V P | -- C I L | CGG (N ₂) CCG |
| Hap1p | S C T I C R K R K V K C D K | -- L R P H -- | C Q Q C T K T G V A H L C H Y | | CGG (N ₆) CCG (CGC, AGG, TGG) |
| Cep3p | F C S V C T F R R K V K C D R | -- M I P -- | C G N C R K R G Q D S E C M K | | TGT (N ₉) CCG |



Since GAL4 has a type of DNA-binding domain not seen in animals we will discuss DNA sequence recognition and more common types of DNA-binding domains tomorrow.

Summary

RNA polymerases and RNA pol II promoters.

TAFs

Transcriptional regulation in yeast.

Uses of GAL4

DNA looping is the simplest explanation for GAL4 action.




Figure 6.22 DNA looping

Transcription factors bound at distant enhancers are able to interact with general transcription factors at the promoter because the intervening DNA can form loops. There is therefore no fundamental difference between the action of transcription factors bound to DNA just upstream of the promoter and to distant enhancers.

Condensation and decondensation of chromatin in the cell cycle.






FIGURE 8 Representation of the stages in folding of a nucleosomal chain (*left*) via the zig-zag ribbon into the helical ribbon (*right*) conformation. To avoid confusion, the DNA is shown only in places, and some underlying nucleosomes have been shaded.

- Chromatin condenses during mitosis and decondenses in interphase when most genes are expressed.
- Histone deacetylases (HDACs) facilitate chromatin condensation and contribute to gene repression (Hda1, Rpd3).
- Histone acetyl transferases (HATs), like GCN5 in the SAGA complex, acetylate histone tails and contribute to chromatin decondensation and gene activation.
- Nucleosome remodelling complexes (Swi/Snf complex), can remove nucleosomes from a promoter region or deposit them there.

Imposing a requirement for nucleosome modifiers for *GAL1* activation.

Neither GCN5 histone acetyltransferase nor the nucleosome remodelling complex Swi/Snf complex are required for GAL1 activation nor to activate 95% of yeast genes.

Nevertheless GAL4 recruits these activities to the promoter and they become necessary if GAL4 sites are weakened or when chromosomes condense for mitosis.



Unusual chromosomes in *Saccharomyces cerevisiae*

- The nuclear membrane was seen only after the electron microscope was invented. The nuclear membrane does not break down at mitosis.
- The genome is small and divided among 16 chromosomes. *S. cerevisiae* lacks a Histone protein 1 that binds the linker DNA between nucleosomes and causes them to condense more. Cell division takes place without formation of visible metaphase chromosomes.
- The fission yeast *Schizosaccharomyces pombe* has fewer chromosomes and these become visible at metaphase. It is quite different from budding yeast in DNA sequence and was promoted as being more like mammals by Paul Nurse and others who used it to study the cell cycle, ("*Schizophrenic pombe*").

26.15 Stringent Control by Stable RNA Transcription

- Poor growth conditions cause bacteria to produce the small-molecule regulators (p)ppGpp.
- **Stringent response** – The ability of a bacterium to shut down synthesis of ribosomes and tRNA in a poor growth medium.
- The trigger for the reaction is the entry of uncharged tRNA into the ribosomal A site.
- (p)ppGpp competes with ATP during formation of the open complex during transcription initiation by RNA polymerase and inhibits the reaction.

26.15 Stringent Control by Stable RNA Transcription





Figure 26.40: Stringent factor catalyzes the synthesis of pppGpp and ppGpp; ribosomal proteins can dephosphorylate pppGpp to ppGpp.

- **Relaxed mutants** – In *E. coli*, these do not display the stringent response to starvation for amino acids (or other nutritional deprivation).
- **Stringent factor** – The protein RelA, which is associated with ribosomes. It synthesizes ppGpp and pppGpp when an uncharged tRNA enters the ribosome.

26.16 r-Protein Synthesis Is Controlled by Autoregulation

- Translation of an r-protein operon can be controlled by a product of the operon that binds to a site on the polycistronic mRNA.

When rRNA is available, the r-proteins associate with it. Translation of mRNA continues.



When no rRNA is available, r-proteins accumulate. An r-protein binds to mRNA and prevents translation.




Figure 26.42: Translation of the r-protein operons is autogenously controlled and responds to the level of rRNA.

26.10 The Operator Competes with Low-Affinity Sites to Bind Repressor

- Proteins that have a high affinity for a specific DNA sequence also have a low affinity for other DNA sequences.
- Every base pair in the bacterial genome is the start of a low-affinity binding site for repressor.

| DNA | Repressor | Repressor + inducer |
|-----------------|--------------------|---------------------|
| Operator | 2×10^{13} | 2×10^{10} |
| Other DNA | 2×10^6 | 2×10^6 |
| Specificity | 10^7 | 10^4 |
| Operators bound | 96% | 3% |
| Operon is: | repressed | induced |

Figure 26.23: lac repressor binds strongly and specifically to its operator, but is released by inducer. All equilibrium constants are in M⁻¹.

26.10 The Operator Competes with Low-Affinity Sites to Bind Repressor




Figure 26.24: Virtually all the repressor in the cell is bound to DNA.

- The large number of low-affinity sites ensures that all repressor protein is bound to DNA.
- Repressor binds to the operator by moving from a low-affinity site rather than by equilibrating from solution.

26.10 The Operator Competes with Low-Affinity Sites to Bind Repressor

- In the absence of inducer, the operator has an affinity for repressor that is 10^7 times that of a low-affinity site.
- The level of 10 repressor tetramers per cell ensures that the operator is bound by repressor 96% of the time.
- Induction reduces the affinity for the operator to 10^4 times that of low-affinity sites, so that operator is bound only 3% of the time.

26.1 Introduction

negative control of transcription by a repressor removed by an inducer (lac repressor + IPTG),

positive control of transcription by activator and inducer (CAP + cAMP),

(also possible, lower half,

negative control of transcription by a repressor and corepressor

and positive control of transcription by having an activator that is removed)







Figure 26.04: Regulatory circuits can be designed from all possible combinations of positive and negative control with inducible and repressible control.

Spare slides!


**DNA is superior to dsRNA as a genomic material because of the major groove.
Specific base recognition by proteins is easy in DNA.**



dsRNA
A-helix
Deep and narrow
Major groove.



DNA
B-form
Wide and
Accessible
Major groove.



Wide major groove means DNA sequence can be recognized by DNA-binding proteins.




FIGURE 6-10 Chemical groups exposed in the major and minor grooves from the edges of the base pairs. The letters in red identify hydrogen bond acceptors (A), hydrogen bond donors (D), nonpolar hydrogens (H), and methyl groups (M).

26.1 Introduction

- **regulator gene** – A gene that codes for a product (typically protein) that controls the expression of other genes (usually at the level of transcription).
- **structural gene** – A gene that codes for any RNA or protein product other than a regulator.




Figure 26.01: A regulator gene codes for a protein that acts at a target site on DNA.

26.1 Introduction

- ***trans-acting*** – A product that can function on any copy of its target DNA. This implies that it is a diffusible protein or RNA.
- ***cis-acting*** – A site that affects the activity only of sequences on its own molecule of DNA (or RNA); this property usually implies that the site does not code for protein.

26.1 Introduction




Figure 26.04: Regulatory circuits can be designed from all possible combinations of positive and negative control with inducible and repressible control.

- Gene regulation *in vivo* can utilize any of these mechanisms, resulting in all four combinations: **negative inducible**, **negative repressible**, **positive inducible**, and **positive repressible**.

Beginnings of genetics in *Saccharomyces cerevisiae*.

- Yeast genetics began only in the 1950s - much later than *E. coli* genetics. Commercial strains of brewer's and baker's yeast are stable diploids or tetraploids and could not be studied genetically.
- Occasional meiosis in diploid stains gives an ascus containing four haploid spores two with *a* and two with α mating types but these immediately mate back to diploids (*Ascomycete* family of fungi). Pure haploid cultures could not be grown from spores because after the first haploid cell division the mother cell switches mating type and can mate with cell it just budded off.
- Yeast genetics required mutants in a gene called *HO* (*homothallism* – is a plant term for having both types of sexual parts on one organism). The *ho* mutation keep haploid cell lines stable because it prevents mating type switching. Winge, (Carlsberg labs, Copenhagen), Lindegren, (Anheuser Busch, Carbondale, Illinois).
- Much of yeast genetics is done with the haploid strains that have only one copy of each gene (same as *E. coli*) but can be mated to give diploids and sporulated back to haploids by nitrogen starvation.

Genome sizes increase faster than gene numbers among the main model organisms.

| Sequenced genomes vary from 470 to 30,000 genes | | | |
|---|--------------|---------|-------------|
| Species | Genomes (Mb) | Genes | Lethal loci |
| <i>Mycoplasma genitalium</i> | 0.58 | 470 | ~300 |
| <i>Rickettsia prowazekii</i> | 1.11 | 834 | |
| <i>Haemophilus influenzae</i> | 1.83 | 1,743 | |
| <i>Methanococcus jannaschi</i> | 1.66 | 1,738 | |
| <i>B. subtilis</i> | 4.2 | 4,100 | |
| <i>E. coli</i> | 4.6 | 4,288 | 1,800 |
| <i>S. cerevisiae</i> | 13.5 | 6,034 | 1,090 |
| <i>S. pombe</i> | 12.5 | 4,929 | |
| <i>A. thaliana</i> | 119 | 25,498 | |
| <i>O. sativa (rice)</i> | 466 | ~30,000 | |
| <i>D. melanogaster</i> | 165 | 13,601 | 3,100 |
| <i>C. elegans</i> | 97 | 18,424 | |
| <i>H. sapiens</i> | 3,300 | ~25,000 | |

Bacterial minimum 1800.

Yeast has 6,000 genes. About 2X basic eukaryotic minimum set of 2,700 or so.

Simpler animals like *Drosophila* have 2X as many genes as yeast.

Humans and other mammals have only 2X *Drosophila*.

4,600 kb

13,500 kb

3x genome size of *E. coli*

165,000 kb

10X genome size of yeast

3,000,000 kb

20X *Drosophila* genome