




SOLVENT SELECTION

Petr Beňovský



IMPORTANCE OF SOLVENTS

- Reaction medium (transport, combine, separate)
 - Dissolution
 - (In)solubility
 - Kinetics
 - Health
 - Safety
 - Environmental aspects
 - Course of the reaction
 - Cost (purchase, recycle, dispose)
- 

PERSPECTIVE ON SOLVENT SELECTION

| ASPECT | COMMENT |
|---|--|
| Safety | Avoid solvents that are toxic or highly flammable |
| Promote high-yielding reactions | Compatible with desired chemistry; Can isolate product in good yield? |
| Convenient (minimize processing operations) | Isolate product from reaction solvent? Operate at high concentrations? |
| Water miscibility | Azeotroping ability Control amount of water |
| Cost of bulk and recoverability | More important at the end of development cycle |
| Environmental | Ethics, and cost of recovery and non-compliance |
| Long-term availability | |
| Acceptability for human use | |
| Water as solvent | But, recovery of product from aqueous layer can be costly, plus cost of disposal |
| Neat reactions | |
| Ionic liquids | |

SOLVENT SELECTION GUIDES

Almost every company created specific solvent guide
 SmithKline Beecham – Curzons, A.D. *et al* *Clean Products and Processes* 1, 82 (1999)

| SOLVENT | | Waste | Impact | Health | Safety |
|-----------|------------------------------------|-------|--------|--------|--------|
| Alcohols | Ethylene glycol | 4 | 9 | 8 | 10 |
| | 1-Butanol | 5 | 7 | 8 | 8 |
| | Diethylene glycol mono butyl ether | 5 | 8 | 8 | 10 |
| | Ethanol / IMS | 3 | 7 | 9 | 6 |
| | 2-Propanol | 3 | 10 | 7 | 7 |
| | Methanol | 3 | 8 | 4 | 8 |
| | 2-Methoxy ethanol | 4 | 9 | 2 | 7 |
| Esters | Butyl acetate | 7 | 7 | 7 | 6 |
| | Propyl acetate | 7 | 6 | 7 | 6 |
| | Isopropyl acetate | 5 | 7 | 7 | 6 |
| | Ethyl acetate | 4 | 9 | 7 | 4 |
| | Methyl acetate | 2 | 6 | 5 | 5 |
| Aromatics | Xylene | 8 | 4 | 5 | 5 |
| | Toluene | 7 | 3 | 5 | 4 |

SOLVENT SELECTION GUIDES

Sanofi – Prat D. *et al Org.Process Res. Dev.* 17, 1517 (2013)

| Solvents Guide | ETHERS: OVERVIEW | | | | |  SANOFI |
|-----------------------------------|------------------------|-----------------|-------------|--------|---------------|---|
| Name | Overall ranking | ICH limit (ppm) | Occ. health | Safety | Environment | Other concern |
| Diethyl ether | Banned | 5000 | OEBV2 | SHB5 | EHB2 | Peroxides, VOC |
| Diisopropyl ether | Substitution advisable | Not listed | OEBV2 | SHB5 | EHB3 | Peroxides |
| Dibutyl ether | Substitution advisable | Not listed | OEBV2 | SHB5 | EHB3 | Peroxides, odor |
| THF | Substitution advisable | 720 | OEBV3 Sk | SHB4 | EHB2 | VOC, miscible with water, peroxides |
| Methyl-THF | Recommended | Not listed | OEBV2 | SHB4 | EHB3 | Peroxides, cost |
| Dioxane | Substitution requested | 380 | OEBV3 Sk | SHB5 | EHB2 | Miscible with water, peroxides |
| Anisole | Recommended | 5000 | OEBV2 | SHB3 | EHB2 | Odor |
| MTBE | Substitution advisable | 5000 | OEBV3 Sk | SHB5 | EHB3 | VOC |
| ETBE | Substitution requested | Not listed | OEBV4 | SHB5 | EHB3 | Peroxides, lack of data |
| CPME | Substitution requested | Not listed | OEBV3 | SHB5 | EHB3 | Peroxides, one supplier only |
| Dimethoxy ethane | Substitution requested | 100 | OEBV4 G2 | SHB4 | EHB2 | CMR (R1B), peroxides |
| Diglyme | Substitution requested | Not listed | OEBV4 G2 | SHB4 | EHB2 | CMR (R1B), peroxides |
| Diethoxymethane | Substitution requested | Not listed | OEBV4 | SHB5 | Not available | Reactive, considered as CMR |

SOLVENT SELECTION GUIDES

GlaxoSmithKline – Henderson, R.K. *et al Green Chemistry* 13, 854 (2011)

| Solvent | Cas number | Melting point °C | Boiling Point °C | Waste | Environmental Impact | Health | Flammability & Explosio | Reactivity/ Stability | Life Cycle Score | Legislation Flag |
|-------------------------|------------|------------------|------------------|-------|----------------------|--------|-------------------------|-----------------------|------------------|------------------|
| Isopropyl acetate | 108-21-4 | -73 | 89 | 5 | 7 | 7 | 6 | 9 | 7 | |
| Dimethyl carbonate | 616-38-6 | -1 | 91 | 4 | 8 | 7 | 6 | 10 | 8 | |
| Ethyl acetate | 141-78-6 | -84 | 77 | 4 | 8 | 8 | 4 | 8 | 6 | |
| t-Butylmethyl ether | 1634-04-4 | -109 | 55 | 4 | 5 | 5 | 3 | 9 | 8 | |
| 2-Methyltetrahydrofuran | 96-47-9 | -137 | 78 | 4 | 5 | 4 | 3 | 6 | 4 | |
| Dichloromethane | 75-09-2 | -95 | 40 | 3 | 6 | 4 | 6 | 9 | 7 | |
| Chloroform | 67-66-3 | -64 | 61 | 3 | 6 | 3 | 6 | 9 | 6 | |

SOLVENT SELECTION GUIDES

ACS Green Chemistry Institute® Roundtable Solvent Selection Guide, March 2011

| Substance Information | | | Scoring Information | | | | |
|-----------------------|-------------------------|------------|---------------------|--------|-----------|-------------|-------------|
| Solvent Class | Solvent Name | CAS Number | Safety | Health | Env (Air) | Env (Water) | Env (Waste) |
| Acid | ACETIC ACID | 64-19-7 | 3 | 6 | 6 | 3 | 6 |
| Acid | ACETIC ANHYDRIDE | 108-24-7 | 3 | 6 | 6 | 2 | 7 |
| Acid | FORMIC ACID | 64-18-6 | 2 | 6 | 5 | 4 | 7 |
| Acid | METHANE SULPHONIC ACID | 75-75-2 | | | 6 | 6 | 10 |
| Acid | PROPIONIC ACID | 79-09-4 | 2 | 5 | 6 | 4 | 6 |
| Alcohol | 1-BUTANOL | 71-36-3 | 3 | 5 | 5 | 5 | 3 |
| Alcohol | 1-PROPANOL | 71-23-8 | 4 | 4 | 6 | 2 | 6 |
| Alcohol | 2-BUTANOL | 78-92-2 | 4 | 5 | 6 | 3 | 5 |
| Alcohol | 2-METHOXYETHANOL | 109-86-4 | 4 | 9 | 5 | 3 | 7 |
| Alcohol | BENZYL ALCOHOL | 100-51-6 | 4 | 3 | 4 | 2 | 4 |
| Alcohol | ETHANOL | 64-17-5 | 4 | 3 | 5 | 1 | 6 |
| Alcohol | ETHYLENE GLYCOL | 107-21-1 | 3 | 3 | 5 | 1 | 7 |
| Alcohol | ISOAMYL ALCOHOL | 123-51-3 | 3 | 4 | 5 | 3 | 4 |
| Alcohol | ISOBUTANOL | 78-83-1 | 3 | 5 | 4 | 3 | 3 |
| Alcohol | ISOPROPYL ALCOHOL (IPA) | 67-63-0 | 5 | 5 | 6 | 2 | 6 |
| Alcohol | METHANOL | 67-56-1 | 3 | 5 | 6 | 3 | 6 |
| Alcohol | T-BUTANOL | 75-65-0 | 3 | 5 | 7 | 2 | 6 |

SOLVENT CONSIDERATIONS

Watch out hydrocarbon solvents with even number of carbons (toxicity, electrostatic buildup);

Classification of solvents – ICH Harmonised Guideline Q3C – Impurities: Guideline for Residual Solvents

Class 1 – solvents to be avoided (known human carcinogens, strongly suspected human carcinogens, and/or environmental hazards, e.g. carbon tetrachloride (concentration limit 4 ppm), 1,2-dichloroethane (5 ppm), 1,1,1-trichloroethane (1500 ppm), benzene (2 ppm))

Class 2 – solvents to be limited (non-genotoxic animal carcinogens, agents of irreversible toxicity, e.g. acetonitrile (410 ppm), chlorobenzene (360 ppm), chloroform (60 ppm), *N,N*-dimethylformamide (880 ppm), hexane (290 ppm), methanol (3000 ppm), *N*-methylpyrrolidone (530 ppm), toluene (890 ppm))


Class 3 – solvents with low toxic potential (permissible daily exposure 50 mg or more per day, e.g. acetic acid, acetone, ethyl acetate, heptane, 2-propanol, triethylamine)

Solvents for which no adequate toxicological data was found – a manufacturer is asked to supply justification for residual levels of these solvents (e.g. diisopropyl ether, petroleum ether, trifluoroacetic acid)

SOLVENT CONSIDERATIONS




The best reaction solvent is the one that crystallizes the product directly from the reaction;

Novartis – the Dimroth rearrangement – temperature and solubility turned out to be the most important – the product simply precipitated from the reaction mixture;



Fischer, R.W. *Org. Process Res. Dev.* 5, 581 (2001)

SOLVENT CONSIDERATIONS



| Solvent | Yield (% after 48 h) |
|--------------|----------------------|
| Benzene | 0 |
| Toluene | 20 |
| Methanol | 80 |
| THF | 91 |
| Methanol-THF | 93 |

Dyson, P.J.; Jessop, P.G. *Catal. Sci. Technol.* **6**, 3302 (2016)



SOLVENT CONSIDERATIONS

Homogeneous vs. Heterogeneous
Reactions using gases

Insolubility is sometimes advantageous (the Schotten-Baumann reaction, the Finkelstein reaction)

Menshutkin (1890)

The reaction rate of the reaction rate of triethylamine with alkyl halides providing quaternary ammonium salts strongly depends on a solvent (hexane **1**, acetone **338**, benzyl alcohol **739**)

Solvents commonly used in **academia** are not often welcomed for **industrial** applications


Safety first !



SOLVENT CONSIDERATIONS


The Finkelstein reaction

The reaction driven to completion by exploiting the different solubility of used and formed halides



SOLVENT CONSIDERATIONS

The Halex reaction




toluene no reaction

water no reaction

DMF satisfying conversion, KCl precipitates out of the reaction mixture

MIXING OF HETEROGENEOUS MIXTURES



Anderson, N.G. *Practical Process Research and Development, A guide for organic chemists*, 2nd Edition, Elsevier 2000

SOLVENT CONSIDERATIONS


| Solvent | TWA (ppm) |
|------------------|-----------|
| Acetone | 500 |
| EtOAc | 400 |
| MeOH | 200 |
| <i>t</i> -BuOH | 100 |
| MTBE | 50 |
| MeCN | 20 |
| DMF | 10 |
| Pyridine | 1 |
| 2-Methoxyethanol | 0.1 |

TWA = Time-Weighted Average shift for five days for nearly safe exposure over an 8 h shift for five days for nearly all workers



SOLVENT CONSIDERATIONS

Also, always consider physical-chemical properties like

- Flash point
 - Flammability
 - Boiling point
 - Melting point
 - Electrostatic charge accumulation
 - Recycling potential
 - Cost of solvent
 - Environmental aspects
 - Cost of disposal
 - Polarity
- 

SOLVENT CONSIDERATIONS

Solvents **rarely used** in the pharmaceutical industry

| Solvent | Disadvantage | Alternative replacement |
|------------------|--|--------------------------------------|
| Diethylether | Flammable | MTBE |
| Diisopropylether | Peroxide formation | MTBE |
| Hexane | Electrostatic charge Neurological toxicity | Heptanes, <i>i</i> -octane |
| Chloroform | Mutagenicity, environmental aspects, toxicity | Dichloromethane, 2-MeTHF, toluene |
| Benzene | Toxicity | Toluene |
| Ethylene glycol | Toxicity | 1,2-Propandiol |
| Acetonitrile | Animal teratogen, potential acetamide generation (genotoxic) | 2-propanol, acetone - water |

SOLVENT CONSIDERATIONS

Solvents **preferred** for process development (Pfizer)

| Preferred | Usable | Undesirable |
|--------------------------|--------------------|------------------------|
| Water | Cyclohexane | Pentane |
| Acetone | Heptane | Hexanes |
| Ethanol | Toluene | Diisopropyl ether |
| 2-Propanol | Methyl cyclohexane | Diethyl ether |
| Ethyl acetate | MTBE | Dichloroethane |
| <i>i</i> -Propyl acetate | <i>i</i> -Octane | Dichloromethane |
| Methanol | 2-MeTHF | Chloroform |
| Methyl ethyl ketone | DMSO | DMF |
| <i>n</i> -Butanol | AcOH | NMP |
| <i>t</i> -Butanol | Ethylene glycol | 1,4-Dioxane |
| | | Benzene |
| | | Carbon tetrachloride |



SOLVENT CONSIDERATIONS

Polarity of solvents

Hughes-Ingold rules ((de)stabilization of transition state)

Aliphatic nucleophilic substitution and elimination reactions;
Considering **pure electrostatic interactions** between ions or dipolar molecules and solvent molecules in **initial** and **transition** states;



SOLVENT CONSIDERATIONS

Hughes-Ingold rules


| Reaction type | Initial reactants | Activated complex | Charge alteration during activation | Effect of increased solvent polarity on rate ^{a)} |
|----------------------|-----------------------------------|---|-------------------------------------|--|
| (a) S _N 1 | R—X | R ^{δ+} ... X ^{δ-} | Separation of unlike charges | Large increase |
| (b) S _N 1 | R—X ⁺ | R ^{δ+} ... X ^{δ+} | Dispersal of charge | Small decrease |
| (c) S _N 2 | Y + R—X | Y ^{δ+} ... R ... X ^{δ-} | Separation of unlike charges | Large increase |
| (d) S _N 2 | Y ⁻ + R—X | Y ^{δ-} ... R ... X ^{δ-} | Dispersal of charge | Small decrease |
| (e) S _N 2 | Y + R—X ⁺ | Y ^{δ+} ... R ... X ^{δ+} | Dispersal of charge | Small decrease |
| (f) S _N 2 | Y ⁻ + R—X ⁺ | Y ^{δ-} ... R ... X ^{δ+} | Destruction of charge | Large decrease |



SOLVENT CONSIDERATIONS



Hughes-Ingold rules


- An **increase** in solvent **polarity** results in an **increase** in the **rates** of those reactions in which the charge density is **greater** in the activated complex than in the initial reactant molecules;
 - An **increase** in solvent **polarity** results in a **decrease** in the **rates** of those reactions in which the charge density is **lower** than in the initial reactant molecules;
 - A change in solvent polarity will have **negligible effect** on the **rates** of those reactions that involve **little** or **no** change in the charge density from reactants to the activated complex.
- 

SOLVENT CONSIDERATIONS

Polarity of solvents

Reichardt, C. *Pure Appl. Chem* 76, 1903 (2004)

Reichardt, C. *Solvents and Solvent Effects in Organic Chemistry*,
3 rd Ed., Wiley-VCH, 2003



SOLVENT CONSIDERATIONS

Polarity of solvents

$$E_T(30) = h \cdot c \cdot N_A \cdot \nu_{max} \quad [\text{kcal} \cdot \text{mol}^{-1}]$$

Could be used even in solid rare gas matrices to establish polarity of rare gases (Xe → Kr → Ar → Ne decreasing polarizability)

$E_T(30)$ recommended and used for monitoring of a quality of disinfection solutions of alcohol/water in order to prevent SARS-CoV 2 infections


SOLVENT CONSIDERATIONS

Polarity of solvents

E_T^N parameter – the negative solvatochromism of the $\pi \rightarrow \pi^*$ shifts of solutions of the betaine dye – more polar solvents stabilize the ground energy of the polar dye, producing thus greater shift in the position of $\pi \rightarrow \pi^*$ absorption relative to that found for solutions of the dye in tetramethylsilane.


Colors of this dye in a solvent are indicative of the polarities of the solvent and solvent combination used to dissolve it.

SOLVENT CONSIDERATIONS



Reichardt, C. *J.Org.Chem* 87, 1616 (2022)

SOLVENT CONSIDERATIONS



Reichardt, C. *Pure Appl. Chem.* 76, 1903 (2004)

SOLVENT CONSIDERATIONS

Polarity of solvents

| Solvent | Polarity E_T^N | Solubility in water (wt%) | Bp of water-solvent azeotrope | wt% of water removed by azeotrope | ICH solvent class |
|---------------------------------|------------------|---------------------------|-------------------------------|-----------------------------------|-------------------|
| Water | 1.000 | - | None | None | |
| EtOH | 0.654 | ∞ | 78 °C | 4.0 | 3 |
| AcOH | 0.648 | ∞ | 77 °C | 97 | 3 |
| DMF | 0.404 | ∞ | None | None | 2 |
| Acetone | 0.355 | ∞ | None | None | 3 |
| CH ₂ Cl ₂ | 0.309 | 1.3 | 38 °C | 1.5 | 2 |
| Toluene | 0.099 | 0.06 | 84 °C | 13.5 | 2 |
| Et ₃ N | 0.043 | 5.5 | 75 °C | 10 | 3 |
| Heptane(s) | 0.012 | 0.0004 | 79 °C | 12.9 | 3 |
| Cyclohexane | 0.006 | 0.006 | 69 °C | 9 | 2 |

SOLVENT CONSIDERATIONS

Polarity of solvent mixtures

| Solvent | Polarity E_T^N | Solvent mixture | Calculated E_T^N |
|---|------------------|--|--------------------|
| MeOH | 0.762 | EtOH:H ₂ O = 6.9:3.1 | 0.762 |
| EtOH | 0.654 | Acetone:H ₂ O = 4.6:5.4 | 0.654 |
| H ₂ O:CH ₂ Cl ₂ = 0.2:99.8 | 0.310 | H ₂ O:MIBK = 1.9:98.1 | 0.283 |
| H ₂ O:CH ₂ Cl ₂ = 0.2:99.8 | 0.310 | H ₂ O:EtOAc = 3.3:96.7 | 0.253 |
| H ₂ O:CH ₂ Cl ₂ = 0.2:99.8 | 0.310 | H ₂ O:2-MeTHF = 5.3:94.7 | 0.223 |
| <i>i</i> -PrOAc | 0.210 | Heptanes:EtOAc = 0.8:9.2 | 0.210 |
| MeOH:H ₂ O = 7:1 | 0.792 | EtOH:H ₂ O = 5:3 | 0.783 |

SOLVENT CONSIDERATIONS

- Tendency of solvents to form **azeotropes with water** is considered advantageous (it is not practical and economical to dry solvents using drying agents on large scale);
- **Be careful** – dependence on pressure (**breaking the azeotrope**)

Effect of reducing distillation pressure on EtOAc – water:

| Pressure (mm) | Bp (°C) | Water in azeotrope (wt%) |
|---------------|---------|--------------------------|
| 760 | 70.4 | 8.5 |
| 250 | 42.6 | 6.3 |
| 25 | 1.9 | 3.6 |

IMPURITIES IN SOLVENTS

- Absolute solvents are rather expensive, common solvents contains some amount of water, for certain operations they should be dried (azeotropic distillation, molecular sieves, use of an excess of cheap reagent);

Denatured solvents (ethanol)


Stabilizers (e.g. BHT in THF)

Tendency to form (hydrogen)peroxides (diisopropyl ether, butadiene, acetaldehyde, 1,4-dioxane, styrene, acrylonitrile, 2-butanol, benzyl alcohol, THF, MIBK, 2-propyl alcohol)

IMPURITIES IN SOLVENTS

- Degradation of solvents (ethyl acetate, DMF)


Side reactions (reesterifications, reaction with dichloromethane)



Avoid unwanted formation of esters of sulfonic acids (potentially mutagenic)

SACRIFICIAL SOLVENTS


About 50 eq. of 98% H_2SO_4 at 50 °C for 3 h followed by an aqueous quench provided ring sulfonation in the product; 5-6 eq. of 98% H_2SO_4 in toluene at 70 °C for 2 h – sulfonation of the product was significantly diminished.





Harrington, P.J. *et al* *Org. Process Res. Dev.* 10, 1157 (2006)

IONIC LIQUIDS

- Ionic liquids are ionic compounds (salts) which are liquids below 100 °C. More commonly, ionic liquids have melting points below room temperature.



IONIC LIQUIDS




SUPERCRITICAL FLUIDS

| SCF | Name | T_c (°C) | p_c (bar) | d_c (g/mL) | MW | μ (De- bye) ^a | Cost ^b (\$/kg) |
|------------------|---------------------|---------------|----------------|-----------------|-------|------------------------------------|------------------------------|
| Ar | argon | -122.5 | 48.6 | 0.531 | 39.95 | 0 | 6 |
| CO ₂ | carbon dioxide | 31.1 | 73.8 | 0.466 | 44.01 | 0 | 3 |
| HCl | hydrogen chloride | 51.5 | 82.6 | 0.42 | 36.46 | 1.08 | 20 |
| HBr | hydrogen bromide | 90.0 | 85.5 | n.a. | 80.91 | 0.82 | 50 |
| HI | hydrogen iodide | 150.7 | 83 | n.a. | 127.9 | 0.44 | n.a. |
| H ₂ O | water | 374.0 | 220.6 | 0.322 | 18.02 | 1.85 | n.a. |
| NH ₃ | ammonia | 132.4 | 113.2 | 0.235 | 17.03 | 1.47 | 3 |
| N ₂ O | nitrous oxide | 36.4 | 72.5 | 0.453 | 44.01 | 0.167 | 50 |
| Kr | krypton | -63.76 | 54.9 | 0.912 | 83.80 | 0 | 3000 |
| SF ₆ | sulfur hexafluoride | 45.5 | 37.6 | 0.737 | 146.1 | 0 | 50 |
| Xe | xenon | 16.6 | 58.3 | 1.099 | 131.3 | 0 | 4000 |

Jessop, P.G.; Leitner, W. *Chemical Synthesis Using Supercritical Fluids*, Wiley-VCH 1999

SUPERCRITICAL CARBON DIOXIDE



Peach, J.; Eastoe, J. *Beilstein J.Org.Chem.* 10, 1878 (2014)

Beckman, E.J. *J.Supercritical Fluids* 28, 121 (2004)



SUPERCRITICAL CARBON DIOXIDE

FOOD INDUSTRY

Coffee decaffeination

Tea decaffeination

Fatty acids from spent barley


Vitamin E, hops, spices extraction

Nicotine extraction

Natural insecticide/pesticide extraction



SUPERCRITICAL CARBON DIOXIDE



Thomas Swan & Co., Ltd.

Licence, P. *et al* *Green Chem.* 5, 99 (2003)



GAS EXPANDED LIQUIDS (GXLs)

Combination of a classical solvent and supercritical solvent (usually scCO_2) – adjustable polarity;

