

CONDITIONAL CONTROL OF GENE EXPRESSION IN THE MOUSE

Mark Lewandoski

One of the most powerful tools that the molecular biology revolution has given us is the ability to turn genes on and off at our discretion. In the mouse, this has been accomplished by using binary systems in which gene expression is dependent on the interaction of two components, resulting in either transcriptional transactivation or DNA recombination. During recent years, these systems have been used to analyse complex and multi-staged biological processes, such as embryogenesis and cancer, with unprecedented precision. Here, I review these systems and discuss certain studies that exemplify the advantages and limitations of each system.

CELL AUTONOMOUS

A genetic trait in which only genotypically mutant cells show the mutant phenotype.

CELL NON-AUTONOMOUS

A cell non-autonomous trait is one in which genotypically mutant cells cause other cells (regardless of their genotype) to show a mutant phenotype.

Section of Genetics of Vertebrate Development, Laboratory of Cancer and Developmental Biology, National Cancer Institute—Frederick Cancer Research and Development Center, Frederick, Maryland 21702-1201, USA. e-mail: mlewandoski@ncifcrf.gov

MOUSE GENOMIC TECHNOLOGIES


Our ability to produce mice that carry altered genomic loci or foreign transgenes has facilitated the study of many biological processes, from embryogenesis to animal behaviour. However, not all biological processes can be accessed and studied by gene-inactivation or transgene-expression strategies. For example, many genes have several roles during embryogenesis and adulthood; knocking out these genes can result in an embryonic-lethal phenotype, which reflects the earliest non-redundant role of the gene, and precludes an analysis of its function in later development. Similarly, embryonic lethality due to the expression of a transgene renders it useless for the study of adult biology. Of the phenotypes that are caused by such genetic manipulations, many are difficult to interpret because they might be due to CELL-AUTONOMOUS, CELL-NON-AUTONOMOUS or systemic effects in the whole animal. To circumvent these limitations, mouse geneticists have long searched for experimental ‘switches’ that allow genes to be activated or silenced where and when the investigator chooses.

The ideal genetic switch results in low or zero basal gene expression when ‘off’ and high levels of gene expression when ‘on’. Furthermore, the switch should be reversible and specific for the target gene, and should not interfere with other cellular components or with general metabolism. The earliest attempts to develop such a genetic switch made use of single transgenes that

had promoters that could be induced by heavy metals, heat shock, interferon or steroids. These transgenes fell somewhat short of the ideal because they produced pleiotropic effects or showed high basal activity in the absence of induction¹.

However, the criteria for an ideal genetic switch have largely been met by binary transgenic systems, in which gene expression is controlled by the interaction of two components: an ‘effector’ transgene, the product of which acts on a ‘target’ transgene (BOX 1). Binary transgenic systems fall into two categories. In one, the effector transactivates transcription of the target transgene. In the other, the effector is a site-specific DNA recombinase that rearranges the target gene, thereby activating or silencing it. Both systems have their advantages and limitations, as discussed in this review. During recent years, the number of these binary systems has increased, and the more established systems, such as Cre/*loxP* and the TetR-based transactivators (see below), have been refined. Furthermore, these systems have been increasingly used to solve otherwise intractable problems in transgenic mice. Here, I review the current systems for conditional gene expression, focusing on specific examples that illustrate details in experimental design that might be useful to those considering the use of such technologies.

Box 1 | Binary transgenics in the mouse



In the ideal conditional system, investigators should be able to activate or silence a transgene where and when they choose. The breakthrough model, on which all the systems in this review are based, is a binary system in which gene expression is controlled through the interaction of two components: the product of an ‘effector’ transgene acting on a ‘target’ transgene. The effector transgene alone should not affect the expression of endogenous genes; its expression should therefore be innocuous. Control of target transgene expression should be wholly conditional on the presence of the effector. So, expression of the target transgene should only occur in doubly transgenic F₁ progeny derived from mouse lines that carry either the target transgene or the effector transgene, or in the presence of an exogenous inducer that allows the appropriate interaction of the effector on the target. These criteria have been largely met by developing binary systems with components derived from systems that are too evolutionarily distant from the mouse to have any unintended interactions with the mouse genome. In the example shown (see figure), the effector gene product–target gene interaction results in target activation but, in some cases (such as when using the site-specific recombinases, Cre or Flp), the effector can also inactivate the target transgene. In most of the inducible binary systems, the effector gene product–target gene interaction depends on the presence of an exogenously added inducer, but withdrawal of an inducer can also result in effector action (see text for more details). TSP, tissue-specific promoter.

Transcriptional transactivation

In the first binary systems, transgene transcription was transactivated by viral proteins^{2,3}, but these systems were not further used because of high basal levels of target-gene expression in the absence of transgene transactivation², or because expression of the transactivator caused tumorigenesis⁴. One exception to this has been the VP16 protein of herpes simplex virus, which has been used to control the transcription of target transgenes that contain the VP16 response element^{5–7}. However, the use of this system is limited because VP16 is lethal when expressed in pre-implantation embryos, presumably because of the SQUELCHING

SQUELCHING
The titration of interacting molecules that are out of equilibrium by the overexpression of an interacting regulatory partner molecule, which can result in pleiotropic effects.

due to its interaction with mouse transcription factors^{8,9}. Squelching might also occur when chimeric transactivators that contain the VP16-activating domain are highly expressed, and *in vitro* solutions to this problem have been reported^{9,10}. But, generally, such interactions have not impeded the use of these chimeric transactivators in transgenic mouse studies¹¹ (described in more detail below) (TABLE 1).

Transcriptional transactivation is used more widely than DNA recombination to bring about transgene activation in mice, mainly because transgene activation by DNA recombination is irreversible, which results in permanent transgene activation in any cell lineage. By contrast, transcriptional transactivation occurs only when the transactivator is present, and, if an inducible system is used, then transgene activation can be reversed at a given time (see below), which provides the investigator with more control over transgene expression.

TetR-based systems. The most widely used binary transcription transactivation systems are the tetracycline-dependent regulatory systems developed by Manfred Gossen and Hermann Bujard¹² (see link to **Hermann Bujard’s trouble-shooting guide**). In these systems, the effector is a fusion of sequences that encode the VP16 transactivation domain and the *Escherichia coli* tetracycline repressor (TetR) protein, which specifically binds both tetracycline and the 19-bp operator sequences (*tetO*) of the *tet* operon in the target transgene, which results in its transcription (FIG. 1). There are two versions of this system. In the original system, the tetracycline-controlled transactivator (tTA) cannot bind DNA when the inducer is present (‘tet-off’) (FIG. 1a), whereas in a modified version, the ‘reverse tTA’ (rtTA) binds DNA only when the inducer is present (‘tet-on’)¹³ (FIG. 1b). The current inducer of choice is doxycycline (Dox) because of its low cost, commercial availability, and because it efficiently activates rtTA and inactivates tTA at doses that are well below cytotoxic levels^{11,13}.

The principal difference between tTA and rtTA is the kinetics of target transgene induction. In the tTA system, transgene suppression requires the continuous administration of Dox; induction occurs on Dox removal and is, therefore, dependent on the rate of Dox clearance. In adult transgenic mice, induction can take from 24 hours to 1 week, depending on the tissue of interest¹⁴. However, in the rtTA system, the addition of Dox can rapidly induce transgene expression, even within an hour¹⁵, whereas subsequent repression depends on the clearance of Dox from the tissue of interest.

Although Dox can cross the placenta and efficiently regulate gene expression during embryogenesis, these systems have been underused to study embryogenesis *per se* and have mostly been used to suppress the embryonic expression of lethal transgenes. In one study by Lee *et al.*¹⁶, which possibly subjected the system to the most stringent test, tTA was used to control the expression of a target transgene that encoded the cell-autonomous, lethal DIPHTHERIA TOXIN A SUBUNIT (DTA). Mouse lines that carried the *tetO*–DTA target transgene were produced at a ten-fold lower frequency compared with this lab’s production

Table 1 | Examples of tissue-specific tTA and rtTA mice*

Tissue-specific promoter	Tissue of expression	Transactivator	Target gene	Reference
<i>Lap</i> (<i>Cebpb</i>)	Liver	tTA	Luciferase	14
<i>Myh6</i>	Heart	tTA	Diphtheria toxin A	16
<i>Ednrb</i>	Embryonic neural crest	tTA, rtTA	<i>Ednrb</i>	17
<i>Camk2a</i>	Hippocampus, striatum	tTA	Mutated calcineurin	19
<i>Camk2a</i>	Hippocampus, cortex, striatum, olfactory bulb, cerebellum	rtTA	Calcineurin inhibitor	18
<i>MMTV-LTR</i>	Submandibular gland	tTA	<i>SV40</i>	20
<i>Tyr</i>	Melanocytes	rtTA	<i>H-ras</i> ^{V12G}	21
<i>Eμ-Sr</i>	Haematopoietic cells	tTA	<i>Myc</i>	22
<i>MMTV-LTR</i>	Haematopoietic cells	tTA	<i>Bcr-Abl1</i>	23
<i>Camk2a</i>	Hippocampus, cortex, striatum	tTA	<i>Hdh</i>	24
<i>Lap</i>	Liver	tTA	<i>Nfkb1a</i> , luciferase	28
<i>Fabp</i>	Small intestine, colonic epithelium	rtTA	Cre recombinase	34
<i>RB</i>	Retina, cerebellum, monocytes	rtTA	Cre recombinase	116
<i>Wap</i>	Mammary gland	rtTA	Cre recombinase	116

*These are just a few examples of many such mouse lines to illustrate the range of tissues in which tissue-specific transactivator expression can be achieved. (*Bcr-Abl1*, breakpoint cluster region-Abelson protein1; *Camk2a*, Ca²⁺/calmodulin-dependent protein kinase II, alpha; *Cebpb*, CCAAT/enhancer-binding protein, beta; *Ednrb*, endothelin receptor-B; *E μ -Sr*, immunoglobulin heavy chain enhancer-SR α promoter; *Fabp*, fatty-acid-binding protein; *Hdh*, huntingtin; *H-ras*, Harvey rat sarcoma virus oncogene; *Lap*, liver activator protein; *Myh6*, myosin heavy chain, cardiac muscle, alpha; *MMTV-LTR*, mouse mammary tumour virus-long terminal repeat; *Myc*, myelocytomatosis oncogene; *Nfkb1a*, nuclear factor of kappa light chain gene enhancer in B-cells inhibitor, alpha; *RB*, retinoblastoma; rtTA, reverse tetracycline-controlled transactivator; *SV40*, simian virus 40; tTA, tetracycline-controlled transactivator; *Tyr*, tyrosinase; *Wap*, whey acidic protein.)

of other transgenic lines, presumably because the leaky expression of the transgene caused cell ablation and death¹⁶. However, the *tetO*-DTA transgene was regulated properly by tTA in the lines that were established, and its specific induction in the hearts of adult mice caused cardiomyopathies¹⁶.

This system has been effectively used during embryogenesis to study endothelin receptor type B (*Ednrb*) signalling¹⁷, which is required for the normal development of neural-crest derivatives. Activation of an *Ednrb* target transgene through rtTA, or its rapid inactivation through tTA, within hours of Dox administration, showed that *Ednrb* is required for neural-crest migration during embryonic days 10–12.5. To achieve this tight regulation and to make all *Ednrb* expression dependent on Dox regulation, gene targeting was used to produce three mouse lines in which the wild-type *Ednrb* allele was replaced with sequences that encoded one of the two transactivators or the *Ednrb*^{tetO} target allele. The SELECTION CASSETTE used to target the *Ednrb* locus, which allowed targeted embryonic stem (ES) cells to be selected, was designed so that it could be removed in transgenic mouse lines by Cre-mediated recombination (as described in more detail below). Its removal was required because the presence of the cassette reduced the expression of the transactivator-encoding transgenes. However, on its removal from the *Ednrb*^{tetO} allele, an unacceptable increase in the basal activity of this allele occurred. As a result, an *Ednrb*^{tetO} allele that still contained the selection cassette was used in experimental crosses. This study exemplifies the importance of having a

versatile experimental design that allows targeted alleles to be altered after transgenic mice are produced; the experiment would not have succeeded had the investigators failed to design their selection cassettes for removal, or had they removed them in ES cells before producing transgenic mice.

Theoretically, it is possible to induce and then silence *Ednrb*^{tetO} expression with only tTA or rtTA, but the kinetics of Dox depletion is too slow for either this induction (through tTA) or repression (through rtTA) to be useful in embryological studies. However, in studies of cancer, neurological disease and memory^{18,19}, such kinetics are not an impediment, and the ability to reverse target-gene expression has produced impressive results. Inducing and then silencing oncogene expression with either tTA or rtTA in both solid tumours^{20,21} and leukaemias^{22,23} has shown that, in most cases, the oncogene that induces tumorigenesis is also required for tumour maintenance. In a mouse model of **Huntington disease** that relied on the tTA activation of a mutant *huntingtin* transgene, Dox treatment over 16 weeks resulted in transgene repression, and the reversal of the neuropathology and behavioural abnormalities that characterize the disease²⁴. These studies indicate that future cancer or Huntington disease therapies that target the activity of a single molecule might be clinically effective.

The 'tet-on' and 'tet-off' systems continue to be modified with different transactivator domains, trans-repression domains, codon-usage modifications, and variations to the configuration of *tetO* sequences^{10,11,25,26} (see also links to **Knoll GmbH** and **ClonTech**). One useful variant allows two target transgenes to be co-regulated

DIPHThERIA TOXIN A SUBUNIT (DTA). Diphtheria toxin consists of two subunits, A and B. The B-subunit binds receptors on the surface of the target cell, facilitating the entry of the A-subunit, which ADP-ribosylates elongation factor 2, thus preventing protein synthesis. The gene that encodes the A-subunit is often used as a cell-autonomous toxin in transgenic ablation experiments.

SELECTION CASSETTE
A DNA fragment that contains a transgene, which, when expressed, allows the selection of a subset of cells that have integrated the DNA fragment into their genomes.




Figure 1 | The tetracycline-responsive regulatory system for transcriptional transactivation. a | The effector is a tetracycline-controlled transactivator (tTA) of transcription that consists of a chimeric construct of the *Escherichia coli* Tn10 *tetR* gene (purple) and the VP16 transactivation domain (orange). In the absence of the inducer, doxycycline (Dox), tTA dimers specifically bind to seven tandemly repeated 19-bp *tetO* sequences (*tetO*₇), thereby activating transcription from a minimal promoter (TATA) and driving expression of the target transgene that encodes the gene of interest (target ORF). When bound to Dox, tTA undergoes a conformational change and cannot bind *tetO* sequences. **b** | In the reverse tTA (rtTA) system, the *tetR* gene has been mutated so that it binds *tetO* sequences and activates transcription only in the presence of Dox. **c** | By flanking *tetO* sequences with two minimal promoters, two target transgenes can be activated by the tTA (or rtTA) systems. In the example shown, tTA drives transcription of a reporter gene, as well as the target gene of interest. (bp, base pair; ORF, open reading frame; pA, polyadenylation site; *tetO*, the 19-bp operator sequences of the *tet* operon; *tetR*, tetracycline repressor; TSP, tissue-specific promoter; VP16, transactivation domain of the herpes simplex virus protein VP16.)

POSITION EFFECTS
The effect of the local chromosomal environment on the levels or pattern of transgene expression, possibly because of local chromatin configuration or nearby *cis*-acting regulatory elements.

by flanking the *tetO* sequences with two minimal promoters²⁷ (FIG. 1c). This enables transactivation in transgenic mice to be monitored when one promoter drives a reporter gene, such as luciferase²⁸. In another variation, the DNA-binding specificity of the transactivator has been modified so that it will discriminate between altered *tetO* sequences, allowing an investigator to switch transactivation between two different target transgenes by altering Dox levels²⁹. Finally, a random mutational screen has generated rtTA variants with increased Dox sensitivity. These variants should mediate a greater responsiveness to Dox in those mouse tissues, such as

brain, to which Dox has limited access²⁵. Although these variants have yet to be used in transgenic mouse studies, the continuing evolution of these systems *in vitro* will soon affect the precision with which we will be able to dissect biological problems *in vivo*.

GAL4-based systems. In *Saccharomyces cerevisiae*, the transcriptional activator Gal4 directs the transcription of Gal4-responsive genes by binding to upstream activator sequences (UASs). The Gal4/UAS system was used to direct transgene activation in the first non-viral binary system to be used in mice³⁰ (FIG. 2a). This approach has been used to study the role of **Sonic hedgehog** (Shh) in the dorsal–ventral patterning of the mouse central nervous system (CNS). In previous studies, Shh, which is normally expressed along the ventral midline of the CNS, had been ectopically expressed along the dorsal midline with a single *Shh* transgene driven by *Wnt1* regulatory sequences (*Wnt1–Shh*)³¹. However, because of an embryonic-lethal phenotype, only a limited number of founder embryos could be analysed. Furthermore, the phenotype varied between these founder embryos presumably because of POSITION EFFECTS. These constraints were bypassed by activating a UAS–*Shh* target transgene with a Gal4 effector transgene that was expressed under the control of a *Wnt1* promoter (*Wnt1–GAL4*) to generate many mutant embryos with a highly reproducible phenotype³². It is noteworthy that the initiation of β-galactosidase activity through the transactivation of UAS–*lacZ* by *Wnt1–GAL4* was delayed by ~24 hours relative to that from a single *Wnt1–lacZ* transgene, a time delay which could be significant in some developmental studies.

One example of an inducible Gal4-based system is the GLVP system, which is named after the constituent parts of the chimeric effector protein (FIG. 2). Because of the ligand-binding domain of the GLVP transactivator, the target transgene can be induced by the synthetic steroids, RU486 or ZK98.734 (FIG. 2b). Recently, the topical application of ZK98.734 to mice that express GLVP specifically in the epidermis was used to activate transforming growth factor-β1 (*Tgfb1*) expression in the skin³⁵. Previous reports using single transgenes had given contradictory results as to whether Tgf-β1 inhibited or stimulated epidermal growth³⁶. By temporally controlling *Tgfb1* expression, locally and in a dose-dependent manner with topical applications of ZK98.734, Tgf-β1 was shown to inhibit epidermal growth³⁵. In general, the use of the GLVP system has been limited to controlling gene expression in adult mice^{33,35,37}, as the abortigenic properties of these synthetic steroids, which act as antiprogestins, limit their use in embryos; although their co-injection with progesterone has been shown to rescue embryonic development (D. R. Roop, personal communication).

In summary, the advantages of using transcriptional transactivation to conditionally control gene expression are: the reversibility of the expression of the target gene; the sensitivity of transgene activation levels to




Figure 2 | Transactivation based on the Gal4/UAS system. **a** | The effector is the Gal4 transcriptional transactivator (blue circles) from *Saccharomyces cerevisiae*, which binds as a homodimer to a 17-bp upstream activating sequence (UAS). The target transgene is activated by Gal4 homodimers that bind to four tandemly repeated UASs placed immediately upstream of a minimal promoter (TATA). **b** | GLVP is an inducible Gal4/UAS-based system. In the original version, the transactivator is a chimeric construct (GLVP) that consists of the herpes simplex virus VP16 activation domain (orange), the Gal4 DNA-binding domain (blue) and the ligand-binding domain of the progesterone receptor (LBD Δ 42) (green), mutated such that it fails to bind progesterone but can bind antiprogestins, such as RU486. GLVP is believed to be sequestered in the cytoplasm by its binding to heat-shock proteins (HSPs) 70 and 90. In the presence of RU486, GLVP is released, becomes localized to the nucleus and transactivates the UAS-bearing target transgene. Several versions of the transactivator exist owing to the modification of different domains¹⁴⁶ (see link to **GeneSwitch™**, Valentis, Inc.). (ORF, open reading frame; pA, polyadenylation site; TSP, tissue-specific promoter.)

inducer concentration; and the ability to control the expression of more than one target transgene. These features make this strategy well suited to gain-of-function experiments, in which phenotypes are generated by transgene overexpression or ectopic expression. A potential limitation is that the binary nature of these systems compounds twofold the basic problem of any transgenic experiment: the susceptibility of transgene regulatory elements to position effects. To achieve correct regulation, the transgenes could be flanked with INSULATOR SITES^{26,33} or inserted into a known locus that provides low basal and high activated expression³⁴, but often the best strategy is to test different combinations of effector and target lines.

INSULATOR SITE
DNA sequence that blocks the interaction between *cis*-acting regulatory elements. These sites are sometimes used to protect transgenes from genomic position effects.

CONSERVATIVE DNA RECOMBINATION
A DNA recombination reaction in which there is no net change in base pairs between the products and the reactants.

Site-specific DNA recombination

Two members of the integrase family of site-specific recombinases, Cre from bacteriophage P1 (REF. 38) and Flp from *Saccharomyces cerevisiae*³⁹, are used at present to conditionally control gene expression by site-specific DNA recombination. Cre and Flp recombinases catalyse a CONSERVATIVE DNA RECOMBINATION event between two 34-bp recognition sites (*loxP* and *FRT*, respectively) that differ in primary sequence but share secondary structure (FIG. 3). Because these recombinases do not require accessory proteins or high-energy cofactors for their activity, they are suitable for use in the foreign milieu of the mouse cell. However, Flp has been modified to alter its temperature optimum for improved activity at 37 °C — the body temperature of mice⁴⁰. Also, many Cre variants now being used have been modified with a heterologous nuclear localization sequence (NLS)^{41,42}, which has been found to increase Cre-mediated recombination in some *in vitro* assays⁴¹, but not in others⁴³. As wild-type Cre contains basic peptides that direct it to the mammalian nucleus⁴³, the advantages of using such modified Cre variants *in vivo* is questionable.

Originally, the expression of *cre* or *FLP* in mice was expected to be innocuous because it was thought that the mouse genome would not harbour endogenous *loxP* or *FRT* sites. However, these recombinases have been shown *in vitro* to mediate recombination between degenerate *loxP/FRT* sequences (pseudosites) with reduced efficiency^{44,45}, and pseudo-*loxP* sites have been identified in the mouse genome⁴⁴. A recent study reported that high levels of *cre* expression in cultured cells leads to growth arrest and DNA damage, indicating that recombination between these pseudo-sites might occur *in vitro*⁴⁶. In one case, *cre* expression in the male germ line resulted in chromosomal aberrations and sterility, indicating that similar interactions might sometimes occur *in vivo*⁴⁷. However, the widespread expression of *cre* in many other transgenic mouse lines (see link to the **Cre transgenic database**), including both male and female germ lines^{48,49}, attests to the rare occurrence of such interactions *in vivo*.

The ability to use Cre- or Flp-mediated DNA recombination to invert or delete DNA fragments *in cis*, or to switch DNA fragments *in trans* has many applications beyond conditional gene expression, which include chromosome engineering (see accompanying review by Yu and Bradley on p780 of this issue) or the recycling of selectable markers in cultured cells⁵⁰. Here, I focus on the applications of this system for activating or silencing gene expression by deleting DNA fragments that have been flanked by directly repeated *loxP* or *FRT* sites (so called ‘floxed’ or ‘flrtd’ alleles, respectively) (FIG. 3). The *Cre/loxP* system is the most frequently used owing to historical contingency, as it was the first well-developed recombination system. Many more effector and target transgenic mouse lines now exist, to which newly generated mouse lines can be crossed; however, the general strategies are the same for the Flp/*FRT* system.

Tissue-specific knockouts. The ability to inactivate an endogenous gene in the mouse in a temporally and spatially controlled manner is not only useful for circumventing early lethal phenotypes, but also allows biological questions to be addressed with exquisite accuracy. In such studies, standard gene-targeting techniques are used to produce a mouse in which an essential region of a gene of interest is floxed, so that tissue-specific *cre* expression results in the inactivation of this allele. Before recombination, the conditional allele should have wild-type activity. In most cases, *loxP* sites are placed in introns, but have sometimes been inserted in 5'- (REFS 51,52) or 3'-flanking regions^{53,54} without compromising gene expression. In one published exception⁵⁵, a floxed glucokinase allele resulted in a slight reduction (10–15%) in enzyme activity, which might have been due to extra linker sequences adjacent to the *loxP* site (M. Magnuson, personal communication).

By crossing a mouse line with a conditional allele to an effector mouse line that expresses *cre* in a tissue-specific manner, progeny are produced in which the conditional allele is inactivated only in those tissues or cells that express *cre*. Since the first published tissue-specific knockout⁵⁶, the use of this strategy has steadily increased (see link to the [Cre transgenic database](#)) (TABLE 2), and there are now many instructive examples of its application.

Table 2 | **Examples of tissue-specific *cre*-expressing mice***

Tissue-specific promoter	Tissue/cell of expression	Floxed target gene	Reference
<i>Lck</i>	T cells	DNA polymerase-β	56
<i>Cryaa</i>	Eye lens	<i>SV40</i> (activated)	82
<i>Ins2</i>	Pancreatic β-cells	<i>Gck</i>	55
<i>Alb</i>	Liver	<i>Gck</i>	
<i>Myog</i>	Skeletal muscle	Diphtheria toxin A [†]	83
<i>KRT5</i>	Epidermis	β1-integrin	54
<i>Nes</i>	Neuronal cells	<i>Mecp2</i>	53
<i>Ins2</i>	Pancreatic β-cells	<i>hGH</i> [‡]	79
<i>Gcg</i>	Pancreatic α-cells	<i>hGH</i> [‡]	
<i>Ppy</i>	Pancreatic β-cells	<i>hGH</i> [‡]	
<i>Pdx1</i>	Pancreatic α-cells	<i>hGH</i> [‡]	
<i>En2</i>	Mid/hindbrain	<i>lacZ</i> [‡]	75
<i>Wnt1</i>	Neural crest	<i>lacZ</i> [‡]	78
<i>Camk2a</i>	Forebrain	<i>Hdh</i>	51
<i>Nes</i>	Branchial arch	<i>Fgf8</i>	65
<i>Msx2</i>	Apical ectodermal ridge of limb bud	<i>Fgf8</i>	66
<i>Pax6</i>	Retina	<i>Pax6</i>	137

*These are just a few examples of many such mouse lines to illustrate the range of tissues in which tissue-specific *cre* expression can be achieved. †In these cases, Cre-mediated recombination resulted in gene activation. ‡In these cases, Cre-mediated recombination resulted in gene inactivation. (*Alb*, albumin; *Camk2a*, Ca²⁺/calmodulin-dependent protein kinase II, alpha; *Cryaa*, crystallin alpha A; *En2*, Engrailed 2; *Gcg*, glucagon; *Ins2*, insulin II; *Lck*, lymphocyte-specific tyrosine kinase; *Fgf8*, fibroblast growth factor 8; *hGH*, human growth hormone; *Hdh*, huntingtin; *Mecp2*, methyl-CpG-binding protein 2; *Msx2*, msh-like homeobox gene 2; *Myog*, myogenin; *Nes*, nestin; *Pax6*, paired box gene 6; *Pdx1*, pancreatic and duodenal homeobox factor 1; *Ppy*, pancreatic polypeptide; *SV40*, simian virus 40; *Wnt1*, wingless-related MMTV integration site 1.)

A selectable marker cassette (such as that containing the neomycin resistance gene, *neo*), is routinely used to select for targeted ES-cell clones and is necessarily part of the initial floxed allele. However, the placement of *neo* in introns, or in 5'- or 3'-flanking or untranslated




Figure 3 | Controlling gene expression by DNA recombination. **a** | The 34-bp *loxP* and *FRT* sites each consist of two 13-bp inverse repeats (black) that flank an 8-bp core sequence (red). This core sequence confers directionality to these sites (red arrows). **b** | Dimers of Cre or Flp (pink) catalyse *in cis* the conservative recombination between two directly repeated *loxP* or *FRT* sites (red arrowheads), resulting in the formation of a synaptic structure, the excision of region B and the juxtaposition of regions A and C. If region B is an essential region of a gene, then the recombination event results in gene inactivation. Recombination can also activate gene expression. For example, transcription from a promoter in region C could fail to reach protein-coding sequences in region C if polyadenylation sites exist in region B; excising region B would therefore activate transcription of region C. **c** | Use of three *loxP* sites and a 'partial' recombination event (between *loxP* sites labelled 1 and 2) to remove a neomycin resistance (*neo*) selection cassette and leave an essential region floxed. The other two products of recombination between *loxP* sites 2 and 3, and between 1 and 3 are also shown. (pA, polyadenylation site; TSP, tissue-specific promoter.)

HYPOMORPHIC ALLELE

An allele that results in a reduction, but not the elimination, of wild-type levels of gene product or activity, often causing a less severe phenotype than a loss-of-function (or null) allele.

ALLELIC SERIES

An array of possible mutant forms of a gene, which usually cause several phenotypes.

SEMI-DOMINANT ALLELE

An allele that causes an intermediate, incompletely dominant phenotype in heterozygotes.

ROSA26

A genetic locus that was originally identified by gene-trapping technology, which is constitutively transcribed.

regions, can interfere with the expression of the floxed allele, compromising its wild-type activity in the absence of Cre-mediated recombination (TABLE 3). In some cases, it has been determined that cryptic splice sites in *neo* interfere with normal splicing and therefore reduce wild-type mRNA levels^{57–59}. A common solution to this problem, carried out in the first tissue-specific knockout experiment, is to use three *loxP* sites, positioned such that *neo*, as well as the essential region of the gene of interest, is floxed⁵⁶. A partial Cre-mediated recombination event then removes *neo*, leaving the essential region floxed (FIG. 3c). This is most often achieved *in vitro* in ES cells by transiently introducing a *cre*-expression cassette⁵⁶. However, it can also be achieved *in vivo* by injecting zygotes with a *cre*-expression cassette⁵⁹, or by breeding mice that carry floxed alleles to mice that mosaically express *cre* in the germ line, such that recombination occurs between two of the three *loxP* sites at a reasonable frequency^{59,60}.

However, because the inclusion of *neo* can generate a HYPOMORPHIC ALLELE, retaining it can often be advantageous, because such an allele, in combination with the null and conditional alleles, can be very informative^{57,58,61,62}. One approach, the allelogenic strategy, generates a hypomorphic allele as part of an ALLELIC SERIES from one mouse line by floxing an essential region of the gene of interest and then flrting the *neo* cassette so that it can be removed by FLP-mediated recombination⁵⁷. Three mouse lines can be generated in such an approach. The original line expresses a hypomorphic allele (if *neo* interferes with gene expression); one line expresses a floxed allele, which can be inactivated in a tissue-specific gene manner and is generated by mating

the original line to a *FLP*-expressing mouse; and finally, one line carries a null allele, generated by breeding the original line to a *cre*-expressing mouse. Such an allelic series at the fibroblast growth factor 8 (*Fgf8*) locus⁵⁷ has been instrumental in studying the role of *Fgf8* in gastrulation⁶³, left–right axis determination⁶⁴, branchial arch development⁶⁵ and limb development⁶⁶. An allelic series that includes any desirable genomic alteration can also be generated by introducing specific mutations into the original targeting cassette; the removal of *neo* by Cre-mediated recombination would then leave the specific mutation⁵⁸. Using this approach, an allelic series has been generated at the Fgf receptor 1 (*Fgfr1*) locus, including SEMI-DOMINANT and hypomorphic alleles⁶¹.

To correctly interpret a phenotype caused by tissue-specific Cre-mediated gene inactivation, it is often essential to know in which cells recombination has taken place. This is usually done by monitoring gene expression at the RNA or protein level, but an alternative is to include a reporter construct (such as one that encodes *lacZ*) downstream of the floxed region, so that recombination both inactivates the target gene and activates the reporter⁶⁷. However, care must be taken to ensure that, like *neo*, the reporter does not compromise the expression of the conditional allele before recombination.

Finally, the success of these experiments depends on the availability of tissue-specific regulatory elements to drive *cre* expression. Controlling *cre* expression is one of the most rigorous tests of the tissue specificity of any regulatory element, as nonspecific expression could render a *cre* transgene useless because of the effective irreversibility of the recombination event. Nevertheless, even if the appropriate Cre line is not available, it is logical for investigators to produce floxed alleles rather than standard ‘knockouts’. This is because they can still generate the recombined null allele by using one of the many Cre ‘deletor’ strains (see link to the [Cre transgenic database](#)) and still have mice that carry the floxed allele for a future date when the right *cre*-expressing mouse comes along.

Cell-lineage analysis. Two different approaches have been used to produce mice in which a histological marker can be activated by Cre-mediated recombination in nearly any tissue, thereby reporting Cre activity. Such mice are useful not only for characterizing a newly made *cre*-expressing line, but also for analysing cell lineages, as described below. In one approach, the production of such reporter mice was achieved by using gene targeting to insert Cre-activatable (as described below) reporter constructs that encode β -galactosidase^{68,69}, or the enhanced fluorescent proteins of the green⁷⁰, yellow or cyan varieties⁷¹, into the *ROSA26* locus, which produces constitutive transgene expression. In the other approach, ES clones that contain *cre*-reporter constructs were screened *in vitro* for widespread expression before the production of mice. In this manner, the *Z/AP*⁷² and *Z/EG*⁷³ reporter lines were produced, in which Cre-mediated recombination causes a switch from the expression of the *lacZ* gene to

Table 3 | Loss of target-gene function due to *neo* cassette

Floxed gene	Location and orientation* of <i>neo</i>	Reference
<i>Fgf8</i>	Intron; same orientation	57
<i>N-Myc</i>	Intron; same orientation	58
<i>Fgfr1</i>	Intron; same orientation	61
<i>Gck</i>	Intron; same orientation	55
<i>Pitx2</i>	Intron; same orientation	62
<i>Pck1</i>	Intron; same orientation	147
<i>Krt1-14</i>	Intron; same orientation	90
<i>Krt1-10</i>	Intron; same orientation	89
<i>Fgfr3</i>	Intron; opposite orientation	148
<i>Hnf4</i>	Intron; opposite orientation	149
<i>Brca1</i>	Intron; opposite orientation	59
<i>Slc2a4</i>	3' UTR	150
<i>Fgf8</i>	3' UTR	67
<i>Nodal</i>	3' UTR	151
<i>Scap</i>	5' UTR	152
<i>Bcl2l</i>	5' promoter region	153

*Orientation of *neo* cassette relative to transcription of floxed gene. (*Bcl2l*, B-cell lymphoma; *Brca1*, breast cancer gene 1; *Fgf8*, fibroblast growth factor 8; *Fgfr3*, fibroblast growth factor receptor 3; *Gck*, glucokinase; *Hnf4*, hepatocyte nuclear factor 4; *Krt1-x*, keratin complex 1, gene x; *N-Myc*, neuroblastoma myc-related oncogene; *Pck1*, phosphoenolpyruvate carboxykinase; *Pitx2*, pituitary homeobox 2; *Scap*, Srebp cleavage-activating protein; *Slc2a4*, solute carrier family 2, member 4 (also called *Glut4*); *Srebp*, sterol regulatory element-binding protein; UTR, untranslated region.)




Figure 4 | Fate mapping the descendants of a gene-expression domain. In this example, a tissue-specific promoter (TSP) (for example, the *Wnt1* promoter) controls the expression of a *cre* or *FLP* transgene. **a** | The *cre* (or *FLP*) transgene is not expressed outside the dorsal neural tube. A reporter transgene is transcribed from a ubiquitous promoter (UP), but no functional gene product is produced outside the dorsal neural tube because Cre (or Flp)-mediated recombination is required to remove the floxed (or flirited) Stop sequences that separate the promoter from the reporter coding sequences (red arrowheads, *loxP* or *FRT* sites). **b** | Cre activity in the dorsal neural tube (red) results in the removal of the Stop sequence and reporter activation (blue). **c** | Migrating neural-crest cells (blue arrow) no longer express *cre* or *FLP*, but the reporter remains activated (blue).

that for alkaline phosphatase or enhanced green fluorescent protein, respectively. The availability of different reporter lines is useful because recombination can sometimes be locus dependent⁷⁴.

These reporter lines are useful for linking gene expression to the fate of specific cell lineages because activation of one of these reporter genes indicates that Cre activity is present in that cell or was present in a progenitor cell (FIG. 4). This strategy has been used to fate map the mid-hindbrain border⁷⁵, neural-crest cell derivatives^{76–78}, pancreatic islet cells⁷⁹ and memory T cells⁸⁰. In such experiments, care must be taken to ensure that Cre is not active in a second lineage that is then erroneously assumed to be related to the first lineage. To overcome this problem, it might be useful to insert *cre* into the locus of the gene of interest so that it is free of position effects. As such insertions usually inactivate the gene of interest, one can also compare the fate map of phenotypically wild-type animals that are heterozygous for the *cre* allele, to mutant animals that are homozygous for the *cre* allele. Such an approach has been used to analyse the role of *Hoxb1* on the fate of hindbrain derivatives⁸¹.

Cre-mediated gene activation. The first demonstration of Cre activity in mice, in which a transgene was activated by Cre-mediated recombination⁸², provides

a model for this general strategy. Activation occurs by Cre removing a ‘Stop’ DNA fragment, which prevents transcription and translation because it contains polyadenylation sequences, false translation initiation codons and/or splice donor sequences. Alternatively, recombination can restore an open reading frame that is interrupted by floxed sequences. In this case, the *loxP* site that remains after recombination must be tolerated as part of the opening reading frame⁸³. Probably the best experimental approach to using this strategy is to design the target transgene for near-ubiquitous expression and then make use of the growing battery of mice that express *cre* in a tissue-specific manner (see link to the [Cre transgenic database](#)) to activate the target transgene. The creation of the *Rosa26* knock-in and Z/AP (or Z/EG) reporter mice, described above, exemplify how to ensure the otherwise difficult goal of achieving widespread transgene promoter activity in mice.

Investigators pursuing the Z/AP (or Z/EG) strategy should characterize the final ES clones *in vitro* before producing mice to ensure that ubiquitous expression of the target transgene is retained after Cre-mediated recombination. This will screen out the ~30% of ES clones in which the transgene is silenced after recombination for reasons that are at present unclear (C. Lobe, personal communication). Another advantage to using ES-cell technology to produce these target mice is that one can ensure that the transgene is present in a single copy, thereby avoiding a multiple array that occurs with the production of transgenic mice through PRONUCLEAR INJECTION. Cre-mediated recombination of such multiple arrays can result in chromosome loss⁸⁴, although it should be noted that this does not always occur⁸⁵.

As with the TetR-based system, the most rigorous demonstration of gene expression control has come from studies in which the activated target transgene expresses DTA driven by a tissue-specific promoter^{83,86,87}. Several labs are now producing universal ‘ablator’ mice in which the DTA target is driven by a ubiquitous promoter, such that different Cre effector mice can be used for tissue-specific ablation experiments.

Endogenous genes can also be altered by gene targeting to render their expression conditional on Cre activity. In this case, Cre-mediated recombination results in the excision of a floxed fragment, such as *neo*, targeted to the intron of a specific locus to prevent expression^{58,88}. So, in a design that is complementary to a tissue-specific knockout experiment, gene function can be studied by using Cre-mediated recombination to achieve tissue-specific gene repair. This approach has been used to model the human dominant skin-blistering disorders — epidermolytic hyperkeratosis and epidermolysis bullosa simplex. In these studies^{89,90}, mouse keratin genes that contain point mutations identical to human disease alleles were silenced with a floxed *neo*. Cre activation in the skin of these mice that was produced by the topical application of an inducer (described below), activated the mutant allele, resulting in a phenotype that models the human syndromes^{89,90}.

PRONUCLEAR INJECTION
One of two methods for producing a transgenic mouse line (the other method being through germ-line transmission of transgenic embryonic stem cells). In this approach, DNA is microinjected into the nucleus of a mouse zygote. The DNA integrates randomly, usually into one genomic locus, as a multiple array.




Figure 5 | Inducible Cre systems. **a** | Tissue-specific expression of the reverse transcriptional transactivator (rtTA) results in *cre* expression and recombination of the target gene, only in the presence of the inducer, doxycycline (Dox) (see FIG. 1 for details of the TetR-based system). **b** | The tissue-specific expression of a fusion gene that encodes Cre (pink circles) and a mutated ligand-binding domain (LBD) (green ovals) from the progesterone or the oestrogen receptor results in a chimeric protein that is thought to be sequestered in the cytoplasm by binding to heat-shock proteins (HSP) 70 and 90. In the presence of an inducer (RU486 or tamoxifen, respectively), the chimeric protein results in recombination of the target gene.

Animated online

Temporal control of Cre expression. The first inducible Cre mouse line used the *Mx1* promoter, which can be activated by injecting mice with interferon- α or interferon- β or with synthetic double-stranded RNA (polyinosinic-polycytidylic acid)⁹¹. The use of these mice has been limited to regulating gene expression in the liver^{92–95} and immune system^{96–98} where recombination efficiency is high^{91,94}.

Overlaying temporal regulation with promoter specificity has been achieved in two ways: by combining the *Cre/loxP* system with the TetR-based systems or by using Cre variants that function only in the presence of an exogenous inducer (FIG. 5). In the latter approach, the Cre-coding region is fused with a mutated ligand-binding domain (LBD) from either the progesterone^{99,100} or the oestrogen receptor^{101–106}. These mutated LBDs will bind and become localized to the nucleus only in the presence of the synthetic steroids — RU486 or tamoxifen, respectively — but not in the presence of endogenous steroids. High recombination efficiencies with these Cre-LBD fusion proteins have been obtained in mouse skin^{107,108}, but, in general, recombination mediated by such fusion proteins has been mosaic and dependent on tissue type^{99,103–104} despite nearly toxic doses of inducer^{102,103,108}. Furthermore, the doses required for induction *in utero* are perilously close to abortigenic levels¹⁰². Nevertheless, these systems have been successfully applied to addressing several biological questions^{89,109,110}. In one elegant study, boundaries that define specific structures in the developing limb were fate mapped at different embryological time points by tamoxifen-

induced, Cre-mediated activation of a *lacZ* reporter gene¹¹¹. New progesterone and oestrogen receptor-based Cre-LBD fusion proteins with enhanced sensitivities to their respective inducers are likely to increase the usefulness of this approach^{112,113}. One of these — Cre-ERT2 — has been used to inactivate the mouse retinoid X-receptor- α in mouse adipocytes¹¹⁴ and epidermis¹¹⁵.

Several groups have combined the TetR-based system with *Cre/loxP* so that Dox controls *cre* expression. As recombination results in a permanent genomic change, monitoring Cre-mediated excisions in such transgenic mice is a sensitive assay for the basal, uninduced, expression of the *tetO*-containing promoter from which *cre* is expressed (FIG. 5a). Inducing *cre* expression with Dox in the rtTA system results in tighter regulation of *cre* expression (although not without basal expression)^{34,116,117} in comparison with the tTA system¹¹⁸. Using this essentially tertiary transgenic system complicates mouse husbandry and increases the risk that position effects might occur, although this can be remedied by using a single Cre-rtTA construct^{116,117}. It is also helpful to prescreen *tetO/Cre* floxed target mice to exclude mouse lines in which leaky, uninduced recombination occurs³⁴. A *tetO-cre* transgene that responds well to induction and has low basal activity has been flrtd so that FLP-mediated recombination might be used to replace *cre* at this locus with other potential target genes³⁴.

An approach that obviates the need for producing Cre effector mice and that achieves temporal control of Cre-mediated recombination is the use of viral vectors,

such as adenovirus^{119–121} or herpes simplex¹²², to deliver *cre*. This approach is most useful in adult animals, although it is possible to deliver adenoviral vectors to embryos *in utero*¹²³. Limitations to this approach include complications due to possible host–virus interactions, mosaic recombination in the targeted tissue and variable infectivity between different tissues. Nevertheless, the viral delivery of *cre* to both activate transgenes and inactivate endogenous genes has been used in studies of liver function, cancer and the neurological basis of learning^{124–126}. Limitations can also be turned into advantages depending on the question being posed. For example, the adenoviral delivery of *cre* has been used to generate pancreatic β -cells in which the glucokinase gene is mosaically inactivated, showing that intercellular coupling has little to no function in glucose-stimulated insulin secretion¹²⁷.

The FLP/FRT system. Initial attempts to develop the FLP/FRT system in transgenic mice resulted in either no recombination¹²⁸ or mosaic recombination¹²⁹ because wild-type FLP and an FLP variant (FLP-F70L) have decreased enzyme stability at 37 °C (REF 130). However, a thermostable variant — FLPe — has been generated⁴⁰ that mediates recombination efficiencies *in vivo* resembling those of Cre¹³¹. All three FLP variants have their uses depending on whether an experiment calls for complete or mosaic recombination levels^{40,131}.

Although use of the FLP/FRT system is unlikely to be as widespread as that of the Cre/*loxP* system, it continues to be developed as a genetic tool and will be useful as a supplementary system. For example, FLP has been used in tissue-specific gene-inactivation studies¹³², FLP–LBD fusions have been generated^{133,134} and a *Rosa26*-based FLP reporter mouse has been produced¹³⁵. Recently, fate mapping through FLP-mediated recombination showed that *Wnt1* expression in the neural crest might define a subset of progenitor cells that eventually gives rise to specific types of precerebellar neurons¹³⁶. As the authors point out, FLP-based lineage mapping can now be used to analyse phenotypes caused by Cre-based, tissue-specific gene knockouts.

In summary, the effective irreversibility of site-specific DNA recombination makes systems such as Cre/*loxP* and FLP/FRT uniquely suited for tissue-specific gene-inactivation and cell-lineage studies during embryogenesis and in the adult. However, as the list of tissue-specific *cre* mice grows, we might find that in some cases high recombinase activity leads to chromosomal aberrations^{46,47}. Another limitation is shown by the recent demonstration that recombination efficiencies can be locus dependent⁷⁴, although other studies have shown the efficient recombination of three floxed alleles in the same cell¹³⁷.

The future

To ask increasingly more sophisticated biological questions, we will eventually need to individually control the expression of two or more genes in the same mouse line. For example, inactivating a ligand in a specific tissue, rescue of the resulting phenotype by transactivating

a constitutively active receptor in an adjacent tissue, and then characterizing the final phenotype by DNA-recombination-based fate mapping might be required. Therefore, systems such as the ecdysone-regulated gene switch¹³⁸ or the *lac* operator–repressor¹³⁹, might be increasingly used to address biological questions in transgenic mice. The ecdysone system is a transactivating system, based on a modified *Drosophila* steroid receptor that has been used to achieve high levels of target-gene induction in cultured cells. However, stocks of the ecdysteroid inducer — muristerone A — which is derived from the seeds of the Himalayan kaladana plant, have become depleted, limiting the use of this system in mice, although this barrier might be overcome with a recent report of alternative inducers¹³⁸. The *E. coli* *lac* operator–repressor system has been recently modified for use in mice by altering the *lacI* codon usage, eliminating cryptic splice sites and removing methylation-prone CpG islands¹³⁹. Alternative DNA recombination-based systems, such as β -recombinase¹⁴⁰ and ϕ C31 integrase¹⁴¹ are also being developed as genetic tools. The generation of Cre and FLP variants (or hybrids such as ‘Clp’ and ‘Fre’¹⁴²) that recognize different recognition sequences might also serve as future recombination systems¹⁴³.

All of these systems rely on appropriate promoters and/or enhancers to control the expression of effector transgenes whether they encode either DNA recombinases or transactivators. So, we are often still dependent on the most basic level of transgenesis: the ability to control the tissue-specific transcription of a transgene, ideally free of position effects. Approaches such as gene targeting, ‘promoter traps’ (for more on insertional mutagenesis, see accompanying review by Stanford *et al.* on p756 of this issue), BAC engineering (for more on this strategy, see accompanying review by Copeland *et al.* on p769 of this issue) and traditional transgenics, each with its particular advantages and disadvantages, expand our ability to achieve tissue specificity with simple transgenesis and therefore our control over these binary systems.

Finally, the molecular domains of the two broad categories of conditional transgenesis described here are DNA (site-specific recombination) and RNA (transcriptional transactivation). What about altering activity at the protein level? Technology that allows enzyme stability and activity to be regulated might soon be used in mouse transgenics as well. For example, chemical inducers of dimerization have been used to regulate protein activity¹⁴⁴, and small molecules that regulate specific enzyme activities or biological processes have been identified in chemical genetic screens¹⁴⁵, which can be considered to be a type of ‘conditional epigenetics’ because only protein and not DNA is affected. So, we are approaching an era in which we will not only be able to conditionally control gene expression in mice where and when we choose, but also at the desired level of gene expression. We might be asymptotically approaching this era, because there are always some limitations and unforeseen technical problems to be solved, but with every step the limiting reagent becomes not our tools but, increasingly, our imaginations.

1. Yarranton, G. T. Inducible vectors for expression in mammalian cells. *Curr. Opin. Biotechnol.* **3**, 506–511 (1992).
2. Khillan, J. S. *et al.* Gene transactivation mediated by the TAT gene of human immunodeficiency virus in transgenic mice. *Nucleic Acids Res.* **16**, 1423–1430 (1988).
3. Bieberich, C. J., King, C. M., Tinkle, B. T. & Jay, G. A transgenic model of transactivation by the Tax protein of HTLV-I. *Virology* **196**, 309–318 (1993).
4. Nerenberg, M., Hinrichs, S. H., Reynolds, R. K., Khoury, G. & Jay, G. The *tat* gene of human T-lymphotropic virus type 1 induces mesenchymal tumors in transgenic mice. *Science* **237**, 1324–1329 (1987).
5. Byrne, G. W. & Ruddle, F. H. Multiplex gene regulation: a two-tiered approach to transgene regulation in transgenic mice. *Proc. Natl Acad. Sci. USA* **86**, 5473–5477 (1989).
6. Gardner, D. P., Byrne, G. W., Ruddle, F. H. & Kappen, C. Spatial and temporal regulation of a *lacZ* reporter transgene in a binary transgenic mouse system. *Transgenic Res.* **5**, 37–48 (1996).
7. Rundle, C. H., Macias, M. P., Yueh, Y. G., Gardner, D. P. & Kappen, C. Transactivation of Hox gene expression in a VP16-dependent binary transgenic mouse system. *Biochim. Biophys. Acta* **1398**, 164–178 (1998).
8. Yueh, Y. G., Yaworsky, P. J. & Kappen, C. Herpes simplex virus transcriptional activator VP16 is detrimental to preimplantation development in mice. *Mol. Reprod. Dev.* **55**, 37–46 (2000).
9. Shockett, P., Difilippantonio, M., Hellman, N. & Schatz, D. G. A modified tetracycline-regulated system provides autoregulatory, inducible gene expression in cultured cells and transgenic mice. *Proc. Natl Acad. Sci. USA* **92**, 6522–6526 (1995).
10. Baron, U., Gossen, M. & Bujard, H. Tetracycline-controlled transcription in eukaryotes: novel transactivators with graded transactivation potential. *Nucleic Acids Res.* **25**, 2723–2729 (1997).
11. Baron, U. & Bujard, H. Tet repressor-based system for regulated gene expression in eukaryotic cells: principles and advances. *Methods Enzymol.* **327**, 401–421 (2000).
A useful primer of the TetR-based systems for investigators who are considering this approach.
12. Gossen, M. & Bujard, H. Tight control of gene expression in mammalian cells by tetracycline-responsive promoters. *Proc. Natl Acad. Sci. USA* **89**, 5547–5551 (1992).
13. Gossen, M. *et al.* Transcriptional activation by tetracyclines in mammalian cells. *Science* **268**, 1766–1769 (1995).
14. Kistner, A. *et al.* Doxycycline-mediated quantitative and tissue-specific control of gene expression in transgenic mice. *Proc. Natl Acad. Sci. USA* **93**, 10933–10938 (1996).
15. Hasan, M. T., Schornig, K., Berger, S., Graewe, W. & Bujard, H. Long-term, noninvasive imaging of regulated gene expression in living mice. *Genesis* **29**, 116–122 (2001).
This study provides a model for the non-invasive imaging of conditional gene expression in mice, in which the TetR system was used to simultaneously regulate two transgenes, which encode Cre and luciferase, from a bidirectional promoter. A photon-imaging system was then used to follow the expression of luciferase in living mice.
16. Lee, P. *et al.* Conditional lineage ablation to model human diseases. *Proc. Natl Acad. Sci. USA* **95**, 11371–11376 (1998).
17. Shin, M. K., Levorse, J. M., Ingram, R. S. & Tilghman, S. M. The temporal requirement for endothelin receptor-B signalling during neural crest development. *Nature* **402**, 496–501 (1999).
This paper reports the use of gene targeting to produce three mouse lines in which wild-type *EdnrB* alleles were replaced with transgenes that encode either tTA or rtTA, or a *tetO-EdnrB* target gene. These lines were crossed to produce mice in which all *EdnrB* activity was doxycycline regulatable.
18. Malleret, G. *et al.* Inducible and reversible enhancement of learning, memory, and long-term potentiation by genetic inhibition of calcineurin. *Cell* **104**, 675–686 (2001).
The value of being able to induce and then repress transgene transactivation is shown in this paper, as well as in references 19–24.
19. Mansuy, I. M., Mayford, M., Jacob, B., Kandel, E. R. & Bach, M. E. Restricted and regulated overexpression reveals calcineurin as a key component in the transition from short-term to long-term memory. *Cell* **92**, 39–49 (1998).
20. Ewald, D. *et al.* Time-sensitive reversal of hyperplasia in transgenic mice expressing SV40 T antigen. *Science* **273**, 1384–1386 (1996).
21. Chin, L. *et al.* Essential role for oncogenic Ras in tumour maintenance. *Nature* **400**, 468–472 (1999).
22. Felsner, D. W. & Bishop, J. M. Reversible tumorigenesis by MYC in hematopoietic lineages. *Mol. Cell* **4**, 199–207 (1999).
23. Huettner, C. S., Zhang, P., Van Etten, R. A. & Tenen, D. G. Reversibility of acute B-cell leukaemia induced by BCR-ABL1. *Nature Genet.* **24**, 57–60 (2000).
24. Yamamoto, A., Lucas, J. J. & Hen, R. Reversal of neuropathology and motor dysfunction in a conditional model of Huntington's disease. *Cell* **101**, 57–66 (2000).
25. Urlinger, S. *et al.* Exploring the sequence space for tetracycline-dependent transcriptional activators: novel mutations yield expanded range and sensitivity. *Proc. Natl Acad. Sci. USA* **97**, 7963–7968 (2000).
Investigators considering a TetR-based approach should review the improved rTA variants described here, as well as in reference 26.
26. Wells, K. D., Foster, J. A., Moore, K., Pursel, V. G. & Wall, R. J. Codon optimization, genetic insulation, and an rTA reporter improve performance of the tetracycline switch. *Transgenic Res.* **8**, 371–381 (1999).
27. Baron, U., Freundlieb, S., Gossen, M. & Bujard, H. Co-regulation of two gene activities by tetracycline via a bidirectional promoter. *Nucleic Acids Res.* **23**, 3605–3606 (1995).
28. Lavon, I. *et al.* High susceptibility to bacterial infection, but no liver dysfunction, in mice compromised for hepatocyte NF- κ B activation. *Nature Med.* **6**, 573–577 (2000).
29. Baron, U. *et al.* Generation of conditional mutants in higher eukaryotes by switching between the expression of two genes. *Proc. Natl Acad. Sci. USA* **96**, 1013–1018 (1999).
30. Ornitz, D. M., Moreadith, R. W. & Leder, P. Binary system for regulating transgene expression in mice: targeting *int-2* gene expression with yeast GAL4/UAS control elements. *Proc. Natl Acad. Sci. USA* **88**, 698–702 (1991).
31. Echelard, Y. *et al.* Sonic hedgehog, a member of a family of putative signaling molecules, is implicated in the regulation of CNS polarity. *Cell* **75**, 1417–1430 (1993).
32. Rowitch, D. H. *et al.* Sonic hedgehog regulates proliferation and inhibits differentiation of CNS precursor cells. *J. Neurosci.* **19**, 8954–8965 (1999).
33. Wang, Y., DeMayo, F. J., Tsai, S. Y. & O'Malley, B. W. Ligand-inducible and liver-specific target gene expression in transgenic mice. *Nature Biotechnol.* **15**, 239–243 (1997).
34. Saam, J. R. & Gordon, J. I. Inducible gene knockouts in the small intestinal and colonic epithelium. *J. Biol. Chem.* **274**, 38071–38082 (1999).
35. Wang, X. J., Liefer, K. M., Tsai, S. O'Malley, B. W. & Roop, D. R. Development of gene-switch transgenic mice that inducibly express transforming growth factor β 1 in the epidermis. *Proc. Natl Acad. Sci. USA* **96**, 8483–8488 (1999).
In this paper, topical application of a synthetic steroid was used to control *Tgfb1* expression in mouse skin through the transactivator GLVPC, showing that Tgf- β 1 inhibits epidermal cell growth.
36. Wang, X. Role of TGF β signaling in skin carcinogenesis. *Microsc. Res. Tech.* **52**, 420–429 (2001).
37. Pierson, T. M. *et al.* Regulable expression of inhibin A in wild-type and inhibin α null mice. *Mol. Endocrinol.* **14**, 1075–1085 (2000).
38. Sauer, B. & Henderson, N. Cre-stimulated recombination at *loxP*-containing DNA sequences placed into the mammalian genome. *Nucleic Acids Res.* **17**, 147–161 (1989).
39. O'Gorman, S., Fox, D. T. & Wahl, G. M. Recombinase-mediated gene activation and site-specific integration in mammalian cells. *Science* **251**, 1351–1355 (1991).
40. Buchholz, F., Angrand, P. O. & Stewart, A. F. Improved properties of FLP recombinase evolved by cycling mutagenesis. *Nature Biotechnol.* **16**, 657–662 (1998).
DNA-shuffling techniques are used here to modify the FLP recombinase so that it can function optimally at the body temperature of mice.
41. Gu, H., Zou, Y. R. & Rajewsky, K. Independent control of immunoglobulin switch recombination at individual switch regions evidenced through Cre-*loxP*-mediated gene targeting. *Cell* **73**, 1155–1164 (1993).
42. Lewandoski, M., Meyers, E. N. & Martin, G. R. Analysis of Fgf8 gene function in vertebrate development. *Cold Spring Harb. Symp. Quant. Biol.* **62**, 159–168 (1997).
43. Le, Y., Gagneten, S., Tombaccini, D., Bethke, B. & Sauer, B. Nuclear targeting determinants of the phage P1 *cre* DNA recombinase. *Nucleic Acids Res.* **27**, 4703–4709 (1999).
44. Thyagarajan, B., Guimaraes, M. J., Groth, A. C. & Calos, M. P. Mammalian genomes contain active recombinase recognition sites. *Gene* **244**, 47–54 (2000).
45. Senecoff, J. F. & Cox, M. M. Directionality in FLP protein-promoted site-specific recombination is mediated by DNA-DNA pairing. *J. Biol. Chem.* **261**, 7380–7386 (1986).
46. Loonstra, A. *et al.* Growth inhibition and DNA damage induced by Cre recombinase in mammalian cells. *Proc. Natl Acad. Sci. USA* **98**, 9209–9214 (2001).
47. Schmidt, E. E., Taylor, D. S., Prigge, J. R., Barnett, S. & Capecchi, M. R. Illegitimate Cre-dependent chromosome rearrangements in transgenic mouse spermatids. *Proc. Natl Acad. Sci. USA* **97**, 13702–13707 (2000).
48. O'Gorman, S., Dagenais, N. A., Qian, M. & Marchuk, Y. Protamine-Cre recombinase transgenes efficiently recombine target sequences in the male germ line of mice, but not in embryonic stem cells. *Proc. Natl Acad. Sci. USA* **94**, 14602–14607 (1997).
49. Lewandoski, M., Wassarman, K. M. & Martin, G. R. *Zp3-cre*, a transgenic mouse line for the activation or inactivation of *loxP*-flanked target genes specifically in the female germ line. *Curr. Biol.* **7**, 148–151 (1997).
50. Abuin, A. & Bradley, A. Recycling selectable markers in mouse embryonic stem cells. *Mol. Cell. Biol.* **16**, 1851–1856 (1996).
51. Dragatsis, I., Levine, M. S. & Zeitlin, S. Inactivation of *Hdh* in the brain and testis results in progressive neurodegeneration and sterility in mice. *Nature Genet.* **26**, 300–306 (2000).
52. Sakai, T. *et al.* Plasma fibronectin supports neuronal survival and reduces brain injury following transient focal cerebral ischemia but is not essential for skin-wound healing and hemostasis. *Nature Med.* **7**, 324–330 (2001).
53. Guy, J., Hendrich, B., Holmes, M., Martin, J. E. & Bird, A. A mouse *Mecp2*-null mutation causes neurological symptoms that mimic Rett syndrome. *Nature Genet.* **27**, 322–326 (2001).
54. Brakebusch, C. *et al.* Skin and hair follicle integrity is crucially dependent on β 1 integrin expression on keratinocytes. *EMBO J.* **19**, 3990–4003 (2000).
55. Postic, C. *et al.* Dual roles for glucokinase in glucose homeostasis as determined by liver and pancreatic β -cell-specific gene knock-outs using Cre recombinase. *J. Biol. Chem.* **274**, 305–315 (1999).
56. Gu, H., Marth, J. D., Orban, P. C., Mossmann, H. & Rajewsky, K. Deletion of a DNA polymerase β -gene segment in T cells using cell type-specific gene targeting. *Science* **265**, 103–106 (1994).
This paper describes the first tissue-specific gene inactivation and provides a model for using three *loxP* sites to remove the selection cassette from the conditional allele *in vitro*. See references 59 and 60 for variations on this strategy.
57. Meyers, E. N., Lewandoski, M. & Martin, G. R. An Fgf8 mutant allelic series generated by Cre- and Flp-mediated recombination. *Nature Genet.* **18**, 136–141 (1998).
An example of the allelogetic strategy as applied to the *Fgf8* locus, in which three alleles (hypomorphic, conditional and null) were generated from one mouse line.
58. Nagy, A. *et al.* Dissecting the role of N-myc in development using a single targeting vector to generate a series of alleles. *Curr. Biol.* **8**, 661–664 (1998).
A demonstration of how to produce an allelic series, which includes alleles with any desirable mutation, using the *N-Myc* locus as an example.
59. Xu, X. *et al.* Direct removal in the mouse of a floxed *neo* gene from a three-*loxP* conditional knockout allele by two novel approaches. *Genesis* **30**, 1–6 (2001).
60. Holzenberger, M. *et al.* Cre-mediated germline mosaicism: a method allowing rapid generation of several alleles of a target gene. *Nucleic Acids Res.* **28**, E92 (2000).
61. Partanen, J., Schwartz, L. & Rossant, J. Opposite phenotypes of hypomorphic and Y766 phosphorylation site mutations reveal a function for Fgr1 in anteroposterior patterning of mouse embryos. *Genes Dev.* **12**, 2332–2344 (1998).
62. Gage, P. J., Suh, H. & Camper, S. A. Dosage requirement of Pitx2 for development of multiple organs. *Development* **126**, 4643–4651 (1999).
63. Sun, X., Meyers, E. N., Lewandoski, M. & Martin, G. R. Targeted disruption of Fgf8 causes failure of cell migration in the gastrulating mouse embryo. *Genes Dev.* **13**, 1834–1846 (1999).
64. Meyers, E. N. & Martin, G. R. Differences in left-right axis pathways in mouse and chick: functions of FGF8 and SHH. *Science* **285**, 403–406 (1999).
65. Trumpp, A., Depew, M. J., Rubenstein, J. L., Bishop, J. M. & Martin, G. R. Cre-mediated gene inactivation demonstrates that FGF8 is required for cell survival and patterning of the first branchial arch. *Genes Dev.* **13**, 3136–3148 (1999).
66. Lewandoski, M., Sun, X. & Martin, G. R. Fgf8 signalling from the AER is essential for normal limb development. *Nature Genet.* **26**, 460–463 (2000).

67. Moon, A. M. & Capocchi, M. R. Fgf8 is required for outgrowth and patterning of the limbs. *Nature Genet.* **26**, 455–459 (2000).
68. Soriano, P. Generalized *lacZ* expression with the ROSA26 Cre reporter strain. *Nature Genet.* **21**, 70–71 (1999).
This reference and reference 72 provide two different models of how to achieve the otherwise difficult goal of near-ubiquitous transgene expression in the mouse. Such expression is ideal for producing mice that are useful as breeding partners for those that carry different tissue-specific cre or FLP transgenes.
69. Mao, X., Fujiwara, Y. & Orkin, S. H. Improved reporter strain for monitoring Cre recombinase-mediated DNA excisions in mice. *Proc. Natl Acad. Sci. USA* **96**, 5037–5042 (1999).
70. Mao, X., Fujiwara, Y., Chapdelaine, A., Yang, H. & Orkin, S. H. Activation of *EGFP* expression by Cre-mediated excision in a new ROSA26 reporter mouse strain. *Blood* **97**, 324–326 (2001).
71. Srinivas, S. *et al.* Cre reporter strains produced by targeted insertion of *EYFP* and *ECFP* into the ROSA26 locus. *BMC Dev. Biol.* **1**, 4 (2001).
72. Lobe, C. G. *et al.* Z/AP, a double reporter for cre-mediated recombination. *Dev. Biol.* **208**, 281–292 (1999).
73. Novak, A., Guo, C., Yang, W., Nagy, A. & Lobe, C. G. Z/EG, a double reporter mouse line that expresses enhanced green fluorescent protein upon Cre-mediated excision. *Genesis* **28**, 147–155 (2000).
74. Vooijs, M., Jonkers, J. & Berns, A. A highly efficient ligand-regulated Cre recombinase mouse line shows that LoxP recombination is position dependent. *EMBO Rep.* **2**, 292–297 (2001).
75. Zinyk, D. L., Mercer, E. H., Harris, E., Anderson, D. J. & Joyner, A. L. Fate mapping of the mouse midbrain-hindbrain constriction using a site-specific recombination system. *Curr. Biol.* **8**, 665–668 (1998).
76. Epstein, J. A. *et al.* Migration of cardiac neural crest cells in Splotch embryos. *Development* **127**, 1869–1878 (2000).
77. Chai, Y. *et al.* Fate of the mammalian cranial neural crest during tooth and mandibular morphogenesis. *Development* **127**, 1671–1679 (2000).
78. Jiang, X., Rowitch, D. H., Soriano, P., McMahon, A. P. & Sucov, H. M. Fate of the mammalian cardiac neural crest. *Development* **127**, 1607–1616 (2000).
79. Herrera, P. L. Adult insulin- and glucagon-producing cells differentiate from two independent cell lineages. *Development* **127**, 2317–2322 (2000).
80. Jacob, J. & Baltimore, D. Modelling T-cell memory by genetic marking of memory T cells *in vivo*. *Nature* **399**, 593–597 (1999).
81. O’Gorman, S. & Barcace, E. A. Multiple neuronal identities and migrations specified by a single Hox gene in mice. *Development* (in the press).
82. Lakso, M. *et al.* Targeted oncogene activation by site-specific recombination in transgenic mice. *Proc. Natl Acad. Sci. USA* **89**, 6232–6236 (1992).
This first demonstration of Cre-mediated transgene activation provides a model for the general strategy. For an elaboration of this model, see reference 88, and for an alternative model, see reference 83.
83. Grieshammer, U., Lewandoski, M., Prevet, D., Oppenheim, R. W. & Martin, G. R. Muscle-specific cell ablation conditional upon Cre-mediated DNA recombination in transgenic mice leads to massive spinal and cranial motoneuron loss. *Dev. Biol.* **197**, 234–247 (1998).
84. Lewandoski, M. & Martin, G. R. Cre-mediated chromosome loss in mice. *Nature Genet.* **17**, 223–225 (1997).
85. Garrick, D., Fiering, S., Martin, D. I. & Whitelaw, E. Repeat-induced gene silencing in mammals. *Nature Genet.* **18**, 56–59 (1998).
86. Drago, J. *et al.* Targeted expression of a toxin gene to D1 dopamine receptor neurons by cre-mediated site-specific recombination. *J. Neurosci.* **18**, 9845–9857 (1998).
87. Lee, K. J., Dietrich, P. & Jessell, T. M. Genetic ablation reveals that the roof plate is essential for dorsal interneuron specification. *Nature* **403**, 734–740 (2000).
88. Dragatsis, I. & Zeitlin, S. A method for the generation of conditional gene repair mutations in mice. *Nucleic Acids Res.* **29**, E10 (2001).
89. Arin, M. J., Longley, M. A., Wang, X. J. & Roop, D. R. Focal activation of a mutant allele defines the role of stem cells in mosaic skin disorders. *J. Cell Biol.* **152**, 645–649 (2001).
90. Cao, T., Longley, M. A., Wang, X. J. & Roop, D. R. An inducible mouse model for epidermolysis bullosa simplex: implications for gene therapy. *J. Cell Biol.* **152**, 651–656 (2001).
91. Kuhn, R., Schwenk, F., Aguet, M. & Rajewsky, K. Inducible gene targeting in mice. *Science* **269**, 1427–1429 (1995).
92. Gerber, H. P. *et al.* VEGF is required for growth and survival in neonatal mice. *Development* **126**, 1149–1159 (1999).
93. Alonzi, T. *et al.* Essential role of STAT3 in the control of the acute-phase response as revealed by inducible gene activation in the liver. *Mol. Cell. Biol.* **21**, 1621–1632 (2001).
94. Rohlmann, A., Gotthardt, M., Hammer, R. E. & Herz, J. Inducible inactivation of hepatic *LRP* gene by cre-mediated recombination confirms role of LRP in clearance of chylomicron remnants. *J. Clin. Invest.* **101**, 689–695 (1998).
95. Sjogren, K. *et al.* Liver-derived insulin-like growth factor I (IGF-I) is the principal source of IGF-I in blood but is not required for postnatal body growth in mice. *Proc. Natl Acad. Sci. USA* **96**, 7088–7092 (1999).
96. Lam, K. P. & Rajewsky, K. Rapid elimination of mature autoreactive B cells demonstrated by Cre-induced change in B cell antigen receptor specificity *in vivo*. *Proc. Natl Acad. Sci. USA* **95**, 13171–13175 (1998).
97. Lam, K. P., Kuhn, R. & Rajewsky, K. *In vivo* ablation of surface immunoglobulin on mature B cells by inducible gene targeting results in rapid cell death. *Cell* **90**, 1073–1083 (1997).
98. Radtke, F. *et al.* Deficient T cell fate specification in mice with an induced inactivation of *Notch1*. *Immunity* **10**, 547–558 (1999).
99. Kellendonk, C. *et al.* Inducible site-specific recombination in the brain. *J. Mol. Biol.* **285**, 175–182 (1999).
100. Kellendonk, C. *et al.* Regulation of Cre recombinase activity by the synthetic steroid RU 486. *Nucleic Acids Res.* **24**, 1404–1411 (1996).
101. Feil, R. *et al.* Ligand-activated site-specific recombination in mice. *Proc. Natl Acad. Sci. USA* **93**, 10887–10890 (1996).
102. Danielian, P. S., Muccino, D., Rowitch, D. H., Michael, S. K. & McMahon, A. P. Modification of gene activity in mouse embryos *in utero* by a tamoxifen-inducible form of Cre recombinase. *Curr. Biol.* **8**, 1323–1326 (1998).
103. Brocard, J. *et al.* Spatio-temporally controlled site-specific somatic mutagenesis in the mouse. *Proc. Natl Acad. Sci. USA* **94**, 14559–14563 (1997).
104. Schwenk, F., Kuhn, R., Angrand, P. O., Rajewsky, K. & Stewart, A. F. Temporally and spatially regulated somatic mutagenesis in mice. *Nucleic Acids Res.* **26**, 1427–1432 (1998).
105. Metzger, D., Clifford, J., Chiba, H. & Chambon, P. Conditional site-specific recombination in mammalian cells using a ligand-dependent chimeric Cre recombinase. *Proc. Natl Acad. Sci. USA* **92**, 6991–6995 (1995).
106. Zhang, Y. *et al.* Inducible site-directed recombination in mouse embryonic stem cells. *Nucleic Acids Res.* **24**, 543–548 (1996).
107. Metzger, D. & Chambon, P. Site- and time-specific gene targeting in the mouse. *Methods* **24**, 71–80 (2001).
108. Vasioukhin, V., Degenstein, L., Wise, B. & Fuchs, E. The magical touch: genome targeting in epidermal stem cells induced by tamoxifen application to mouse skin. *Proc. Natl Acad. Sci. USA* **96**, 8551–8556 (1999).
109. Zhang, Y., Wienands, J., Zurn, C. & Reth, M. Induction of the antigen receptor expression on B lymphocytes results in rapid competence for signaling of SLP-65 and Syk. *EMBO J.* **17**, 7304–7310 (1998).
110. Imai, T., Jiang, M., Kastner, P., Chambon, P. & Metzger, D. Selective ablation of retinoid X receptor α in hepatocytes impairs their lifespan and regenerative capacity. *Proc. Natl Acad. Sci. USA* **98**, 4581–4586 (2001).
111. Kimmel, R. A. *et al.* Two lineage boundaries coordinate vertebrate apical ectodermal ridge formation. *Genes Dev.* **14**, 1377–1389 (2000).
112. Wunderlich, F. T., Wildner, H., Rajewsky, K. & Edenhofer, F. New variants of inducible Cre recombinase: a novel mutant of Cre-PR fusion protein exhibits enhanced sensitivity and an expanded range of inducibility. *Nucleic Acids Res.* **29**, E47 (2001).
Investigators considering inducible cre experiments should review these improved Cre-PR variants, as well as the Cre-ER (T2) variant in references 113–115.
113. Indra, A. K. *et al.* Temporally-controlled site-specific mutagenesis in the basal layer of the epidermis: comparison of the recombinase activity of the tamoxifen-inducible Cre-ER(T) and Cre-ER(T2) recombinases. *Nucleic Acids Res.* **27**, 4324–4327 (1999).
114. Imai, T., Jiang, M., Chambon, P. & Metzger, D. Impaired adipogenesis and lipolysis in the mouse upon selective ablation of the retinoid X receptor α mediated by a tamoxifen-inducible chimeric Cre recombinase (Cre-ERT2) in adipocytes. *Proc. Natl Acad. Sci. USA* **98**, 224–228 (2001).
115. Li, M. *et al.* Skin abnormalities generated by temporally controlled RXR α mutations in mouse epidermis. *Nature* **407**, 633–636 (2000).
116. Utomo, A. R., Nikitin, A. Y. & Lee, W. H. Temporal, spatial, and cell type-specific control of Cre-mediated DNA recombination in transgenic mice. *Nature Biotechnol.* **17**, 1091–1096 (1999).
117. Holzenberger, M., Zaoui, R., Leneuve, P., Hamard, G. & Le Bouc, Y. Ubiquitous postnatal LoxP recombination using a doxycycline auto-inducible Cre transgene (DAI-Cre). *Genesis* **26**, 157–159 (2000).
118. St-Onge, L., Furth, P. A. & Gruss, P. Temporal control of the Cre recombinase in transgenic mice by a tetracycline responsive promoter. *Nucleic Acids Res.* **24**, 3875–3877 (1996).
119. Rohlmann, A., Gotthardt, M., Willnow, T. E., Hammer, R. E. & Herz, J. Sustained somatic gene inactivation by viral transfer of Cre recombinase. *Nature Biotechnol.* **14**, 1562–1565 (1996).
120. Wang, Y., Krushel, L. A. & Edelman, G. M. Targeted DNA recombination *in vivo* using an adenovirus carrying the cre recombinase gene. *Proc. Natl Acad. Sci. USA* **93**, 3932–3936 (1996).
121. Agah, R. *et al.* Gene recombination in postmitotic cells. Targeted expression of Cre recombinase provokes cardiac-restricted, site-specific rearrangement in adult ventricular muscle *in vivo*. *J. Clin. Invest.* **100**, 169–179 (1997).
122. Brooks, A. I., Mukherjee, B., Panahian, N., Cory-Slechta, D. & Federoff, H. J. Nerve growth factor somatic mosaicism produced by herpes virus-directed expression of cre recombinase. *Nature Biotechnol.* **15**, 57–62 (1997).
123. Christensen, G., Minamisawa, S., Gruber, P. J., Wang, Y. & Chien, K. R. High-efficiency, long-term cardiac expression of foreign genes in living mouse embryos and neonates. *Circulation* **101**, 178–184 (2000).
124. Brooks, A. I., Cory-Slechta, D. A. & Federoff, H. J. Gene–experience interaction alters the cholinergic septohippocampal pathway of mice. *Proc. Natl Acad. Sci. USA* **97**, 13378–13383 (2000).
125. Shibata, H. *et al.* Rapid colorectal adenoma formation initiated by conditional targeting of the *Apc* gene. *Science* **278**, 120–123 (1997).
126. Lee, Y. H., Sauer, B., Johnson, P. F. & Gonzalez, F. J. Disruption of the *c/ebp* α gene in adult mouse liver. *Mol. Cell. Biol.* **17**, 6014–6022 (1997).
127. Piston, D. W., Knobel, S. M., Postic, C., Shelton, K. D. & Magnuson, M. A. Adenovirus-mediated knockout of a conditional glucokinase gene in isolated pancreatic islets reveals an essential role for proximal metabolic coupling events in glucose-stimulated insulin secretion. *J. Biol. Chem.* **274**, 1000–1004 (1999).
128. Ludwig, D. L., Stringer, J. R., Wight, D. C., Doetschman, H. C. & Duffy, J. J. FLP-mediated site-specific recombination in microinjected murine zygotes. *Transgenic Res.* **5**, 385–395 (1996).
129. Dymecki, S. M. Fip recombinase promotes site-specific DNA recombination in embryonic stem cells and transgenic mice. *Proc. Natl Acad. Sci. USA* **93**, 6191–6196 (1996).
130. Buchholz, F., Ringrose, L., Angrand, P. O., Rossi, F. & Stewart, A. F. Different thermostabilities of FLP and Cre recombinases: implications for applied site-specific recombination. *Nucleic Acids Res.* **24**, 4256–4262 (1996).
131. Rodriguez, C. I. *et al.* High-efficiency deleter mice show that FLP is an alternative to Cre-loxP. *Nature Genet.* **25**, 139–140 (2000).
132. Vooijs, M., van der Valk, M., te Riele, H. & Berns, A. Fip-mediated tissue-specific inactivation of the retinoblastoma tumor suppressor gene in the mouse. *Oncogene* **17**, 1–12 (1998).
133. Logie, C. & Stewart, A. F. Ligand-regulated site-specific recombination. *Proc. Natl Acad. Sci. USA* **92**, 5940–5944 (1995).
134. Logie, C., Nichols, M., Myles, K., Funder, J. W. & Stewart, A. F. Positive and negative discrimination of estrogen receptor agonists and antagonists using site-specific DNA recombinase fusion proteins. *Mol. Endocrinol.* **12**, 1120–1132 (1998).
135. Farley, F. W., Soriano, P., Steffen, L. S. & Dymecki, S. M. Widespread recombinase expression using FLPeR (flipper) mice. *Genesis* **28**, 106–110 (2000).
136. Rodriguez, C. I. & Dymecki, S. M. Origin of the precerebellar system. *Neuron* **27**, 475–486 (2000).
Fip expression is used here to permanently mark Wnt1-expressing cells and to trace their descendants. This paper contains an informative discussion of the advantages and potential

- artefacts of this type of cell-lineage analysis.**
137. Marquardt, T. *et al.* Pax6 is required for the multipotent state of retinal progenitor cells. *Cell* **105**, 43–55 (2001).
138. Saez, E. *et al.* Identification of ligands and coligands for the ecdysone-regulated gene switch. *Proc. Natl Acad. Sci. USA* **97**, 14512–14517 (2000).
139. Cronin, C. A., Gluba, W. & Scrable, H. The *lac* operator–repressor system is functional in the mouse. *Genes Dev.* **15**, 1506–1517 (2001).
- This paper reports a modified *lac* operator–repressor system that functions in the mouse and was used to regulate a coat-colour transgene, such that its expression is IPTG responsive.**
140. Diaz, V. *et al.* New insights into host factor requirements for prokaryotic β -recombinase-mediated reactions in mammalian cells. *J. Biol. Chem.* **276**, 16257–16264 (2001).
141. Thyagarajan, B., Olivares, E. C., Hollis, R. P., Ginsburg, D. S. & Calos, M. P. Site-specific genomic integration in mammalian cells mediated by phage ϕ C31 integrase. *Mol. Cell. Biol.* **21**, 3926–3934 (2001).
142. Shaikh, A. C. & Sadowski, P. D. Chimeras of the Flp and Cre recombinases: tests of the mode of cleavage by Flp and Cre. *J. Mol. Biol.* **302**, 27–48 (2000).
143. Hartung, M. & Kisters-Woike, B. Cre mutants with altered DNA binding properties. *J. Biol. Chem.* **273**, 22884–22891 (1998).
144. Spencer, D. M. Creating conditional mutations in mammals. *Trends Genet.* **12**, 181–187 (1996).
145. Peterson, R. T., Link, B. A., Dowling, J. E. & Schreiber, S. L. Small molecule developmental screens reveal the logic and timing of vertebrate development. *Proc. Natl Acad. Sci. USA* **97**, 12965–12969 (2000).
146. Wang, Y., Xu, J., Pierson, T., O'Malley, B. W. & Tsai, S. Y. Positive and negative regulation of gene expression in eukaryotic cells with an inducible transcriptional regulator. *Gene Ther.* **4**, 432–441 (1997).
147. She, P. *et al.* Phosphoenolpyruvate carboxykinase is necessary for the integration of hepatic energy metabolism. *Mol. Cell. Biol.* **20**, 6508–6517 (2000).
148. Wang, Y. *et al.* A mouse model for achondroplasia produced by targeting fibroblast growth factor receptor 3. *Proc. Natl Acad. Sci. USA* **96**, 4455–4460 (1999).
149. Hayhurst, G. P., Lee, Y. H., Lambert, G., Ward, J. M. & Gonzalez, F. J. Hepatocyte nuclear factor 4 α (nuclear receptor 2A1) is essential for maintenance of hepatic gene expression and lipid homeostasis. *Mol. Cell. Biol.* **21**, 1393–1403 (2001).
150. Abel, E. D. *et al.* Cardiac hypertrophy with preserved contractile function after selective deletion of GLUT4 from the heart. *J. Clin. Invest.* **104**, 1703–1714 (1999).
151. Lowe, L. A., Yamada, S. & Kuehn, M. R. Genetic dissection of nodal function in patterning the mouse embryo. *Development* **128**, 1831–1843 (2001).
152. Matsuda, M. *et al.* SREBP cleavage-activating protein (SCAP) is required for increased lipid synthesis in liver induced by cholesterol deprivation and insulin elevation. *Genes Dev.* **15**, 1206–1216 (2001).
153. Rucker, E. B. *et al.* Bcl-x and Bax regulate mouse primordial germ cell survival and apoptosis during embryogenesis. *Mol. Endocrinol.* **14**, 1038–1052 (2000).

Acknowledgements

I would like to apologize to colleagues whose work is not men-

tioned due to lack of space. Many thanks to H. Bujard, S. Dymecki, C. Lobe, M. Magnuson, S. O'Gorman, D. Roop and D. Rowitch for helpful discussion and correspondence, and for sharing unpublished data. I would also like to thank G. Martin, S. Sharam, C. Stewart, L. Tessarollo, T. Yamaguchi and members of my lab for comments on the manuscript, and G. Martin for years of discussions, support and encouragement.

Online links

DATABASES

The following terms in this article are linked online to:

MGI: <http://www.informatics.jax.org/Ednrb> | [Fgfr1](http://www.informatics.jax.org/Fgfr1) | [Hoxb1](http://www.informatics.jax.org/Hoxb1) | [huntingtin](http://www.informatics.jax.org/huntingtin) | [Mx1](http://www.informatics.jax.org/Mx1) | [Sonic](http://www.informatics.jax.org/Sonic) | [hedgehog](http://www.informatics.jax.org/hedgehog) | [Tgfb1](http://www.informatics.jax.org/Tgfb1) | [Wnt1](http://www.informatics.jax.org/Wnt1)

OMIM: <http://www.ncbi.nlm.nih.gov/Omim/> | [Huntington disease](http://www.ncbi.nlm.nih.gov/Omim/Huntington%20disease)

Saccharomyces genome database: <http://genome-www.stanford.edu/Saccharomyces/Gal4>

FURTHER INFORMATION

ClonTech: <http://www.clontech.com/tet/index.shtml>

Cre transgenic database: <http://www.mshri.on.ca/nagy/default.htm>

GeneSwitch™: <http://www.geneswitch.com/default.htm>

Hermann Bujard's trouble-shooting guide: <http://www.zmbh.uni-heidelberg.de/bujard/Homepage.html>

Knoll GmbH: <http://www.basf-pharma.com/tet/index.html>

Mark Lewandoski's lab: <http://ccr.ncifcrf.gov/staff/staff.asp?name=lewandoski>