




9. MECHANISMS OF DEVELOPMENT I – REGULATION OF LIMB DEVELOPMENT BY FIBROBLAST GROWTH FACTORS (FGF)

4 receptors: FGFR1-4
22 ligands: FGF1-23






Initiation: *Tbx5* - dependent





A Induction

B Progression



Outgrowth: *Tbx5* - independent



wt

Fgf10^{-/-}

wt



Fgf10^{-/-}

c

d

e



f



AER

k







l



wt

Hindlimb

Fgf8/Fgf4^{-/-}





Double KO
No *Fgf8* expression
No *Fgf4* expression

Transient *Fgf8* expression
Precocious, transient *Fgf4* expression


wt

Hindlimb

Fgf8/Fgf4^{-/-}




a Normal FL and HL




AER KO mutant phenotypes

b *Fgf8* single KO FL



c *Fgf4; Fgf8* double KO FL




d *Fgf8* single KO HL



e *Fgf4; Fgf8* double KO HL



How do the limbs grow?



FGFR3



Tyrosine Kinase Group

Tyrosine Kinase Like Group



Group of Homologs of Yeast Sterile Protein Kinases



Casein Protein Kinases

PKA, PKG & PKC Containing Group

Calcium/Calmodulin Dependent Protein Kinase Group

CDK, MAP, GSK3 & CLK Containing Group







FGFR3-related skeletal dysplasia

Hypochondroplasia
Achondroplasia
SADDAN
Thanatophoric Dysplasia

STATURE



FGFR3-related skeletal dysplasia





Achondroplasia

Thanatophoric Dysplasia





healthy

- short long bones
- brachydactyly
- macrocephaly





TD



- low nasal bridge
- spinal stenosis
- temporal lobe malformations




Loss-of-function

vs. Gain-of-function

A**B**





healthy





TD


FGFR3 inhibits chondrocyte proliferation by arresting their cell cycle in G1 phase




....via inhibition of cdk activity necessary for progression through the G1 phase of a cell cycle



FGF alters the cartilage-like phenotype of chondrocytes





.....via MMP-mediated degradation of extracellular matrix






FGF2 activates Erk and p38 MAPK, PLC γ and PKB in chondrocytes

FGF2


C1 C2 1' 5' 10' 30' 1h 2h 4h 8h





.....but only Ras/Erk activity is involved in FGF-induced growth arrest



Erk MAP kinase activity is necessary for FGFR3 phenotype in cartilage




C-type Natriuretic Peptide (CNP) over-expression results in skeleton overgrowth in mice




wild-type

CNP ↑



CNP over-expression???

CNP rescues dwarfism caused by ACH mutation in FGFR3




wild-type


$Fgfr3^{Ach}$

$Fgfr3^{Ach/CNP}$ ↑






CNP counteracts FGF2-mediated chondrocyte growth arrest through cGMP-dependent pathway




CNP antagonizes FGF2-mediated loss of cartilage extracellular matrix in chondrocytes





CNP counteracts FGF2-mediated activation of Erk MAP kinase in chondrocytes




CNP inhibits Erk MAP kinase module at the Raf level




Is protein kinase C (PKC) involved in FGFR3-mediated activation of Erk in chondrocytes?







| | | | | | | | | | | | |
|------------------|---|---|---|---|----|----|----|---|----|----|---|
| FGF2 | - | + | + | + | + | + | + | + | + | + | + |
| Bis I (μ M) | - | - | 1 | 5 | 10 | 20 | 50 | - | - | - | - |
| SU5402(μ M) | - | - | - | - | - | - | - | 1 | 10 | 20 | - |
| vehicle(DMSO) | - | - | + | + | + | + | + | + | + | + | + |


kinase






| | | | | | | | | | |
|-------|---|----|-----|-----|----|---|-----|-----|---|
| FGF2 | - | 5' | 10' | 30' | 1h | - | 30' | 30' | - |
| Bis I | - | - | - | - | - | - | - | + | + |





Protein kinase C inhibitor Bisindolylmaleimide I (Bis I) suppresses the FGF2-mediated activation of Erk MAP kinase pathway in chondrocytes by preventing the SHP2 association with FRS2 and Gab1 adaptor proteins




FGFR3 associates with STAT1 and acts as STAT1-kinase in chondrocytes




Chronic FGF stimulus inhibits cytokine/STAT signaling in chondrocytes




Chronic FGF stimulus inhibits cytokine/STAT signaling in chondrocytes



FGF2 causes premature senescence in chondrocytes



FGF2 signals towards the cytoskeleton in chondrocytes




Where is Wnt?




FGF2: C1 15' 1h 3h 6h 12h 24h 48h 72h C2 C3


b-catenin



actin



2001



2008

