Global analysis. Exercises 8

1) Find $\int_M \omega$ for

- $\omega = (x y)dx + (x + y)dy$ and M is the segment AB with A = (2,3) and B = (3,5);
- $\omega = ydx + xdy$ and M is a quarter of the circle $g(t) = (R\cos t, R\sin t), t \in (0, \frac{\pi}{2});$
- $\omega = xdx + ydy + (x + y 1)dz$ and M is the segment AB with A = (1, 1, 1)and B = (2, 3, 4).

2) Find

$$\int_M dx_3 \wedge dx_4 + x_1 x_3 dx_2 \wedge dx_4,$$

where $M = \{(x_1, x_2, x_3, x_4) \in \mathbb{R}^4 | x_1^2 + x_2^2 = 1, x_3^2 + x_4^2 = 1\}$ and the orientation is given by the parameterization $g(u, v) = (\cos u, \sin u, \cos v, \sin v), (u, v) \in (0, 2\pi) \times (0, 2\pi).$

- 3) Using the Stokes Theorem, find $\int_M \omega$, where
 - $\omega = (x^2 + y^2)dx + (x^2 y^2)dy$ and M is the circuit of the triangle with the vertexes A = (0, 0), B = (1, 0), C = (0, 1);
 - $\omega = (x^2 + y^2)dx + (x^2 y^2)dy$ and M is the graph of the function y = 1 |1 x|, $x \in (0, 2)$, starting at (0, 0);
 - $\omega = xydy \wedge dx + yzdz \wedge dx + xzdx \wedge dy$ and M is the surface of the pyramid bounded by the planes x = 0, y = 0, z = 0, x + y + z = 0.