



Short communication

Clenbuterol enhances the production of kynurenic acid in brain cortical slices and glial cultures

Elżbieta Luchowska¹, Renata Kloc¹, Sebastian Wnuk¹, Bartosz Olajossy¹, Marian Wielosz¹, Ewa M. Urbańska^{1,2}

¹Department of Pharmacology and Clinical Pharmacology, Skubiszewski Medical School, Jaczewskiego 8, PL 20-090 Lublin, Poland

²Department of Toxicology, Institute of Agricultural Medicine, Jaczewskiego 2, PL 20-950 Lublin, Poland

Correspondence: Ewa M. Urbańska, e-mail: emurbanska@poczta.onet.pl

Abstract:

The effect of a β_2 -adrenergic agonist, clenbuterol on the production of a glutamate receptor antagonist, kynurenic acid was studied *in vitro*. Clenbuterol enhanced the production of kynurenic acid in brain cortical slices (0.1–1.0 mM) and in glial cultures (1–50 μ M). Timolol, a non-selective β -adrenergic antagonist prevented this effect. The presented data indicate a novel mechanism of action of β_2 -adrenoceptor agonists and suggest that an increased formation of the endogenous glutamate receptor antagonist, kynurenic acid could partially contribute to their neuroprotective activity.

Key words:

β -adrenergic agonist, kynurenic acid, glial culture, *in vitro*

Abbreviations: GFAP – glial fibrillary acidic protein, HPLC – high performance liquid chromatography, KAT – kynurenine aminotransferase, KRB – Krebs-Ringer buffer, NGF – nerve growth factor, NMDA – N-methyl-D-aspartate

Introduction

Kynurenic acid is the only known endogenous compound able to block the glycine site of N-methyl-D-aspartate (NMDA) receptors and to antagonize α_7 subtype of cholinergic nicotinic receptors [15]. It is formed in the brain and in the periphery [10, 15, 18] from a direct bioprecursor, L-kynurenine, *via* enzymatic reaction mediated by kynurenine aminotransfe-

rases I and II (KAT I and II). Kynurenic acid exerts anticonvulsant and neuroprotective activity in experimental models, and its disturbed metabolism was suggested to be one of the causative factors in human neuropathology [15, 17]. Kynurenic acid may also inhibit glutamate release, and this effect is not shared by other NMDA or α_7 cholinergic antagonists [2]. It blocks pre-synaptic NMDA autoreceptors already at low, submicromolar concentrations, thereby regulating glutamate release [12]. Kynurenic acid formation is controlled by several distinct intra- and intercellular mechanisms, such as e.g. endogenous excitatory amino acid agonists or D,L-homocysteine [3, 8, 13, 17]. Here, we demonstrate that β_2 -adrenergic agonist clenbuterol enhances the production of kynurenic acid *ex vivo*, in rat brain cortical slices and in mixed glial cultures.

Materials and Methods

Animals

Cortical slices were obtained from male Wistar rats, weighing 220–250 g. Newborn Wistar rats (postnatal day 1–2) were used for the preparation of glial cultures. Animals were housed under standard laboratory conditions, at ambient temperature of 20°C, with food and water available *ad libitum*. All experimental procedures were approved by the Local Ethics Committee in Lublin and are in agreement with European Communities Council Directive on the use of animals in experimental studies.

Substances

L-kynurenine (sulfate salt), kynurenic acid, L-pyruvate, pyridoxal-5'-phosphate, 2-mercaptoethanol, clenbuterol and timolol and cellulose membrane dialysis tubing were obtained from Sigma-Aldrich (St. Louis, USA), whereas all high pressure liquid chromatography (HPLC) reagents were obtained from J.T. Baker Laboratory Chemicals. All other chemicals were purchased from Sigma-Aldrich (St. Louis, USA).

In vitro experiments – cortical slices

Production of kynurenic acid in freshly prepared cortical slices was performed as previously described [8]. Each well contained 6 slices, and each concentration of the drug was studied in 6 wells (N = 6). Briefly, slices were incubated (2 h, 37°C) in Krebs-Ringer buffer (KRB), pH 7.4 (118.5 mM NaCl, 4.75 mM KCl, 1.18 mM MgSO₄, 1.77 mM CaCl₂, 12.9 mM NaH₂PO₄, 3 mM Na₂HPO₄ and 5 mM D-glucose) saturated with 95% O₂/5% CO₂, in the presence of L-kynurenine (10 μM) and various concentrations of the studied substances. Kynurenic acid was separated from the incubation medium and quantified fluorimetrically using a high performance liquid chromatography (HPLC) system (Varian, USA) and ESA catecholamine HR-80, 3 μm, C18 reverse-phase column. The mean control production of kynurenic acid in the presence of 10 μM L-kynurenine was 2.98 ± 0.61 pmol/h/well.

In vitro experiments – glial cultures

Mixed glial cultures were obtained from cerebral hemispheres of rat pups (postnatal day 1–2) as de-

scribed by others [5, 19], with our modification. Briefly, brain tissue of newborn rats was seeded at a density of one brain/75 cm² culture flasks (Falcon, Switzerland) containing 10 ml of Basal Medium Eagle with Earle's salts (BME, Biochrom AG, Berlin), supplemented with 2 mM L-glutamine, penicillin-streptomycin (500 IU/ml – 500 UG/ml; Gibco-BRL) and containing 10% heat-inactivated fetal bovine serum (Gibco, InVitrogen). The medium was replaced twice a week. Cultures were maintained at 37°C in a 5% CO₂/95% air atmosphere. After 14–15 days, the cells were replated on poly-L-lysine-coated 24-microwell plates (Nunc), and the medium was replaced every 3 days. After 1 week, a confluent astrocytic monolayer developed with scattered oligodendrocytes and microglia on top. Approx. 75–80% of the cells were glial fibrillary acidic protein (GFAP) – positive (as revealed by immunostaining). The production of kynurenic acid was assessed in tissue cultures after 20–22 days *in vitro*. On the day of experiment, the culture medium was changed and the studied substances were added to cultures. After 22 h, the medium was changed into the freshly prepared KRB of the above composition. Cells were further incubated in the presence of L-kynurenine (final concentration 10 μM; 2 h) and the studied substances (total incubation time 24 h). Each concentration of the drug was studied in 6 wells (N = 6). The mean control production of kynurenic acid by glial cells in the presence of 10 μM L-kynurenine was 11.8 ± 1.25 pmol/h/well.

Activities of kynurenine aminotransferases (KATs)

The activities of KAT I and KAT II were assayed in the dialyzed homogenate of cortical brain tissue, as described before [8]. Briefly, freshly obtained cortical brain tissue was homogenized (1:9; w/v) in 5 mM Tris-acetate buffer, pH 8.0, containing 50 μM pyridoxal-5'-phosphate and 10 mM 2-mercaptoethanol. The resulting homogenate was centrifuged (12,000 rpm for 10 min), and the supernatant was dialyzed overnight at 8°C, using cellulose membrane dialysis tubing, against 4 l of the buffer composed as above. The enzyme preparation was incubated in the reaction mixture containing 2 μM L-kynurenine, 1 mM pyruvate, 70 μM pyridoxal-5'-phosphate, 150 mM Tris-acetate buffer, at pH of 7.0 or 9.5, for KAT II or KAT I, respectively (all given concentrations are the final ones) and solutions of the tested drugs at varying con-

centrations. When KAT II activity was assayed, glutamine (final concentration 2 mM), the KAT I inhibitor, was added to the samples. Further procedures were performed as described above. The mean control activity of KAT I and KAT II in the presence of 2 μ M kynurenic acid was 1.72 ± 0.09 and 0.78 ± 0.07 pmol of kynurenic acid/mg of tissue/h, respectively.

Each experiment was repeated at least twice. Statistical comparisons were performed using one-way analysis of variance (ANOVA) with Bonferroni *post-hoc* test.

Results

Clenbuterol enhanced the kynurenic acid synthesis in cortical slices (0.25–1.0 mM) and in glial cultures (1–50 μ M) (Fig. 1). The non-selective β -adrenergic antagonist timolol, used at the 50 μ M concentration in cortical slices and at 5 μ M concentration in glial cultures, has reversed its action in both experiments (Fig. 1). KAT I and KAT II activities were not altered by clenbuterol at up to 0.5 mM concentration (Tab. 1).

Discussion

Central noradrenergic system may modulate neuronal survival, neuroplasticity and neurogenesis [14]. Neurons and glial cells express several subtypes of functional α - and β -adrenergic receptors, moreover, β_2 -adrenergic receptors are located predominantly on the glial cells [14]. Agonists of β_2 -adrenoceptors display neuroprotective effects *in vitro* and *in vivo* [7, 14]. Clenbuterol protected mouse cerebral cortex and rat hippocampus from ischemic damage and attenuated hippocampal neuronal loss induced by L-glutamate in culture [16]. These effects seemed to result from the induction of nerve growth factor (NGF) synthesis [4, 16]. Moreover, the compound was shown to increase brain levels of tryptophan [11]. Clinically, drugs activating β_2 -adrenergic receptors are commonly used to treat and to prevent the attacks of bronchial asthma due to their potent peripheral effects, including bronchodilation and enhancement of mucociliary clearance [1].




Fig. 1. The effect of clenbuterol on the kynurenic acid production in rat brain cortical slices and in mixed glial cultures. Cortical slices were incubated for 2 h in KRB, pH 7.4, in the presence of 10 μ M L-kynurenic acid and 50–1000 μ M clenbuterol. Timolol (50 μ M) was added to the incubation media 10 min prior to clenbuterol. Glial cells were incubated in the presence 0.1–50 μ M clenbuterol for 24 h. L-kynurenic acid (final concentration 10 μ M) was present in the incubation media during the last 2 h of incubation. Timolol (5 μ M) was added to the incubation media 10 min prior to clenbuterol. * $p < 0.05$ vs. control (ANOVA); ^a $p < 0.05$ vs. clenbuterol (ANOVA)

Tab. 1. The effect of clenbuterol on the activity of kynurenic acid aminotransferases (KATs) I and II

Clenbuterol (μ M)	KAT I (% of control)	KAT II (% of control)
10	101.8 \pm 4.0	98.5 \pm 3.7
50	97.8 \pm 5.3	102.7 \pm 4.8
100	98.1 \pm 5.5	95.2 \pm 8.5
500	104.7 \pm 4.5	95.7 \pm 6.2

Data are presented as the percentage of control values \pm SD

The data presented herein suggest a novel central mechanism of action of β_2 -adrenergic agonists involving an increased production of glutamate receptor antagonist, kynurenic acid. Timolol, a non-selective β -adrenergic antagonist reversed the action of clenbuterol *in vitro*; KATs activities were not changed by clenbuterol. Thus, the observed increase in the kynurenic acid production seems to result from a direct stimulation of the β_2 -adrenergic receptors and not from the activation of kynurenic acid biosynthetic enzymes. The regulation of kynurenic acid synthesis is governed by various extracellular and intracellular mechanisms. Notably, only few drugs, e.g. some antiepileptics, are able to stimulate kynurenic acid production, but when used at higher, close to millimolar concentrations [9]. The concentrations of clenbuterol effective in glial cultures (1–50 μ M) are comparable with the neuroprotective concentrations of clenbuterol (1 μ M) re-

ported by others [7]. The underlying cellular mechanism most probably is associated with an increased formation of intracellular cAMP, the second messenger mediating the effects of β_2 -adrenergic stimulation [6, 7]. Further experiments will clarify this issue.

An enhanced synthesis of kynurenic acid may be one of the mechanisms contributing to the neuroprotective effects of β -adrenoceptor agonists. Development of drugs modifying noradrenergic receptor activity and/or downstream signaling, especially in astrocytes, was suggested for treatment of several neurological and psychiatric disorders and as a novel therapy in neurodegeneration [7]. Our current research focuses on the assessment of the effects exerted by a broad range of β -adrenergic agonists on the brain synthesis of kynurenic acid *in vitro* and identification of cellular mechanisms underlying this action.

Acknowledgments:

This study was supported by the grant from the State Committee for Scientific Research, KBN 2P05F 04429 and grant from Medical University School in Lublin, DS. 450/08.

References:

- Anderson GP: Interactions between corticosteroids and β -adrenergic agonists in asthma disease induction, progression, and exacerbation. *Am J Respir Crit Care Med*, 2000, 161, S188–196.
- Carpenedo R, Pittaluga A, Cozzi A, Attucci S, Galli A, Raiteri M, Moroni F: Presynaptic kynurenate-sensitive receptors inhibit glutamate release. *Eur J Neurosci*, 2001, 13, 2141–2147.
- Chmiel-Perzyńska I, Perzyński A, Wielosz M, Urbańska EM: Hyperglycemia enhances the inhibitory effect of mitochondrial toxins and D,L-homocysteine on the brain production of kynurenic acid. *Pharmacol Rep*, 2007, 59, 268–273.
- Culmsee C, Stumm RK, Schafer MK, Weihe E, Kriegstein J: Clenbuterol induces growth factor mRNA, activates astrocytes, and protects rat brain tissue against ischemic damage. *Eur J Pharmacol*, 1999, 379, 33–45.
- Giulian D, Baker TJ: Characterization of amoeboid microglia isolated from developing mammalian brain. *J Neurosci*, 1986, 6, 2163–2178.
- Hertz L, Chen Y, Gibbs ME, Zang P, Peng L: Astrocytic adrenoceptors: a major drug target in neurological and psychiatric disorders? *Curr Drug Targets CNS Neurol Disord*, 2004, 3, 239–267.
- Junker V, Becker A, Hühne R, Zembatov M, Ravati A, Culmsee C, Kriegstein J: Stimulation of β -adrenoceptors activates astrocytes and provides neuroprotection. *Eur J Pharmacol*, 2002, 446, 25–36.
- Kocki T, Luchowski P, Luchowska E, Wielosz M, Turski WA, Urbańska EM: L-cysteine sulphinate, endogenous sulphur-containing amino acid, inhibits rat brain kynurenic acid production via selective interference with kynurenine aminotransferase II. *Neurosci Lett*, 2003, 346, 97–100.
- Kocki T, Wielosz M, Turski WA, Urbańska EM: Enhancement of brain kynurenic acid production by anticonvulsants—novel mechanism of antiepileptic activity? *Eur J Pharmacol*, 2006, 41, 147–151.
- Kuc D, Rahnama M, Tomaszewski T, Rzeski W, Wejksza K, Urbanik-Sypniewska T, Parada-Turska J et al.: Kynurenic acid in human saliva – does it influence oral microflora? *Pharmacol Rep*, 2006, 58, 393–398.
- Lenard NR, Gettys TW, Dunn AJ: Activation of β_2 - and β_3 -adrenergic receptors increases brain tryptophan. *J Pharmacol Exp Ther*, 2003, 305, 653–659.
- Luccini E, Musante V, Neri E, Raiteri M, Pittaluga A: N-methyl-D-aspartate autoreceptors respond to low and high agonist concentrations by facilitating, respectively, exocytosis and carrier-mediated release of glutamate in rat hippocampus. *J Neurosci Res*, 2007, 85, 3657–3665.
- Luchowska E, Luchowski P, Paczek R, Ziembowicz A, Kocki T, Turski WA, Wielosz M et al.: Dual effect of DL-homocysteine and S-adenosylhomocysteine on brain synthesis of the glutamate receptor antagonist, kynurenic acid. *J Neurosci Res*, 2005, 79, 375–382.
- Marien MR, Colpaert FC, Rosenquist AC: Noradrenergic mechanisms in neurodegenerative diseases: a theory. *Brain Res Brain Res Rev*, 2004, 45, 38–78.
- Németh H, Toldi J, Vecsei L: Role of kynurenines in the central and peripheral nervous systems. *Curr Neurovasc Res*, 2005, 2, 249–260.
- Semkova I, Schilling M, Henrich-Noack P, Rami A, Kriegstein J: Clenbuterol protects mouse cerebral cortex and rat hippocampus from ischemic damage and attenuates glutamate neurotoxicity in cultured hippocampal neurons by induction of NGF. *Brain Res*, 1996, 717, 44–54.
- Stazka J, Luchowski P, Wielosz M, Kleinrok Z, Urbańska EM: Endothelium-dependent production and liberation of kynurenic acid by rat aortic rings exposed to L-kynurenine. *Eur J Pharmacol*, 2002, 448, 133–137.
- Urbańska EM, Luchowski P, Luchowska E, Pniewski J, Woźniak R, Chodakowska-Żebrowska M, Łazarewicz J: Serum kynurenic acid positively correlates with cardiovascular disease risk factor, homocysteine a study in stroke patients. *Pharmacol Rep*, 2006, 58, 501–511.
- Vincent VAM, Van Dam A-M, Persoons JHA, Schotanus K, Steinbusch HWN, Schoffeleer ANM, Berkenbosch F: Gradual inhibition of inducible nitric oxide synthase but not interleukin-1 β production in rat microglial cells of endotoxin-treated mixed glial cell cultures. *Glia*, 1996, 17, 94–102.

Received:

February 27, 2008; in revised form: May 13, 2008.